

MEASURED QUANTUM GROUPOIDS IN ACTION

MICHEL ENOCK

ABSTRACT. Franck Lesieur had introduced in his thesis (now published in an expended and revised version in the *Mémoires de la SMF* (2007)) a notion of measured quantum groupoid, in the setting of von Neumann algebras and a simplification of Lesieur's axioms is presented in an appendix of this article. We here develop the notions of actions, crossed-product, and obtain a biduality theorem, following what had been done by Stefaan Vaes for locally compact quantum groups. Moreover, we prove that the inclusion of the initial algebra into its crossed-product is depth 2, which gives a converse of a result proved by Jean-Michel Vallin and the author. More precisely, to any action of a measured quantum groupoid, we associate another measured quantum groupoid. In particular, starting from an action of a locally compact quantum group, we obtain a measured quantum groupoid canonically associated to this action; when the action is outer, this measured quantum groupoid is the initial locally compact quantum group.

arXiv:0710.5364v2 [math.OA] 19 Sep 2008

1991 *Mathematics Subject Classification.* 46L55, 46L89.

Key words and phrases. measured quantum groupoids, actions, crossed-product, biduality theorem, depth 2 inclusions.

1. Introduction	6
1.1.	6
1.2.	6
1.3.	6
1.4.	6
1.5.	6
1.6.	7
2. Preliminaries	7
2.1. Spatial theory [C1], [S2], [T]	8
2.2. Jones' basic construction and operator-valued weights; depth 2 inclusions	10
2.3. Relative tensor product [C1], [S2], [T]	12
2.4. Fiber product [V1], [EV]	15
2.5. Slice maps [E4]	16
2.6. Vaes' Radon-Nikodym theorem	18
3. Measured quantum groupoids	18
3.1. Definition	18
3.2. Definition	19
3.3. Algebras and Hopf-bimodules associated to a pseudo-multiplicative unitary	20
3.4. Fundamental example	22
3.5. Definitions ([L1], [L2])	22
3.6. Theorem([L1], [L2])	23
3.7. Definitions	23
3.8. Theorem ([L2], [E6])	23
3.9. Notations	25
3.10. Theorem ([L2])	25
3.11. Theorem ([L2])	27
3.12. Theorem([L2])	27
3.13. Example	28
3.14. Example	29
3.15. Theorem([EV], [E4] 8.2 and 8.3)	29
4. Left invariance revisited	30
4.1. Definitions and notations	30
4.2. Lemma	31
4.3. Lemma	31
4.4. Definitions and lemma	31
4.5. Proposition	32
4.6. Proposition	33
4.7. Proposition	34
4.8. Theorem	35
4.9. Proposition	35
4.10. Proposition	36
4.11. Proposition	36
4.12. Theorem	39
5. Corepresentations of measured quantum groupoids	39
5.1. Definition	39
5.2. Theorem	40
5.3. Corollary	40

5.4.	Proposition	41
5.5.	Proposition	42
5.6.	Example	42
5.7.	Proposition	43
5.8.	Proposition	43
5.9.	Proposition	44
5.10.	Theorem	44
5.11.	Theorem	45
6.	Actions of measured quantum groupoids	45
6.1.	Definition	45
6.2.	Example	46
6.3.	Example	46
6.4.	Example	46
6.5.	Example	46
6.6.	Proposition	46
6.7.	Theorem	47
6.8.	Proposition	47
6.9.	Definition	48
6.10.	Example	48
6.11.	Definition	48
6.12.	Proposition	48
6.13.	Proposition	48
6.14.	Definition	49
7.	Some technical properties of actions	49
7.1.	Definition	49
7.2.	Proposition	49
7.3.	Definition	50
7.4.	Lemma	50
7.5.	Lemma	50
7.6.	Proposition	51
7.7.	Proposition	52
7.8.	Corollary	52
8.	The standard implementation of an action : the case of a δ -invariant weight	52
8.1.	Definition	52
8.2.	Example	53
8.3.	Lemma	53
8.4.	Proposition	54
8.5.	Proposition	56
8.6.	Theorem	58
8.7.	Lemma	58
8.8.	Theorem	59
8.9.	Corollary	61
8.10.	Corollary	62
8.11.	Corollary	62
9.	Crossed-product and dual actions	63
9.1.	Definition	63
9.2.	Example	63
9.3.	Example	63
9.4.	Theorem	63

9.5. Example	64
9.6. Example	64
9.7. Proposition	64
9.8. Theorem	65
9.9. Definition	65
9.10. Lemma	65
9.11. Proposition	66
9.12. Lemma	66
10. An auxilliary weight on the crossed-product	68
10.1. Proposition	68
10.2. Proposition	68
10.3. Proposition	70
10.4. Lemma	70
10.5. Proposition	71
10.6. Corollary	72
10.7. Theorem	72
10.8. Theorem	73
10.9. Proposition	73
10.10. Definition	73
10.11. Proposition	73
10.12. Corollary	74
11. Biduality	74
11.1. Lemma	74
11.2. Proposition	75
11.3. Proposition	76
11.4. Proposition	77
11.5. Theorem	78
11.6. Theorem	79
11.7. Theorem	79
11.8. Theorem	80
11.9. Theorem	80
11.10. Remark	80
12. Characterization of crossed-products	80
12.1. Notations	81
12.2. Lemma	81
12.3. Theorem	83
12.4. Corollary	84
12.5. Corollary	85
13. Dual weight; bidual weight; depth 2 inclusion associated to an action	85
13.1. Definition	85
13.2. Example	85
13.3. Theorem	86
13.4. Theorem	86
13.5. Proposition	86
13.6. Lemma	87
13.7. Theorem	88
13.8. Theorem	89
13.9. Theorem	90
13.10. Theorem	91

13.11. Remark	91
14. The measured quantum groupoid associated to an action	91
14.1. Theorem	92
14.2. Theorem	92
14.3. Theorem	92
14.4. Theorem	93
14.5. Corollary	94
14.6. Corollary	94
14.7. Proposition	94
14.8. Example	95
14.9. Remark	95
Appendix A. Coinverse and scaling group of a measured quantum groupoid	96
A.1. Lemma	96
A.2. Definition	97
A.3. Proposition	97
A.4. Proposition	97
A.5. Theorem	97
A.6. Theorem	98
A.7. Theorem	98
A.8. Definition	98
A.9. Theorem	99
A.10. Proposition	99
Appendix B. Automorphism groups on the basis of a measured quantum groupoid	100
B.1. Lemma	100
B.2. Proposition	100
B.3. Proposition	101
B.4. Corollary	102
B.5. Lemma	102
B.6. Lemma	103
B.7. Proposition	104
B.8. Theorem	107
References	108

1. INTRODUCTION

1.1. In two articles ([Val2], [Val3]), J.-M. Vallin has introduced two notions (pseudo-multiplicative unitary, Hopf-bimodule), in order to generalize, up to the groupoid case, the classical notions of multiplicative unitary [BS] and of Hopf-von Neumann algebras [ES2] which were introduced to describe and explain duality of groups, and led to appropriate notions of quantum groups ([ES2], [W1], [W2], [BS], [MN], [W3], [KV1], [KV2], [MNW]).

In another article [EV], J.-M. Vallin and the author have constructed, from a depth 2 inclusion of von Neumann algebras $M_0 \subset M_1$, with an operator-valued weight T_1 verifying a regularity condition, a pseudo-multiplicative unitary, which led to two structures of Hopf bimodules, dual to each other. Moreover, we have then constructed an action of one of these structures on the algebra M_1 such that M_0 is the fixed point subalgebra, the algebra M_2 given by the basic construction being then isomorphic to the crossed-product. We construct on M_2 an action of the other structure, which can be considered as the dual action.

If the inclusion $M_0 \subset M_1$ is irreducible, we recovered quantum groups, as proved and studied in former papers ([EN], [E2]).

Therefore, this construction leads to a notion of "quantum groupoid", and a construction of a duality within "quantum groupoids".

1.2. In a finite-dimensional setting, this construction can be mostly simplified, and is studied in [NV1], [BSz1], [BSz2], [Sz], [Val4], [Val5], and examples are described. In [NV2], the link between these "finite quantum groupoids" and depth 2 inclusions of II_1 factors is given.

1.3. Franck Lesieur introduced [L1] a notion of "measured quantum groupoids", in which a modular hypothesis on the basis is required. Mimicking in a wider setting the technics of Kustermans and Vaes ([KV1], [KV2]), he obtained then a pseudo-multiplicative unitary, which, as in the locally compact quantum group case, "contains" all the information of the object (the von Neuman algebra, the coproduct, the antipod, the co-inverse). Unfortunately, the axioms chosen then by Lesieur don't fit perfectly with the duality (namely, the dual object does not fit the modular condition on the basis chosen in [L1]), and, in order to get a perfect symmetry between his objects and their duals, Lesieur gave the name of "measured quantum groupoids" to a wider class [L2], whose axioms could be described as the analog of [MNW], in which a duality is defined and studied, the initial objects considered in [L1] being denoted now "adapted measured quantum groupoids". In [E3] had been shown that, with suitable conditions, the objects constructed in [EV] from depth 2 inclusions, are "measured quantum groupoids" in this new setting.

1.4. Unfortunately, the axioms given in [L2] are very complicated, and there was a serious need for simplification. This has been done in [E6], which is given in an appendix of this article.

1.5. Here are developped the notion of action (already introduced in [EV]), crossed-product, etc, following what had been done for locally compact quantum groups in [E1], [ES1], [V2]. Then we prove a Takesaki-like biduality theorem; moreover, we prove that the inclusion of the initial algebra into its crossed-product is depth 2, and, therefore, using ([EV], [E3]), we can construct another measured quantum groupoid, canonically associated to the action. These results generalize both the case of locally compact quantum groups ([V2]) and the case of measured groupoids ([Y1], [Y2], [Y3]).

1.6. This article is organized as follows :

In chapter 2 are recalled all the definitions and constructions needed for that theory, namely Connes-Sauvageot's relative tensor product of Hilbert spaces, fiber product of von Neumann algebras, depth 2 inclusions, and Vaes' Radon-Nykodym theorem.

The chapter 3 is a résumé of the construction of measured quantum groupoids; this theory is developped using the axiomatization given in [E6]; the equivalence with Lesieur's axioms given in [L2] is postponed in an appendix.

Chapter 4 is a technical part, mostly inspired by [KV2], which will be needed in chapter 6.

In chapter 5 is introduced the notion of (unitary) corepresentation of a measured quantum groupoid; this notion will be used throughout the paper.

In chapter 6 is defined the notion of an action of a measured quantum groupoid on a von Neumann algebra; we define, using chapter 4, the notion of integrable action.

In chapter 7, we define some technical properties of actions (which will be proved to be always satisfied in chapter 11).

Chapter 8 is very technical; we prove, when there exists a δ -invariant weight, that there is a standard implementation of an action.

In chapter 9 is defined the crossed-product of a von Neumann algebra by a measured quantum groupoid, via an action, and the dual action of the dual measured quantum groupoid on this crossed-product.

In chapter 10, we construct on the crossed-product an auxilliary weight which will satisfy all the properties needed for applying chapter 8. Therefore, we get that the dual action has a standard implementation. This auxilliary weight will be used in chapter 13 to define the dual weight.

In chapter 11, we obtain a Takesaki-like biduality theorem : the double crossed-product is, in a certain sense, equivalent to the initial algebra, and the bidual action to the initial action.

In chapter 12, a measured quantum groupoid \mathfrak{G} be given, we characterize the crossed products by an action of \mathfrak{G} among von Neumann algebras on which acts the measured quantum groupoid $\widehat{\mathfrak{G}}^c$.

In chapter 13, using the weight defined in chapter 10 and duality theorems proved in chapter 11, we define the dual weight on the crossed-product, and obtain some results on this weight, and then on the bidual weight, which allow us to study the basic construction made from the inclusion of the initial algebra into its crossed-product, and, using chapter 12, to prove that this inclusion is depth 2.

In chapter 14, we apply the results of ([EV], [E4]) to the depth 2 inclusion obtained in chapter 13, which allow us to construct a new measured quantum groupoid. In particular, starting with an action of a locally compact quantum group, we obtain a measured quantum groupoid; this gives a new example of a measured quantum groupoid which should look familiar to specialists of group actions and groupoids.

In the appendix had been postponed the proof that our axioms, developped in [E6], are equivalent with Lesieur's axioms, given in [L2]. Namely, in appendix A, with our axioms, we construct the co-inverse, the scaling group and the antipod, mimicking what Lesieur did for his "adapted measured quantum groupoids". Then, in appendix B, we prove the equivalence of the two sets of axioms.

2. PRELIMINARIES

In this chapter are mainly recalled definitions and notations about Connes' spatial theory (2.1, 2.3) and the fiber product construction (2.4, 2.5) which are the main technical

tools of the theory of measured quantum theory. We recall also some definitions and results about depth 2 inclusions (2.2), and Vaes' Radon-Nykodym theorem (2.6).

2.1. Spatial theory [C1], [S2], [T]. Let N be a von Neumann algebra, and let ψ be a faithful semi-finite normal weight on N ; let \mathfrak{N}_ψ , \mathfrak{M}_ψ , H_ψ , π_ψ , Λ_ψ , J_ψ , Δ_ψ , ... be the canonical objects of the Tomita-Takesaki construction associated to the weight ψ . Let α be a non-degenerate normal representation of N on a Hilbert space \mathcal{H} . We may as well consider \mathcal{H} as a left N -module, and write it then ${}_\alpha\mathcal{H}$. Following ([C1], definition 1), we define the set of ψ -bounded elements of ${}_\alpha\mathcal{H}$ as :

$$D({}_\alpha\mathcal{H}, \psi) = \{\xi \in \mathcal{H}; \exists C < \infty, \|\alpha(y)\xi\| \leq C\|\Lambda_\psi(y)\|, \forall y \in \mathfrak{N}_\psi\}$$

Then, for any ξ in $D({}_\alpha\mathcal{H}, \psi)$, there exists a bounded operator $R^{\alpha, \psi}(\xi)$ from H_ψ to \mathcal{H} , defined, for all y in \mathfrak{N}_ψ by :

$$R^{\alpha, \psi}(\xi)\Lambda_\psi(y) = \alpha(y)\xi$$

If there is no ambiguity about the representation α , we shall write $R^\psi(\xi)$ instead of $R^{\alpha, \psi}(\xi)$. This operator belongs to $Hom_N(H_\psi, \mathcal{H})$; therefore, for any ξ, η in $D({}_\alpha\mathcal{H}, \psi)$, the operator :

$$\theta^{\alpha, \psi}(\xi, \eta) = R^{\alpha, \psi}(\xi)R^{\alpha, \psi}(\eta)^*$$

belongs to $\alpha(N)'$; moreover, $D({}_\alpha\mathcal{H}, \psi)$ is dense ([C1], lemma 2), stable under $\alpha(N)'$, and the linear span generated by the operators $\theta^{\alpha, \psi}(\xi, \eta)$ is a weakly dense ideal in $\alpha(N)'$.

With the same hypothesis, the operator :

$$\langle \xi, \eta \rangle_{\alpha, \psi} = R^{\alpha, \psi}(\eta)^* R^{\alpha, \psi}(\xi)$$

belongs to $\pi_\psi(N)'$. Using Tomita-Takesaki's theory, $\pi_\psi(N)'$ is equal to $J_\psi\pi_\psi(N)J_\psi$, and therefore anti-isomorphic to N (or isomorphic to the opposite von Neumann algebra N^o). We shall consider now $\langle \xi, \eta \rangle_{\alpha, \psi}$ as an element of N^o , and the linear span generated by these operators is a dense algebra in N^o . More precisely ([C1], lemma 4, and [S1], lemme 1.5), we get that $\langle \xi, \eta \rangle_{\alpha, \psi}^o$ belongs to \mathfrak{M}_ψ , and that :

$$\Lambda_\psi(\langle \xi, \eta \rangle_{\alpha, \psi}^o) = J_\psi R^{\alpha, \psi}(\xi)^* \eta$$

If y in N is analytical with respect to ψ , and if $\xi \in D({}_\alpha\mathcal{H}, \psi)$, then we get that $\alpha(y)\xi$ belongs to $D({}_\alpha\mathcal{H}, \psi)$ and that :

$$R^{\alpha, \psi}(\alpha(y)\xi) = R^{\alpha, \psi}(\xi)J_\psi\sigma_{-i/2}^\psi(y^*)J_\psi$$

So, if η is another ψ -bounded element of ${}_\alpha\mathcal{H}$, we get :

$$\langle \alpha(y)\xi, \eta \rangle_{\alpha, \psi}^o = \sigma_{i/2}^\psi(y) \langle \xi, \eta \rangle_{\alpha, \psi}^o$$

There exists ([C1], prop.3) a family $(e_i)_{i \in I}$ of ψ -bounded elements of ${}_\alpha\mathcal{H}$, such that

$$\sum_i \theta^{\alpha, \psi}(e_i, e_i) = 1$$

Such a family will be called an (α, ψ) -basis of \mathcal{H} .

It is possible ([EN] 2.2) to construct an (α, ψ) -basis of \mathcal{H} , $(e_i)_{i \in I}$, such that the operators $R^{\alpha, \psi}(e_i)$ are partial isometries with final supports $\theta^{\alpha, \psi}(e_i, e_i)$ 2 by 2 orthogonal, and such that, if $i \neq j$, then $\langle e_i, e_j \rangle_{\alpha, \psi} = 0$. Such a family will be called an (α, ψ) -orthogonal basis of \mathcal{H} . We have then, for ξ, η in $D({}_\alpha\mathcal{H}, \psi)$:

$$R^{\alpha, \psi}(\xi) = \sum_i \theta^{\alpha, \psi}(e_i, e_i) R^{\alpha, \psi}(\xi) = \sum_i R^{\alpha, \psi}(e_i) \langle \xi, e_i \rangle_{\alpha, \psi}$$

$$\langle \xi, \eta \rangle_{\alpha, \psi} = \sum_i \langle \eta, e_i \rangle_{\alpha, \psi}^* \langle \xi, e_i \rangle_{\alpha, \psi}$$

$$\xi = \sum_i R^{\alpha, \psi}(e_i) J_\psi \Lambda_\psi(\langle \xi, e_i \rangle_{\alpha, \psi}^o)$$

the sums being weakly convergent.

Moreover, we get that, for all n in N , $\theta^{\alpha, \psi}(e_i, e_i) \alpha(n) e_i = \alpha(n) e_i$, and $\theta^{\alpha, \psi}(e_i, e_i)$ is the orthogonal projection on the closure of the subspace $\{\alpha(n) e_i, n \in N\}$.

Let β be a normal non-degenerate anti-representation of N on \mathcal{H} . We may then as well consider \mathcal{H} as a right N -module, and write it \mathcal{H}_β , or consider β as a normal non-degenerate representation of the opposite von Neumann algebra N^o , and consider \mathcal{H} as a left N^o -module.

We can then define on N^o the opposite faithful semi-finite normal weight ψ^o ; we have $\mathfrak{N}_{\psi^o} = \mathfrak{N}_\psi^*$, and the Hilbert space H_{ψ^o} will be, as usual, identified with H_ψ , by the identification, for all x in \mathfrak{N}_ψ , of $\Lambda_{\psi^o}(x^*)$ with $J_\psi \Lambda_\psi(x)$.

From these remarks, we infer that the set of ψ^o -bounded elements of \mathcal{H}_β is :

$$D(\mathcal{H}_\beta, \psi^o) = \{\xi \in \mathcal{H}; \exists C < \infty, \|\beta(y^*)\xi\| \leq C \|\Lambda_\psi(y)\|, \forall y \in \mathfrak{N}_\psi\}$$

and, for any ξ in $D(\mathcal{H}_\beta, \psi^o)$ and y in \mathfrak{N}_ψ , the bounded operator $R^{\beta, \psi^o}(\xi)$ is given by the formula :

$$R^{\beta, \psi^o}(\xi) J_\psi \Lambda_\psi(y) = \beta(y^*) \xi$$

This operator belongs to $\text{Hom}_{N^o}(H_\psi, \mathcal{H})$. Moreover, $D(\mathcal{H}_\beta, \psi^o)$ is dense, stable under $\beta(N)' = P$, and, for all y in P , we have :

$$R^{\beta, \psi^o}(y\xi) = y R^{\beta, \psi^o}(\xi)$$

Then, for any ξ, η in $D(\mathcal{H}_\beta, \psi^o)$, the operator

$$\theta^{\beta, \psi^o}(\xi, \eta) = R^{\beta, \psi^o}(\xi) R^{\beta, \psi^o}(\eta)^*$$

belongs to P , and the linear span generated by these operators is a dense ideal in P ; moreover, the operator-valued product $\langle \xi, \eta \rangle_{\beta, \psi^o} = R^{\beta, \psi^o}(\eta)^* R^{\beta, \psi^o}(\xi)$ belongs to $\pi_\psi(N)$; we shall consider now, for simplification, that $\langle \xi, \eta \rangle_{\beta, \psi^o}$ belongs to N , and the linear span generated by these operators is a dense algebra in N , stable under multiplication by analytic elements with respect to ψ . More precisely, $\langle \xi, \eta \rangle_{\beta, \psi^o}$ belongs to \mathfrak{M}_ψ ([C1], lemma 4) and we have ([S1], lemme 1.5) :

$$\Lambda_\psi(\langle \xi, \eta \rangle_{\beta, \psi^o}) = R^{\beta, \psi^o}(\eta)^* \xi$$

A (β, ψ^o) -basis of \mathcal{H} is a family $(e_i)_{i \in I}$ of ψ^o -bounded elements of \mathcal{H}_β , such that :

$$\sum_i \theta^{\beta, \psi^o}(e_i, e_i) = 1$$

We have then, for all ξ in $D(\mathcal{H}_\beta, \psi^o)$:

$$\xi = \sum_i R^{\beta, \psi^o}(e_i) \Lambda_\psi(\langle \xi, e_i \rangle_{\beta, \psi^o})$$

It is possible to choose the $(e_i)_{i \in I}$ such that the $R^{\beta, \psi^o}(e_i)$ are partial isometries, with final supports $\theta^{\beta, \psi^o}(e_i, e_i)$ 2 by 2 orthogonal, and $\langle e_i, e_j \rangle_{\beta, \psi^o} = 0$ if $i \neq j$; such a family will be then called a (β, ψ^o) -orthogonal basis of \mathcal{H} . We have then

$$R^{\beta, \psi^o}(e_i) = \theta^{\beta, \psi^o}(e_i, e_i) R^{\beta, \psi^o}(e_i) = R^{\beta, \psi^o}(e_i) \langle e_i, e_i \rangle_{\beta, \psi^o}$$

Moreover, we get that, for all n in N , and for all i , we have $\theta^{\beta, \psi^o}(e_i, e_i)\beta(n)e_i = \beta(n)e_i$, and therefore, $\theta^{\beta, \psi^o}(e_i, e_i)$ is the orthogonal projection on the closure of the subspace $\{\beta(n)e_i, n \in N\}$.

2.2. Jones' basic construction and operator-valued weights; depth 2 inclusions.

Let $M_0 \subset M_1$ be an inclusion of von Neumann algebras (for simplification, these algebras will be supposed to be σ -finite), equipped with a normal faithful semi-finite operator-valued weight T_1 from M_1 to M_0 (to be more precise, from M_1^+ to the extended positive elements of M_0 (cf. [T] IX.4.12)). Let ψ_0 be a normal faithful semi-finite weight on M_0 , and $\psi_1 = \psi_0 \circ T_1$; for $i = 0, 1$, let $H_i = H_{\psi_i}$, $J_i = J_{\psi_i}$, $\Delta_i = \Delta_{\psi_i}$ be the usual objects constructed by the Tomita-Takesaki theory associated to these weights. Following ([J], 3.1.5(i)), the von Neumann algebra $M_2 = J_1 M_0' J_1$ defined on the Hilbert space H_1 will be called the basic construction made from the inclusion $M_0 \subset M_1$. We have $M_1 \subset M_2$, and we shall say that the inclusion $M_0 \subset M_1 \subset M_2$ is standard.

Following ([EN] 10.6), for x in \mathfrak{N}_{T_1} , we shall define $\Lambda_{T_1}(x)$ by the following formula, for all z in \mathfrak{N}_{ψ_0} :

$$\Lambda_{T_1}(x)\Lambda_{\psi_0}(z) = \Lambda_{\psi_1}(xz)$$

This operator belongs to $\text{Hom}_{M_0^o}(H_0, H_1)$; if x, y belong to \mathfrak{N}_{T_1} , then $\Lambda_{T_1}(x)\Lambda_{T_1}(y)^*$ belongs to M_2 , and $\Lambda_{T_1}(x)^*\Lambda_{T_1}(y) = T_1(x^*y) \in M_0$.

Using then Haagerup's construction ([T], IX.4.24), it is possible to construct a normal semi-finite faithful operator-valued weight T_2 from M_2 to M_1 ([EN], 10.7), which will be called the basic construction made from T_1 . If x, y belong to \mathfrak{N}_{T_1} , then $\Lambda_{T_1}(x)\Lambda_{T_1}(y)^*$ belongs to \mathfrak{M}_{T_2} , and $T_2(\Lambda_{T_1}(x)\Lambda_{T_1}(y)^*) = xy^*$. As we have, for all normal semi-finite faithful weight φ on M_1 :

$$\frac{d\varphi}{d\varphi^o} = \Delta_\varphi$$

the operator-valued weight T_2 is characterized by the equality ([EN], 10.3) :

$$\frac{d\psi_1 \circ T_2}{d\psi_0^o} = \frac{d\psi_1}{d(\psi_0 \circ T_1)^o} = \Delta_{\psi_1}$$

The operator-valued weight T_2 from M_2 to M_1 will be called the basic construction made from the operator-valued weight T_1 from M_1 to M_0 .

Repeating this construction, we obtain by recurrence successive basic constructions, which lead to Jones' tower $(M_i)_{i \in \mathbb{N}}$ of von Neumann algebras, which is the inclusion :

$$M_0 \subset M_1 \subset M_2 \subset M_3 \subset M_4 \subset \dots$$

which is equipped (for $i \geq 1$) with normal faithful semi-finite operator-valued weights T_i from M_i to M_{i-1} . We define then, by recurrence, the weight $\psi_i = \psi_{i-1} \circ T_i$ on M_i , and we shall write H_i, J_i, Δ_i instead of H_{ψ_i} , etc. We shall define the mirroring j_i on $\mathcal{L}(H_i)$ by $j_i(x) = J_i x^* J_i$, for all x in $\mathcal{L}(H_i)$.

Following ([EN] 10.6), for x in \mathfrak{N}_{T_i} , we shall define $\Lambda_{T_i}(x)$ by the following formula, for all z in $\mathfrak{N}_{\psi_{i-1}}$:

$$\Lambda_{T_i}(x)\Lambda_{\psi_{i-1}}(z) = \Lambda_{\psi_i}(xz)$$

Then, $\Lambda_{T_i}(x)$ belongs to $\text{Hom}_{M_{i-1}^o}(H_{i-1}, H_i)$; if x, y belong to \mathfrak{N}_{T_i} , then $\Lambda_{T_i}(x)^*\Lambda_{T_i}(y) = T_i(x^*y)$, and $\Lambda_{T_i}(x)\Lambda_{T_i}(y)^*$ belongs to M_{i+1} ; more precisely, it belongs to $\mathfrak{M}_{T_{i+1}}$, and $T_{i+1}(\Lambda_{T_i}(x)\Lambda_{T_i}(y)^*) = xy^*$.

Let $M_0 \subset M_1$ be an inclusion of von Neumann algebras, equipped with a normal semi-finite faithful operator-valued weight T_1 from M_1 to M_0 ; following ([GHJ] 4.6.4), we shall

say that the inclusion $M_0 \subset M_1$ is depth 2 if the inclusion (called the derived tower) :

$$M'_0 \cap M_1 \subset M'_0 \cap M_2 \subset M'_0 \cap M_3$$

is standard, and, following ([EN], 11.12), we shall say that the operator-valued weight T_1 is regular if both restrictions $T_{2|M'_0 \cap M_2}$ and $T_{3|M'_0 \cap M_3}$ are semi-finite.

By Tomita-Takesaki theory, the Hilbert space H_1 bears a natural structure of $M_1 - M_1^o$ -bimodule, and, therefore, by restriction, of $M_0 - M_0^o$ -bimodule. Let us write r for the canonical representation of M_0 on H_1 , and s for the canonical antirepresentation given, for all x in M_0 , by $s(x) = J_1 r(x)^* J_1$. Let us have now a closer look to the subspaces $D(H_{1s}, \psi_0^o)$ and $D({}_r H_1, \psi_0)$. If x belongs to $\mathfrak{N}_{T_1} \cap \mathfrak{N}_{\psi_1}$, we easily get that $J_1 \Lambda_{\psi_1}(x)$ belongs to $D({}_r H_1, \psi_0)$, with :

$$R^{r, \psi_0}(J_1 \Lambda_{\psi_1}(x)) = J_1 \Lambda_{T_1}(x) J_0$$

and $\Lambda_{\psi_1}(x)$ belongs to $D(H_{1s}, \psi_0)$, with :

$$R^{s, \psi_0^o}(\Lambda_{\psi_1}(x)) = \Lambda_{T_1}(x)$$

In ([E4], 2.3) was proved that the subspace $D(H_{1s}, \psi_0^o) \cap D({}_r H_1, \psi_0)$ is dense in H_1 ; let us write down and precise this result :

2.2.1. Proposition. *Let us keep on the notations of this paragraph; let $\mathcal{T}_{\psi_1, T_1}$ be the algebra made of elements x in $\mathfrak{N}_{\psi_1} \cap \mathfrak{N}_{T_1} \cap \mathfrak{N}_{\psi_1}^* \cap \mathfrak{N}_{T_1}^*$, analytical with respect to ψ_1 , and such that, for all z in \mathbb{C} , $\sigma_z^{\psi_1}(x_n)$ belongs to $\mathfrak{N}_{\psi_1} \cap \mathfrak{N}_{T_1} \cap \mathfrak{N}_{\psi_1}^* \cap \mathfrak{N}_{T_1}^*$. Then :*

- (i) *the algebra $\mathcal{T}_{\psi_1, T_1}$ is weakly dense in M_1 ; it will be called Tomita's algebra with respect to ψ_1 and T_1 ;*
- (ii) *for any x in $\mathcal{T}_{\psi_1, T_1}$, $\Lambda_{\psi_1}(x)$ belongs to $D(H_{1s}, \psi_0) \cap D({}_r H_1, \psi_0)$;*
- (iii) *for any ξ in $D(H_{1s}, \psi_0^o)$, there exists a sequence x_n in $\mathcal{T}_{\psi_1, T_1}$ such that $\Lambda_{T_1}(x_n) = R^{s, \psi_0^o}(\Lambda_{\psi_1}(x))$ is weakly converging to $R^{s, \psi_0^o}(\xi)$ and $\Lambda_{\psi_1}(x_n)$ is converging to ξ .*

Proof. The result (i) is taken from ([EN], 10.12); we get in ([E4], 2.3) an increasing sequence of projections p_n in M_1 , converging to 1, and elements x_n in $\mathcal{T}_{\psi_1, T_1}$ such that $\Lambda_{\psi_1}(x_n) = p_n \xi$. So, (i) and (ii) were obtained in ([E4], 2.3) from this construction. More precisely, we get that :

$$\begin{aligned} T_1(x_n^* x_n) &= \langle R^{s, \psi^o}(\Lambda_{\psi_1}(x_n)), R^{s, \psi_0^o}(\Lambda_{\psi_1}(x_n)) \rangle_{s, \psi_0^o} \\ &= \langle p_n \xi, p_n \xi \rangle_{s, \psi_0^o} \\ &= R^{s, \psi^o}(\xi)^* p_n R^{s, \psi^o}(\xi) \end{aligned}$$

which is increasing and weakly converging to $\langle \xi, \xi \rangle_{s, \psi_0^o}$. □

We finish by writing a proof of this useful lemma, we were not able to find in litterature:

2.2.2. Lemma. *Let $M_0 \subset M_1$ be an inclusion of von neumann algebras, equipped with a normal faithful semi-finite operator-valued weight T from M_1 to M_0 . Let ψ_0 be a normal semi-finite faithful weight on M_0 , and $\psi_1 = \psi_0 \circ T$; if x is in \mathfrak{N}_T , and if y is in $M'_0 \cap M_1$, analytical with respect to ψ_1 , then xy belongs to \mathfrak{N}_T .*

Proof. Let a be in \mathfrak{N}_{ψ_0} ; then xa belongs to \mathfrak{N}_{ψ_1} , and $xya = xay$ belongs to \mathfrak{N}_{ψ_1} ; moreover, let us consider the element $T(y^*x^*xy)$ of the positive extended part of M_0^+ ; we have :

$$\begin{aligned}
\langle T(y^*x^*xy), \omega_{\Lambda_{\psi_0}(a)} \rangle &= \psi_1(a^*y^*x^*xya) \\
&= \|\Lambda_{\psi_1}(xay)\|^2 \\
&= \|J_{\psi_1}\sigma_{-i/2}^{\psi_1}(y^*)J_{\psi_1}\Lambda_{\psi_1}(xa)\|^2 \\
&= \|J_{\psi_1}\sigma_{-i/2}^{\psi_1}(y^*)J_{\psi_1}\Lambda_T(x)\Lambda_{\psi_0}(a)\|^2 \\
&\leq \|\sigma_{-i/2}^{\psi_1}(y^*)\|^2 \|\Lambda_T(x)\Lambda_{\psi_0}(a)\|^2 \\
&= \|\sigma_{-i/2}^{\psi_1}(y^*)\|^2 \langle T(x^*x), \omega_{\Lambda_{\psi_0}(a)} \rangle
\end{aligned}$$

from which we get that $T(y^*x^*xy)$ is bounded and

$$T(y^*x^*xy) \leq \|\sigma_{-i/2}^{\psi_1}(y^*)\|^2 T(x^*x)$$

□

2.3. Relative tensor product [C1], [S2], [T]. Using the notations of 2.1, let now \mathcal{K} be another Hilbert space on which there exists a non-degenerate representation γ of N . Following J.-L. Sauvageot ([S2], 2.1), we define the relative tensor product $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ as

the Hilbert space obtained from the algebraic tensor product $D(\mathcal{H}_{\beta}, \psi^o) \odot \mathcal{K}$ equipped with the scalar product defined, for ξ_1, ξ_2 in $D(\mathcal{H}_{\beta}, \psi^o)$, η_1, η_2 in \mathcal{K} , by

$$(\xi_1 \odot \eta_1 | \xi_2 \odot \eta_2) = (\gamma(\langle \xi_1, \xi_2 \rangle_{\beta, \psi^o}) \eta_1 | \eta_2)$$

where we have identified N with $\pi_{\psi}(N)$ to simplify the notations.

The image of $\xi \odot \eta$ in $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ will be denoted by $\xi \otimes_{\psi} \eta$. We shall use intensively this construction; one should bear in mind that, if we start from another faithful semi-finite normal weight ψ' , we get another Hilbert space $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$; there exists an isomorphism

$U_{\beta, \gamma}^{\psi, \psi'}$ from $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ to $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$, which is unique up to some functorial property ([S2],

2.6) (but this isomorphism does not send $\xi \otimes_{\psi} \eta$ on $\xi \otimes_{\psi'} \eta$!).

When no confusion is possible about the representation and the anti-representation, we shall write $\mathcal{H} \otimes_{\psi} \mathcal{K}$ instead of $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$, and $\xi \otimes_{\psi} \eta$ instead of $\xi \otimes_{\beta \otimes_{\gamma} \mathcal{K}}$.

In ([S1] 2.1), the relative tensor product $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ is defined also, if ξ_1, ξ_2 are in \mathcal{H} , η_1, η_2 are in $D(\gamma \mathcal{K}, \psi)$, by the following formula :

$$(\xi_1 \odot \eta_1 | \xi_2 \odot \eta_2) = (\beta(\langle \eta_1, \eta_2 \rangle_{\gamma, \psi}) \xi_1 | \xi_2)$$

which leads to the definition of a relative flip σ_{ψ} which will be an isomorphism from $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ onto $\mathcal{K}_{\gamma \otimes_{\beta} \mathcal{H}}$, defined, for any ξ in $D(\mathcal{H}_{\beta}, \psi^o)$, η in $D(\gamma \mathcal{K}, \psi)$, by :

$$\sigma_{\psi}(\xi \otimes_{\psi} \eta) = \eta \otimes_{\psi^o} \xi$$

This allows us to define a relative flip ς_{ψ} from $\mathcal{L}(\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}})$ to $\mathcal{L}(\mathcal{K}_{\gamma \otimes_{\beta} \mathcal{H}})$ which sends X in $\mathcal{L}(\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}})$ onto $\varsigma_{\psi}(X) = \sigma_{\psi} X \sigma_{\psi}^*$. Starting from another faithful semi-finite normal weight ψ' , we get a von Neumann algebra $\mathcal{L}(\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}})$ which is isomorphic to $\mathcal{L}(\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}})$,

and a von Neumann algebra $\mathcal{L}(\mathcal{K}_{\gamma \otimes_{\beta} \mathcal{H}})$ which is isomorphic to $\mathcal{L}(\mathcal{K}_{\gamma \otimes_{\beta} \mathcal{H}})$; as we get that :

$$\sigma_{\psi'} \circ U_{\beta, \gamma}^{\psi, \psi'} = U_{\gamma, \beta}^{\psi^o, \psi'^o}$$

we see that these isomorphisms exchange ς_{ψ} and $\varsigma_{\psi'}$. Therefore, the homomorphism ς_{ψ} can be denoted ς_N without any reference to a specific weight.

We recall, following ([S2], 2.2b) that, for all ξ in \mathcal{H} , η in $D(\gamma \mathcal{K}, \psi)$, y in N , analytic with respect to ψ , we have :

$$\beta(y) \xi \otimes_{\psi} \eta = \xi \otimes_{\psi} \gamma(\sigma_{-i/2}^{\psi}(y)) \eta$$

With the notations of 2.1, let $(e_i)_{i \in I}$ a (β, ψ^o) -orthogonal basis of \mathcal{H} ; let us remark that, for all η in \mathcal{K} , we have :

$$e_i \otimes_{\psi} \gamma \eta = e_i \otimes_{\psi} \gamma(\langle e_i, e_i \rangle_{\beta, \psi^o}) \eta$$

On the other hand, $\theta^{\beta, \psi^o}(e_i, e_i)$ is an orthogonal projection, and so is $\theta^{\beta, \psi^o}(e_i, e_i) \otimes_{\gamma} 1$; this last operator is the projection on the subspace $e_i \otimes_{\psi} \gamma(\langle e_i, e_i \rangle_{\beta, \psi^o}) \mathcal{K}$ ([E3], 2.3) and, therefore, we get that $\mathcal{H} \otimes_{\psi} \mathcal{K}$ is the orthogonal sum of the subspaces

$$e_i \otimes_{\psi} \gamma(\langle e_i, e_i \rangle_{\beta, \psi^o}) \mathcal{K}$$

For any Ξ in $\mathcal{H} \otimes_{\psi} \mathcal{K}$, there exist ξ_i in \mathcal{K} , such that $\gamma(\langle e_i, e_i \rangle_{\beta, \psi^o}) \xi_i = \xi_i$ and $\Xi = \sum_i e_i \otimes_{\psi} \xi_i$, from which we get that $\sum_i \|\xi_i\|^2 = \|\Xi\|^2$.

Let us suppose now that \mathcal{K} is a $N - P$ bimodule; that means that there exists a von Neumann algebra P , and a non-degenerate normal anti-representation ϵ of P on \mathcal{K} , such that $\epsilon(P) \subset \gamma(N)'$. We shall write then $\gamma \mathcal{K}_{\epsilon}$. If y is in P , we have seen that it is possible to define then the operator $1_{\mathcal{H}} \otimes_{\psi} \epsilon(y)$ on $\mathcal{H} \otimes_{\psi} \mathcal{K}$, and we define this way a non-degenerate normal antirepresentation of P on $\mathcal{H} \otimes_{\psi} \mathcal{K}$, we shall call again ϵ for simplification. If \mathcal{H} is a $Q - N$ bimodule, then $\mathcal{H} \otimes_{\psi} \mathcal{K}$ becomes a $Q - P$ bimodule (Connes' fusion of bimodules).

Taking a faithful semi-finite normal weight ν on P , and a left P -module ${}_{\zeta} \mathcal{L}$ (i.e. a Hilbert space \mathcal{L} and a normal non-degenerate representation ζ of P on \mathcal{L}), it is possible then to define $(\mathcal{H} \otimes_{\psi} \mathcal{K}) \otimes_{\nu} {}_{\zeta} \mathcal{L}$. Of course, it is possible also to consider the Hilbert space

$\mathcal{H} \otimes_{\psi} ({}_{\zeta} \mathcal{L} \otimes_{\nu} \mathcal{K})$. It can be shown that these two Hilbert spaces are isomorphic as

$\beta(N)' - \zeta(P)'^o$ -bimodules. (In ([Val1] 2.1.3), the proof, given for $N = P$ abelian can be used, without modification, in that wider hypothesis). We shall write then $\mathcal{H} \otimes_{\psi} \mathcal{K} \otimes_{\nu} {}_{\zeta} \mathcal{L}$

without parenthesis, to emphasise this coassociativity property of the relative tensor product.

Dealing now with that Hilbert space $\mathcal{H} \otimes_{\psi} \mathcal{K} \otimes_{\nu} {}_{\zeta} \mathcal{L}$, there exist different flips, and it is necessary to be careful with notations. For instance, $1_{\beta} \otimes \sigma_{\nu}$ is the flip from this Hilbert space onto $\mathcal{H} \otimes_{\psi} ({}_{\zeta} \mathcal{L} \otimes_{\nu^o} \mathcal{K})$, where γ is here acting on the second leg of ${}_{\zeta} \mathcal{L} \otimes_{\nu^o} \mathcal{K}$ (and

should therefore be written $1_{\zeta \otimes_{\nu^o} \epsilon} \gamma$, but this will not be done for obvious reasons). Here, the parentheses remain, because there is no associativity rule, and to remind that γ is not acting on \mathcal{L} . The adjoint of $1_{\beta} \otimes \sigma_{\nu}$ is $1_{\beta} \otimes \sigma_{\nu^o}$.

The same way, we can consider $\sigma_{\psi} \epsilon \otimes_{\zeta} 1$ from $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}} \epsilon \otimes_{\zeta} \mathcal{L}$ onto $(\mathcal{H}_{\gamma \otimes_{\beta} \mathcal{H}}) \epsilon \otimes_{\zeta} \mathcal{L}$.

Another kind of flip sends $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{L}} \zeta \otimes_{\epsilon} \mathcal{K}$ onto $\mathcal{L}_{\zeta \otimes_{\epsilon} \mathcal{H}} (\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}})$. We shall denote this

application $\sigma_{\gamma, \epsilon}^{1,2}$ (and its adjoint $\sigma_{\epsilon, \gamma}^{1,2}$), in order to emphasize that we are exchanging the first and the second leg, and the representations γ and ϵ on the third leg.

If π denotes the canonical left representation of N on the Hilbert space $L^2(N)$, we verify that the application which sends, for all ξ in \mathcal{H} , ψ normal semi-finite faithful weight on N , and x in \mathfrak{N}_{ψ} , the vector $\xi_{\beta \otimes_{\pi} J_{\psi} \Lambda_{\psi}(x)}$ on $\beta(x^*)\xi$, gives an isomorphism of $\mathcal{H}_{\beta \otimes_{\pi} L^2(N)}$,

which sends the antirepresentation of N given by $n \mapsto 1_{\mathcal{H}} \beta \otimes_{\pi} L^2(N)$ on β . The same

way, there exists a canonical identification, as left N -modules, of $L^2(N) \otimes_{\psi} \mathcal{K}$ with \mathcal{K} . For any ξ in $D(\mathcal{H}_{\beta}, \psi^o)$, we define the bounded linear application $\lambda_{\xi}^{\beta, \gamma}$ from \mathcal{K} to $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$

by, for all η in \mathcal{K} , $\lambda_{\xi}^{\beta, \gamma}(\eta) = \xi_{\beta \otimes_{\gamma} \eta}$. We shall write λ_{ξ} if no confusion is possible. We get

([EN], 3.10) :

$$\lambda_{\xi}^{\beta, \gamma} = R^{\beta, \psi^o}(\xi) \otimes_{\psi} 1_{\mathcal{K}}$$

We have :

$$(\lambda_{\xi}^{\beta, \gamma})^* \lambda_{\xi}^{\beta, \gamma} = \gamma(< \xi, \xi >_{\beta, \psi^o})$$

We may define, for any η in $D(\gamma \mathcal{K}, \psi)$, an application $\rho_{\eta}^{\beta, \gamma}$ from \mathcal{H} to $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ by

$\rho_{\eta}^{\beta, \gamma}(\xi) = \xi_{\beta \otimes_{\gamma} \eta}$. We shall write ρ_{η} if no confusion is possible. We get that :

$$(\rho_{\eta}^{\beta, \gamma})^* \rho_{\eta}^{\beta, \gamma} = \beta(< \eta, \eta >_{\gamma, \psi})$$

Let x be an element of $\mathcal{L}(\mathcal{H})$, commuting with the right action of N on \mathcal{H}_{β} (i.e. x belongs to $\beta(N)'$). It is possible to define an operator $x_{\beta \otimes_{\gamma} 1_{\mathcal{K}}}$ on $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$. We can easily evaluate $\|x_{\beta \otimes_{\gamma} 1_{\mathcal{K}}}\|$: for any finite $J \subset I$, for any η_i in \mathcal{K} , we have :

$$\begin{aligned} ((x^* x)_{\beta \otimes_{\gamma} 1_{\mathcal{K}}})(\sum_{i \in J} e_i_{\beta \otimes_{\gamma} \eta_i}) &= \sum_{i \in J} (\gamma(< x e_i, x e_i >_{\beta, \psi^o}) \eta_i | \eta_i) \\ &\leq \|x\|^2 \sum_{i \in J} (\gamma(< e_i, e_i >_{\beta, \psi^o}) \eta_i | \eta_i) \\ &= \|x\|^2 \|\sum_{i \in J} e_i_{\beta \otimes_{\gamma} \eta_i}\| \end{aligned}$$

from which we get $\|x_{\beta \otimes_{\gamma} 1_{\mathcal{K}}}\| \leq \|x\|$.

By the same way, if y commutes with the left action of N on $\gamma \mathcal{K}$ (i.e. $y \in \gamma(N)'$), it is possible to define $1_{\mathcal{H}} \beta \otimes_{\gamma} y$ on $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$, and by composition, it is possible to define then

$x_{\beta \otimes_{\gamma} y}$. If we start from another faithful semi-finite normal weight ψ' , the canonical

isomorphism $U_{\beta, \gamma}^{\psi, \psi'}$ from $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ to $\mathcal{H}_{\beta \otimes_{\gamma} \mathcal{K}}$ sends $x_{\beta \otimes_{\gamma} y}$ on $x_{\beta \otimes_{\gamma} y}$ ([S2], 2.3 and

2.6); therefore, this operator can be denoted $x \underset{N}{\beta \otimes_\gamma} y$ without any reference to a specific weight, and we get $\|x \underset{N}{\beta \otimes_\gamma} y\| \leq \|x\| \|y\|$.

If \mathcal{K} is a Hilbert space on which there exists a non-degenerate representation γ of N , then \mathcal{K} is a $N - \gamma(N)'^o$ bimodule, and the conjugate Hilbert space $\overline{\mathcal{K}}$ is a $\gamma(N)' - N^o$ bimodule, and, ([S2]), for any normal faithful semi-finite weight ϕ on $\gamma(N)'$, the fusion ${}_{\gamma(N)'} \mathcal{K} \underset{\phi^o}{\otimes} \overline{\mathcal{K}}$ is isomorphic to the standard space $L^2(N)$, equipped with its standard left and right representation.

Using that remark, the associativity rule, and the identification (as right N -modules) of $\mathcal{H} \underset{\psi}{\otimes} L^2(N)$ with \mathcal{H} , one gets for any $x \in \beta(N)'$:

$$\|x \underset{N}{\beta \otimes_\gamma} 1_{\mathcal{K}}\| \leq \|x \underset{N}{\beta \otimes_\gamma} 1_{\mathcal{K}} \underset{\gamma(N)'^o}{\otimes} 1_{\overline{\mathcal{K}}}\| = \|x \underset{N}{\beta \otimes} 1_{L^2(N)}\| = \|x\|$$

from which we get $\|x \underset{N}{\beta \otimes_\gamma} 1_{\mathcal{K}}\| = \|x\|$.

If \mathcal{H} and \mathcal{K} are finite-dimensional Hilbert spaces, the relative tensor product $\mathcal{H} \underset{\psi}{\beta \otimes_\gamma} \mathcal{K}$ can be identified with a subspace of the tensor Hilbert space $\mathcal{H} \otimes \mathcal{K}$ ([EV] 2.4), the projection on which belonging to $\beta(N) \otimes \gamma(N)$.

2.4. Fiber product [V1], [EV]. Let us follow the notations of 2.3; let now M_1 be a von Neumann algebra on \mathcal{H} , such that $\beta(N) \subset M_1$, and M_2 be a von Neumann algebra on \mathcal{K} , such that $\gamma(N) \subset M_2$. The von Neumann algebra generated by all elements $x \underset{N}{\beta \otimes_\gamma} y$, where x belongs to M_1' , and y belongs to M_2' , will be denoted $M_1' \underset{N}{\beta \otimes_\gamma} M_2'$ (or $M_1' \otimes_N M_2'$ if no confusion if possible), and will be called the relative tensor product of M_1' and M_2' over N . The commutant of this algebra will be denoted $M_1 \underset{N}{\beta * \gamma} M_2$ (or $M_1 *_N M_2$ if no confusion is possible) and called the fiber product of M_1 and M_2 , over N . It is straightforward to verify that, if P_1 and P_2 are two other von Neumann algebras satisfying the same relations with N , we have :

$$M_1 *_N M_2 \cap P_1 *_N P_2 = (M_1 \cap P_1) *_N (M_2 \cap P_2)$$

Moreover, we get that $\varsigma_N(M_1 \underset{N}{\beta * \gamma} M_2) = M_2 \underset{N^o}{\gamma * \beta} M_1$.

In particular, we have :

$$(M_1 \cap \beta(N)') \underset{N}{\beta \otimes_\gamma} (M_2 \cap \gamma(N)') \subset M_1 \underset{N}{\beta * \gamma} M_2$$

and :

$$M_1 \underset{N}{\beta * \gamma} \gamma(N) = (M_1 \cap \beta(N)') \underset{N}{\beta \otimes_\gamma} 1$$

More generally, if β is a non-degenerate normal involutive antihomomorphism from N into a von Neumann algebra M_1 , and γ a non-degenerate normal involutive homomorphism from N into a von Neumann algebra M_2 , it is possible to define, without any reference to a specific Hilbert space, a von Neumann algebra $M_1 \underset{N}{\beta * \gamma} M_2$.

Moreover, if now β' is a non-degenerate normal involutive antihomomorphism from N into another von Neumann algebra P_1 , γ' a non-degenerate normal involutive homomorphism from N into another von Neumann algebra P_2 , Φ a normal involutive homomorphism from M_1 into P_1 such that $\Phi \circ \beta = \beta'$, and Ψ a normal involutive homomorphism from M_2 into P_2 such that $\Psi \circ \gamma = \gamma'$, it is possible then to define a normal involutive homomorphism

(the proof given in ([S1] 1.2.4) in the case when N is abelian can be extended without modification in the general case) :

$$\Phi_{\beta * \gamma} \Psi : M_1 \underset{N}{\beta * \gamma} M_2 \mapsto P_1 \underset{N}{\beta' * \gamma'} P_2$$

In the case when $\gamma \mathcal{K}_\epsilon$ is a $N - P^o$ bimodule as explained in 2.3 and $\zeta \mathcal{L}$ a P -module, if $\gamma(N) \subset M_2$ and $\epsilon(P) \subset M_2$, and if $\zeta(P) \subset M_3$, where M_3 is a von Neumann algebra on \mathcal{L} , it is possible to consider then $(M_1 \underset{N}{\beta * \gamma} M_2) \underset{P}{\epsilon * \zeta} M_3$ and $M_1 \underset{N}{\beta * \gamma} (M_2 \underset{P}{\epsilon * \zeta} M_3)$. The coassociativity property for relative tensor products leads then to the isomorphism of these von Neumann algebra we shall write now $M_1 \underset{N}{\beta * \gamma} M_2 \underset{P}{\epsilon * \zeta} M_3$ without parenthesis.

If M_1 and M_2 are finite-dimensional, the fiber product $M_1 \underset{N}{\beta * \gamma} M_2$ can be identified to a reduced algebra of $M_1 \otimes M_2$ (reduced by a projector which belongs to $\beta(N) \otimes \gamma(N)$) ([EV] 2.4).

2.5. Slice maps [E4]. Let A be in $M_1 \underset{N}{\beta * \gamma} M_2$, ψ a normal faithful semi-finite weight on N , \mathcal{H} an Hilbert space on which M_1 is acting, \mathcal{K} an Hilbert space on which M_2 is acting, and let ξ_1, ξ_2 be in $D(\mathcal{H}_\beta, \psi^o)$; let us define :

$$(\omega_{\xi_1, \xi_2} \underset{\psi}{\beta * \gamma} id)(A) = (\lambda_{\xi_2}^{\beta, \gamma})^* A \lambda_{\xi_1}^{\beta, \gamma}$$

We define this way $(\omega_{\xi_1, \xi_2} \underset{\psi}{\beta * \gamma} id)(A)$ as a bounded operator on \mathcal{K} , which belongs to M_2 , such that :

$$((\omega_{\xi_1, \xi_2} \underset{\psi}{\beta * \gamma} id)(A) \eta_1 | \eta_2) = (A (\xi_1 \underset{\psi}{\beta \otimes \gamma} \eta_1) | \xi_2 \underset{\psi}{\beta \otimes \gamma} \eta_2)$$

One should note that $(\omega_{\xi_1, \xi_2} \underset{\psi}{\beta * \gamma} id)(1) = \gamma(< \xi_1, \xi_2 >_{\beta, \psi^o})$.

Let us define the same way, for any η_1, η_2 in $D(\gamma \mathcal{K}, \psi)$:

$$(id \underset{\psi}{\beta * \gamma} \omega_{\eta_1, \eta_2})(A) = (\rho_{\eta_2}^{\beta, \gamma})^* A \rho_{\eta_1}^{\beta, \gamma}$$

which belongs to M_1 .

We therefore have a Fubini formula for these slice maps : for any ξ_1, ξ_2 in $D(\mathcal{H}_\beta, \psi^o)$, η_1, η_2 in $D(\gamma \mathcal{K}, \psi)$, we have :

$$< (\omega_{\xi_1, \xi_2} \underset{\psi}{\beta * \gamma} id)(A), \omega_{\eta_1, \eta_2} > = < (id \underset{\psi}{\beta * \gamma} \omega_{\eta_1, \eta_2})(A), \omega_{\xi_1, \xi_2} >$$

Let ϕ_1 be a normal semi-finite weight on M_1^+ , and A be a positive element of the fiber product $M_1 \underset{N}{\beta * \gamma} M_2$, then we may define an element of the extended positive part of M_2 , denoted $(\phi_1 \underset{\psi}{\beta * \gamma} id)(A)$, such that, for all η in $D(\gamma L^2(M_2), \psi)$, we have :

$$\|(\phi_1 \underset{\psi}{\beta * \gamma} id)(A)^{1/2} \eta\|^2 = \phi_1(id \underset{\psi}{\beta * \gamma} \omega_\eta)(A)$$

Moreover, then, if ϕ_2 is a normal semi-finite weight on M_2^+ , we have :

$$\phi_2(\phi_1 \underset{\psi}{\beta * \gamma} id)(A) = \phi_1(id \underset{\psi}{\beta * \gamma} \phi_2)(A)$$

and if ω_i be in M_{1*} such that $\phi_1 = \sup_i \omega_i$, we have $(\phi_1 \underset{\psi}{\beta} *_{\gamma} id)(A) = \sup_i (\omega_i \underset{\psi}{\beta} *_{\gamma} id)(A)$.

Let now P_1 be a von Neuman algebra such that :

$$\beta(N) \subset P_1 \subset M_1$$

and let Φ_i ($i = 1, 2$) be a normal faithful semi-finite operator valued weight from M_i to P_i ; for any positive operator A in the fiber product $M_1 \underset{N}{\beta} *_{\gamma} M_2$, there exists an element $(\Phi_1 \underset{N}{\beta} *_{\gamma} id)(A)$ of the extended positive part of $P_1 \underset{N}{\beta} *_{\gamma} M_2$, such that ([E4], 3.5), for all η in $D(\gamma L^2(M_2), \psi)$, and ξ in $D(L^2(P_1)_{\beta}, \psi^o)$, we have :

$$\|(\Phi_1 \underset{N}{\beta} *_{\gamma} id)(A)^{1/2} (\xi \underset{\psi}{\beta} \otimes_{\gamma} \eta)\|^2 = \|\Phi_1(id \underset{\psi}{\beta} *_{\gamma} \omega_{\eta})(A)^{1/2} \xi\|^2$$

If ϕ is a normal semi-finite weight on P , we have :

$$(\phi \circ \Phi_1 \underset{\psi}{\beta} *_{\gamma} id)(A) = (\phi \underset{\psi}{\beta} *_{\gamma} id)(\Phi_1 \underset{N}{\beta} *_{\gamma} id)(A)$$

We define the same way an element $(id \underset{N}{\beta} *_{\gamma} \Phi_2)(A)$ of the extended positive part of $M_1 \underset{N}{\gamma} *_{\beta} P_2$, and we have :

$$(id \underset{N}{\beta} *_{\gamma} \Phi_2)((\Phi_1 \underset{N}{\beta} *_{\gamma} id)(A)) = (\Phi_1 \underset{N}{\beta} *_{\gamma} id)((id \underset{N}{\beta} *_{\gamma} \Phi_2)(A))$$

Let π denotes the canonical left representation of N on the Hilbert space $L^2(N)$; let x be an element of $M_{1\beta} \underset{N}{*} \pi\pi(N)$, which can be identified (2.4) to $M_1 \cap \beta(N)'$, we get that, for e in \mathfrak{N}_{ψ} , we have :

$$(id_{\beta} \underset{\psi}{*} \pi \omega_{J_{\psi} \Lambda_{\psi}(e)})(x) = \beta(ee^*)x$$

Therefore, by increasing limits, we get that $(id_{\beta} \underset{\psi}{*} \pi \psi)$ is the injection of $M_1 \cap \beta(N)'$ into M_1 . More precisely, if x belongs to $M_1 \cap \beta(N)'$, we have :

$$(id_{\beta} \underset{\psi}{*} \pi \psi)(x_{\beta} \otimes_N \pi 1) = x$$

Therefore, if Φ_2 is a normal faithful semi-finite operator-valued weight from M_2 onto $\gamma(N)$, we get that, for all A positive in $M_1 \underset{N}{\beta} *_{\gamma} M_2$, we have :

$$(id_{\beta} \underset{\psi}{*} \gamma \psi \circ \Phi_2)(A)_{\beta} \otimes_N \gamma 1 = (id_{\beta} \underset{\psi}{*} \gamma \Phi_2)(A)$$

With the notations of 2.1, let $(e_i)_{i \in I}$ be a (β, ψ^o) -orthogonal basis of \mathcal{H} ; using the fact (2.3) that, for all η in \mathcal{K} , we have :

$$e_i \underset{\psi}{\beta} \otimes_{\gamma} \eta = e_i \underset{\psi}{\beta} \otimes_{\gamma} \gamma(< e_i, e_i >_{\beta, \psi^o}) \eta$$

we get that, for all X in $M_1 \underset{N}{\beta} *_{\gamma} M_2$, ξ in $D(\mathcal{H}_{\beta}, \psi^o)$, we have

$$(\omega_{\xi, e_i} \underset{\psi}{\beta} *_{\gamma} id)(X) = \gamma(< e_i, e_i >_{\beta, \psi^o}) (\omega_{\xi, e_i} \underset{\psi}{\beta} *_{\gamma} id)(X)$$

2.6. Vaes' Radon-Nikodym theorem. In [V1] is proved a very nice Radon-Nikodym theorem for two normal faithful semi-finite weights on a von Neumann algebra M . If Φ and Ψ are such weights, then are equivalent :

- the two modular automorphism groups σ^Φ and σ^Ψ commute;
- the Connes' derivative $[D\Psi : D\Phi]_t$ is of the form :

$$[D\Psi : D\Phi]_t = \lambda^{it^2/2} \delta^{it}$$

where λ is a non-singular positive operator affiliated to $Z(M)$, and δ is a non-singular positive operator affiliated to M .

It is then easy to verify that $\sigma_t^\Phi(\delta^{is}) = \lambda^{ist} \delta^{is}$, and that

$$[D\Phi \circ \sigma_t^\Psi : D\Phi]_s = \lambda^{ist}$$

$$[D\Psi \circ \sigma_t^\Phi : D\Psi]_s = \lambda^{-ist}$$

Moreover, we have also, for any $x \in M^+$:

$$\Psi(x) = \lim_n \Phi((\delta^{1/2} e_n) x (\delta^{1/2} e_n))$$

where the e_n are self-adjoint elements of M given by the formula :

$$e_n = \frac{2n^2}{\Gamma(1/2)\Gamma(1/4)} \int_{\mathbb{R}^2} e^{-n^2 x^2 - n^4 y^4} \lambda^{ix} \delta^{iy} dx dy$$

The operators e_n are analytic with respect to σ^Φ and such that, for any $z \in \mathbb{C}$, the sequence $\sigma_z^\Phi(e_n)$ is bounded and strongly converges to 1.

In that situation, we shall write $\Psi = \Phi_\delta$ and call δ the modulus of Ψ with respect to Φ ; λ will be called the scaling operator of Ψ with respect to Φ .

Moreover, if $a \in M$ is such that $a\delta^{1/2}$ is bounded and its closure $\overline{a\delta^{1/2}}$ belongs to \mathfrak{N}_Φ , then a belongs to \mathfrak{N}_Ψ . We may then identify $\Lambda_\Psi(a)$ with $\Lambda_\Phi(\overline{a\delta^{1/2}})$, which leads to the identifications of J_Ψ with $\lambda^{i/4} J_\Phi$, and of Δ_Ψ with $J_\Phi \delta^{-1} J_\Phi \Delta_\Phi$.

3. MEASURED QUANTUM GROUPOIDS

In this chapter, we first recall the definition of Hopf-bimodules (3.1). We then give (3.2) the definition of a pseudo-multiplicative unitary, give the fundamental example given by groupoids (3.4), and construct the algebras and the Hopf-bimodules "generated by the left (resp. right) leg" of a pseudo-multiplicative unitary (3.3). We recall the definition of left-(resp. right-) invariant operator-valued weights on a Hopf-bimodule; we then give the definition of a measured quantum groupoid (3.7). We give the essential theorems which found the theory of measured quantum groupoids and its duality (3.8, 3.10, 3.11, 3.12).

3.1. Definition. A quintuplet $(N, M, \alpha, \beta, \Gamma)$ will be called a Hopf-bimodule, following ([Val2], [EV] 6.5), if N, M are von Neumann algebras, α a faithful non-degenerate representation of N into M , β a faithful non-degenerate anti-representation of N into M , with commuting ranges, and Γ an injective involutive homomorphism from M into $M_{\beta^* \alpha} M_N$

such that, for all X in N :

(i) $\Gamma(\beta(X)) = 1_{\beta \otimes \alpha} \beta(X)$

(ii) $\Gamma(\alpha(X)) = \alpha(X)_{\beta \otimes \alpha} 1_N$

(iii) Γ satisfies the co-associativity relation :

$$(\Gamma_{\beta^* \alpha} id) \Gamma = (id_{\beta^* \alpha} \Gamma) \Gamma$$

This last formula makes sense, thanks to the two preceding ones and 2.4. The von Neumann algebra N will be called the basis of $(N, M, \alpha, \beta, \Gamma)$.

If $(N, M, \alpha, \beta, \Gamma)$ is a Hopf-bimodule, it is clear that $(N^\circ, M, \beta, \alpha, \varsigma_N \circ \Gamma)$ is another Hopf-bimodule, we shall call the symmetrized of the first one. (Recall that $\varsigma_N \circ \Gamma$ is a homomorphism from M to $M_{r^*s} M$).

If N is abelian, $\alpha = \beta$, $\Gamma = \varsigma_N \circ \Gamma$, then the quadruplet $(N, M, \alpha, \alpha, \Gamma)$ is equal to its symmetrized Hopf-bimodule, and we shall say that it is a symmetric Hopf-bimodule.

Let \mathcal{G} be a measured groupoid, with $\mathcal{G}^{(0)}$ as its set of units, and let us denote by r and s the range and source applications from \mathcal{G} to $\mathcal{G}^{(0)}$, given by $xx^{-1} = r(x)$ and $x^{-1}x = s(x)$. As usual, we shall denote by $\mathcal{G}^{(2)}$ (or $\mathcal{G}_{s,r}^{(2)}$) the set of composable elements, i.e.

$$\mathcal{G}^{(2)} = \{(x, y) \in \mathcal{G}^2; s(x) = r(y)\}$$

Let $(\lambda^u)_{u \in \mathcal{G}^{(0)}}$ be a Haar system on \mathcal{G} and ν a measure ν on $\mathcal{G}^{(0)}$. Let us write μ the measure on \mathcal{G} given by integrating λ^u by ν :

$$\mu = \int_{\mathcal{G}^{(0)}} \lambda^u d\nu$$

By definition, ν is said quasi-invariant if μ is equivalent to its image under the inverse $x \mapsto x^{-1}$ of \mathcal{G} (see [Ra], [R1], [R2], [C2] II.5, [P] and [AR] for more details, precise definitions and examples of groupoids).

In [Y1], [Y2], [Y3] and [Val2] were associated to a measured groupoid \mathcal{G} , equipped with a Haar system $(\lambda^u)_{u \in \mathcal{G}^{(0)}}$ and a quasi-invariant measure ν on $\mathcal{G}^{(0)}$ two Hopf-bimodules : The first one is $(L^\infty(\mathcal{G}^{(0)}, \nu), L^\infty(\mathcal{G}, \mu), r_{\mathcal{G}}, s_{\mathcal{G}}, \Gamma_{\mathcal{G}})$, where we define $r_{\mathcal{G}}$ and $s_{\mathcal{G}}$ by writing , for g in $L^\infty(\mathcal{G}^{(0)})$:

$$\begin{aligned} r_{\mathcal{G}}(g) &= g \circ r \\ s_{\mathcal{G}}(g) &= g \circ s \end{aligned}$$

and where $\Gamma_{\mathcal{G}}(f)$, for f in $L^\infty(\mathcal{G})$, is the function defined on $\mathcal{G}^{(2)}$ by $(s, t) \mapsto f(st)$; $\Gamma_{\mathcal{G}}$ is then an involutive homomorphism from $L^\infty(\mathcal{G})$ into $L^\infty(\mathcal{G}_{s,r}^2)$ (which can be identified to $L^\infty(\mathcal{G})_{s^*r} L^\infty(\mathcal{G})$).

The second one is symmetric; it is $(L^\infty(\mathcal{G}^{(0)}, \nu), \mathcal{L}(\mathcal{G}), r_{\mathcal{G}}, r_{\mathcal{G}}, \widehat{\Gamma}_{\mathcal{G}})$, where $\mathcal{L}(\mathcal{G})$ is the von Neumann algebra generated by the convolution algebra associated to the groupoid \mathcal{G} , and $\widehat{\Gamma}_{\mathcal{G}}$ has been defined in [Y3] and [Val2].

If (N, M, r, s, Γ) be a Hopf-bimodule with a finite-dimensional algebra M , then, the identification of $M_{\beta^* \alpha} M$ with a reduced algebra $(M \otimes M)_e$ (2.4) leads to an injective homomorphism $\tilde{\Gamma}$ from M to $M \otimes M$ such that $\tilde{\Gamma}(1) = e \neq 1$ and $(\tilde{\Gamma} \otimes id)\tilde{\Gamma} = (id \otimes \tilde{\Gamma})\tilde{\Gamma}$ ([EV] 6.5). Then $(M, \tilde{\Gamma})$ is a weak Hopf \mathbb{C}^* -algebra in the sense of ([BSz1], [BSz2], [Sz]).

3.2. Definition. Let N be a von Neumann algebra; let \mathfrak{H} be a Hilbert space on which N has a non-degenerate normal representation α and two non-degenerate normal anti-representations $\hat{\beta}$ and β . These 3 applications are supposed to be injective, and to commute two by two. Let ν be a normal semi-finite faithful weight on N ; we can therefore construct the Hilbert spaces $\mathfrak{H}_{\beta \otimes \alpha} \mathfrak{H}$ and $\mathfrak{H}_{\alpha \otimes \hat{\beta}} \mathfrak{H}$. A unitary W from $\mathfrak{H}_{\beta \otimes \alpha} \mathfrak{H}$ onto $\mathfrak{H}_{\alpha \otimes \hat{\beta}} \mathfrak{H}$ will be called a pseudo-multiplicative unitary over the basis N , with respect

to the representation α , and the anti-representations $\hat{\beta}$ and β (we shall write it is an $(\alpha, \hat{\beta}, \beta)$ -pseudo-multiplicative unitary), if :

(i) W intertwines $\alpha, \hat{\beta}, \beta$ in the following way : we have, for all $X \in N$:

$$W(\alpha(X)_{\beta \otimes \alpha} 1) = (1_{\alpha \otimes \hat{\beta}} \alpha(X))W$$

$$W(1_{\beta \otimes \alpha} \beta(X)) = (1_{\alpha \otimes \hat{\beta}} \beta(X))W$$

$$W(\hat{\beta}(X)_{\beta \otimes \alpha} 1) = (\hat{\beta}(X)_{\alpha \otimes \hat{\beta}} 1)W$$

$$W(1_{\beta \otimes \alpha} \hat{\beta}(X)) = (\beta(X)_{\alpha \otimes \hat{\beta}} 1)W$$

(ii) The operator W satisfies :

$$(1_{\mathfrak{H}} \alpha \otimes_{\beta} W)(W_{\beta \otimes \alpha} 1_{\mathfrak{H}}) = (W_{\alpha \otimes \hat{\beta}} 1_{\mathfrak{H}}) \sigma_{\alpha, \beta}^{2,3} (W_{\hat{\beta} \otimes \alpha} 1)(1_{\mathfrak{H}} \beta \otimes_{\alpha} \sigma_{\nu^o})(1_{\mathfrak{H}} \beta \otimes_{\alpha} W)$$

Here, $\sigma_{\alpha, \beta}^{2,3}$ goes from $(H_{\alpha \otimes \hat{\beta}} H)_{\beta \otimes \alpha} H$ to $(H_{\beta \otimes \alpha} H)_{\alpha \otimes \hat{\beta}} H$, and $1_{\mathfrak{H}} \beta \otimes_{\alpha} \sigma_{\nu^o}$ goes from $H_{\beta \otimes \alpha} (H_{\alpha \otimes \hat{\beta}} H)$ to $H_{\beta \otimes \alpha} H_{\hat{\beta} \otimes \alpha} H$.

All the properties supposed in (i) allow us to write such a formula, which will be called the "pentagonal relation".

One should note that this definition is different from the definition introduced in [EV] (and repeated afterwards). It is in fact the same formula, the new writing :

$$\sigma_{\alpha, \beta}^{2,3} (W_{\hat{\beta} \otimes \alpha} 1)(1_{\mathfrak{H}} \beta \otimes_{\alpha} \sigma_{\nu^o})$$

is here replacing the rather akward writing :

$$(\sigma_{\nu^o} \alpha \otimes_{\hat{\beta}} 1_{\mathfrak{H}})(1_{\mathfrak{H}} \alpha \otimes_{\hat{\beta}} W) \sigma_{2\nu} (1_{\mathfrak{H}} \beta \otimes_{\alpha} \sigma_{\nu^o})$$

but denotes the same operator, and we suggest the reader to convince himself of this easy fact.

If we start from another normal semi-finite faithful weight ν' on N , we may define, using 2.3, another unitary $W^{\nu'} = U_{\alpha, \hat{\beta}}^{\nu^o, \nu'} W U_{\beta, \alpha}^{\nu', \nu}$ from $\mathfrak{H}_{\beta \otimes \alpha}$ onto $\mathfrak{H}_{\alpha \otimes \hat{\beta}}$. The formulae

which link these isomorphisms between relative product Hilbert spaces and the relative flips allow us to check that this operator $W^{\nu'}$ is also pseudo-multiplicative; which can be resumed in saying that a pseudo-multiplicative unitary does not depend on the choice of the weight on N .

If W is an $(\alpha, \hat{\beta}, \beta)$ -pseudo-multiplicative unitary, then the unitary $\sigma_{\nu} W^* \sigma_{\nu}$ from $\mathfrak{H}_{\hat{\beta} \otimes \alpha}$ to $\mathfrak{H}_{\alpha \otimes \beta}$ is an $(\alpha, \beta, \hat{\beta})$ -pseudo-multiplicative unitary, called the dual of W .

3.3. Algebras and Hopf-bimodules associated to a pseudo-multiplicative unitary. For ξ_2 in $D(\alpha \mathfrak{H}, \nu)$, η_2 in $D(\mathfrak{H}_{\hat{\beta}}, \nu^o)$, the operator $(\rho_{\eta_2}^{\alpha, \hat{\beta}})^* W \rho_{\xi_2}^{\beta, \alpha}$ will be written $(id * \omega_{\xi_2, \eta_2})(W)$; we have, therefore, for all ξ_1, η_1 in \mathfrak{H} :

$$((id * \omega_{\xi_2, \eta_2})(W) \xi_1 | \eta_1) = (W(\xi_1_{\beta \otimes \alpha} \xi_2) | \eta_1_{\alpha \otimes \hat{\beta}} \eta_2)$$

and, using the intertwining property of W with $\hat{\beta}$, we easily get that $(id * \omega_{\xi_2, \eta_2})(W)$ belongs to $\hat{\beta}(N)'$.

If x belongs to N , we have :

$$(id * \omega_{\xi_2, \eta_2})(W)\alpha(x) = (id * \omega_{\xi_2, \alpha(x^*)\eta_2})(W)$$

$$\beta(x)(id * \omega_{\xi_2, \eta_2})(W) = (id * \omega_{\hat{\beta}(x)\xi_2, \eta_2})(W)$$

We shall write $A_w(W)$ the weak closure of the linear span of these operators, which are right $\alpha(N)$ -modules and left $\beta(N)$ -modules. Applying ([E3] 3.6), we get that $A_w(W)$ and $A_w(W)^*$ are non-degenerate algebras (one should note that the notations of ([E3]) had been changed in order to fit with Lesieur's notations). We shall write $\mathcal{A}(W)$ the von Neumann algebra generated by $A_w(W)$. We then have $\mathcal{A}(W) \subset \hat{\beta}(N)'$.

For ξ_1 in $D(\mathfrak{H}_\beta, \nu^o)$, η_1 in $D({}_\alpha\mathfrak{H}, \nu)$, we shall write $(\omega_{\xi_1, \eta_1} * id)(W)$ for the operator $(\lambda_{\eta_1}^{\alpha, \hat{\beta}})^* W \lambda_{\xi_1}^{\beta, \alpha}$; we have, therefore, for all ξ_2, η_2 in \mathfrak{H} :

$$((\omega_{\xi_1, \eta_1} * id)(W)\xi_2|\eta_2) = (W(\xi_1 \underset{\nu}{\beta} \otimes_{\alpha} \xi_2)|\eta_1 \underset{\nu^o}{\alpha} \otimes_{\hat{\beta}} \eta_2)$$

and, using the intertwining property of W with β , we easily get that $(\omega_{\xi_1, \eta_1} * id)(W)$ belongs to $\beta(N)'$.

We shall write $\widehat{A_w(W)}$ the weak closure of the linear span of these operators. It is clear that this weakly closed subspace is a non degenerate algebra; following ([EV] 6.1 and 6.5), we shall write $\widehat{\mathcal{A}(W)}$ the von Neumann algebra generated by $\widehat{A_w(W)}$. We then have $\widehat{\mathcal{A}(W)} \subset \beta(N)'$.

In ([EV] 6.3 and 6.5), using the pentagonal equation, we got that $(N, \mathcal{A}(W), \alpha, \beta, \Gamma)$, and $(N, \widehat{\mathcal{A}(W)}, \alpha, \hat{\beta}, \hat{\Gamma})$ are Hopf-bimodules, where Γ and $\hat{\Gamma}$ are defined, for any x in $\mathcal{A}(W)$ and y in $\widehat{\mathcal{A}(W)}$, by :

$$\Gamma(x) = W^*(1 \underset{N^o}{\alpha} \otimes_{\hat{\beta}} x)W$$

$$\hat{\Gamma}(y) = \sigma_{\nu^o} W(y \underset{N}{\beta} \otimes_{\alpha} 1)W^* \sigma_{\nu}$$

(Here also, we have changed the notations of [EV], in order to fit with Lesieur's notations). In ([EV] 6.1(iv)), we had obtained that x in $\mathcal{L}(\mathfrak{H})$ belongs to $\mathcal{A}(W)'$ if and only if x belongs to $\alpha(N)' \cap \beta(N)'$ and verifies $(x \underset{N^o}{\alpha} \otimes_{\hat{\beta}} 1)W = W(x \underset{N}{\beta} \otimes_{\alpha} 1)$. We obtain the same

way that y in $\mathcal{L}(\mathfrak{H})$ belongs to $\widehat{\mathcal{A}(W)}'$ if and only if y belongs to $\alpha(N)' \cap \hat{\beta}(N)'$ and verifies $(1 \underset{N^o}{\alpha} \otimes_{\hat{\beta}} y)W = W(1 \underset{N}{\beta} \otimes_{\alpha} y)$.

Moreover, we get that $\alpha(N) \subset \mathcal{A} \cap \hat{\mathcal{A}}$, $\beta(N) \subset \mathcal{A}$, $\hat{\beta}(N) \subset \hat{\mathcal{A}}$, and, for all x in N :

$$\Gamma(\alpha(x)) = \alpha(x) \underset{N}{\beta} \otimes_{\alpha} 1$$

$$\Gamma(\beta(x)) = 1 \underset{N}{\beta} \otimes_{\alpha} \beta(x)$$

$$\hat{\Gamma}(\alpha(x)) = \alpha(x) \underset{N}{\hat{\beta}} \otimes_{\alpha} 1$$

$$\hat{\Gamma}(\hat{\beta}(x)) = 1 \underset{N}{\hat{\beta}} \otimes_{\alpha} \hat{\beta}(x)$$

3.4. Fundamental example. Let \mathcal{G} be a measured groupoid; let's use all notations introduced in 3.1. Let us note :

$$\mathcal{G}_{r,r}^2 = \{(x, y) \in \mathcal{G}^2, r(x) = r(y)\}$$

Then, it has been shown [Val2] that the formula $W_{\mathcal{G}}f(x, y) = f(x, x^{-1}y)$, where x, y are in \mathcal{G} , such that $r(y) = r(x)$, and f belongs to $L^2(\mathcal{G}^{(2)})$ (with respect to an appropriate measure, constructed from λ^u and ν), is a unitary from $L^2(\mathcal{G}^{(2)})$ to $L^2(\mathcal{G}_{r,r}^2)$ (with respect also to another appropriate measure, constructed from λ^u and ν).

Let us define $r_{\mathcal{G}}$ and $s_{\mathcal{G}}$ from $L^\infty(\mathcal{G}^{(0)}, \nu)$ to $L^\infty(\mathcal{G}, \mu)$ (and then considered as representations on $\mathcal{L}(L^2(\mathcal{G}, \mu))$, for any f in $L^\infty(\mathcal{G}^{(0)}, \nu)$, by $r_{\mathcal{G}}(f) = f \circ r$ and $s_{\mathcal{G}}(f) = f \circ s$).

We shall identify ([Y3], 3.2.2) the Hilbert space $L^2(\mathcal{G}^{(2)})$ with the relative Hilbert tensor product $L^2(\mathcal{G}, \mu) \underset{L^\infty(\mathcal{G}^{(0)}, \nu)}{\otimes_{r_{\mathcal{G}}}} L^2(\mathcal{G}, \mu)$, and the Hilbert space $L^2(\mathcal{G}_{r,r}^2)$ with the relative Hilbert

tensor product $L^2(\mathcal{G}, \mu) \underset{L^\infty(\mathcal{G}^{(0)}, \nu)}{\otimes_{r_{\mathcal{G}}}} L^2(\mathcal{G}, \mu)$. Moreover, the unitary $W_{\mathcal{G}}$ can be then inter-

preted [Val3] as a pseudo-multiplicative unitary over the basis $L^\infty(\mathcal{G}^{(0)}, \nu)$, with respect to the representation $r_{\mathcal{G}}$, and anti-representations $s_{\mathcal{G}}$ and $r_{\mathcal{G}}$ (as here the basis is abelian, the notions of representation and anti-representations are the same, and the commutation property is fulfilled). So, we get that $W_{\mathcal{G}}$ is a $(r_{\mathcal{G}}, s_{\mathcal{G}}, r_{\mathcal{G}})$ pseudo-multiplicative unitary.

Let us take the notations of 3.3; the von Neumann algebra $\mathcal{A}(W_{\mathcal{G}})$ is equal to the von Neumann algebra $L^\infty(\mathcal{G}, \nu)$ ([Val3], 3.2.6 and 3.2.7); using ([Val3] 3.1.1), we get that the Hopf-bimodule homomorphism Γ defined on $L^\infty(\mathcal{G}, \mu)$ by $W_{\mathcal{G}}$ is equal to the usual Hopf-bimodule homomorphism $\Gamma_{\mathcal{G}}$ studied in [Val2], and recalled in 3.1. Moreover, the von Neumann algebra $\widehat{\mathcal{A}(W_{\mathcal{G}})}$ is equal to the von Neumann algebra $\mathcal{L}(\mathcal{G})$ ([Val3], 3.2.6 and 3.2.7); using ([Val3] 3.1.1), we get that the Hopf-bimodule homomorphism $\widehat{\Gamma}$ defined on $\mathcal{L}(\mathcal{G})$ by $W_{\mathcal{G}}$ is the usual Hopf-bimodule homomorphism $\widehat{\Gamma}_{\mathcal{G}}$ studied in [Y3] and [Val2]. Let us suppose now that the groupoid \mathcal{G} is locally compact in the sense of [R1]; it has been proved in ([E3] 4.8) that $W_{\mathcal{G}}$ satisfies a strong condition of regularity, called "norm regularity".

3.5. Definitions ([L1], [L2]). Let $(N, M, \alpha, \beta, \Gamma)$ be a Hopf-bimodule, as defined in 3.1; a normal, semi-finite, faithful operator valued weight T from M to $\alpha(N)$ is said to be left-invariant if, for all $x \in \mathfrak{M}_T^+$, we have :

$$(id_{\beta * \alpha} T) \underset{N}{\Gamma}(x) = T(x) \underset{N}{\beta \otimes \alpha} 1$$

or, equivalently (2.5), if we write $\Phi = \nu \circ \alpha^{-1} \circ T$:

$$(id_{\beta * \alpha} \Phi) \underset{N}{\Gamma}(x) = T(x)$$

A normal, semi-finite, faithful operator-valued weight T' from M to $\beta(N)$ will be said to be right-invariant if it is left-invariant with respect to the symmetrized Hopf-bimodule, i.e., if, for all $x \in \mathfrak{M}_{T'}^+$, we have :

$$(T' \underset{N}{\beta * \alpha} id) \underset{N}{\Gamma}(x) = 1 \underset{N}{\beta \otimes \alpha} T'(x)$$

or, equivalently, if we write $\Psi = \nu \circ \beta^{-1} \circ T'$:

$$(\Psi \underset{N}{\beta * \alpha} id) \underset{N}{\Gamma}(x) = T'(x)$$

3.6. Theorem([L1], [L2]). Let $(N, M, \alpha, \beta, \Gamma)$ be a Hopf-bimodule, as defined in 3.1, and let T be a left-invariant normal, semi-finite, faithful operator valued weight from M to $\alpha(N)$; let us choose a normal, semi-finite, faithful weight ν on N , and let us write $\Phi = \nu \circ \alpha^{-1} \circ T$, which is a normal, semi-finite, faithful weight on M ; let us write H_Φ , J_Φ , Δ_Φ for the canonical objects of the Tomita-Takesaki theory associated to the weight Φ , and let us define, for x in N , $\hat{\beta}(x) = J_\Phi \alpha(x^*) J_\Phi$.

(i) There exists a unique isometry U from $H_\Phi \otimes_{\beta}^{\nu^o} H_\Phi$ to $H_\Phi \otimes_{\alpha}^{\nu} H_\Phi$, such that, for any (β, ν^o) -orthogonal basis $(\xi_i)_{i \in I}$ of $(H_\Phi)_\beta$, for any a in $\mathfrak{N}_T \cap \mathfrak{N}_\Phi$ and for any v in $D((H_\Phi)_\beta, \nu^o)$, we have

$$U(v \otimes_{\beta}^{\nu^o} \Lambda_\Phi(a)) = \sum_{i \in I} \xi_i \otimes_{\alpha}^{\nu} \Lambda_\Phi((\omega_{v, \xi_i} \otimes_{\beta}^{\nu^o} id)(\Gamma(a)))$$

(ii) Let us suppose there exists a right-invariant normal, semi-finite, faithful operator valued weight T' from M to $\beta(N)$; then this isometry is a unitary, and $W = U^*$ is an $(\alpha, \hat{\beta}, \beta)$ -pseudo-multiplicative unitary from $H_\Phi \otimes_{\alpha}^{\nu} H_\Phi$ to $H_\Phi \otimes_{\beta}^{\nu^o} H_\Phi$ which verifies, for any x, y_1, y_2 in $\mathfrak{N}_T \cap \mathfrak{N}_\Phi$:

$$(i * \omega_{J_\Phi \Lambda_\Phi(y_1^* y_2), \Lambda_\Phi(x)})(W) = (id \otimes_{\beta}^{\nu^o} \omega_{J_\Phi \Lambda_\Phi(y_2), J_\Phi \Lambda_\Phi(y_1)})(\Gamma(x^*))$$

Clearly, the pseudo-multiplicative unitary W does not depend upon the choice of the right-invariant operator-valued weight T' , and, for any y in M , we have :

$$\Gamma(y) = W^*(1 \otimes_{\beta}^{\nu^o} y)W$$

Proof. This is [L2] 3.51 and 3.52. □

3.7. Definitions. Let us take the notations of 3.6; let us write $\Psi = \nu \circ \beta^{-1} \circ T'$. We shall say that ν is relatively invariant with respect to T and T' if the two modular automorphism groups associated to the two weights Φ and Ψ commute; we then write down:

Definition

A measured quantum groupoid is an octuplet $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ such that :

- (i) $(N, M, \alpha, \beta, \Gamma)$ is a Hopf-bimodule, as defined in 3.1,
- (ii) T is a left-invariant normal, semi-finite, faithful operator valued weight T from M to $\alpha(N)$, as defined in 3.5,
- (iii) T' is a right-invariant normal, semi-finite, faithful operator-valued weight T' from M to $\beta(N)$, as defined in 3.5,
- (iv) ν is normal semi-finite faithful weight on N , which is relatively invariant with respect to T and T' .

Remark These axioms are not Lesieur's axioms, given in ([L2], 4.1). The equivalence of these axioms with Lesieur's axioms had been written down in [E6], and is recalled in the appendix (B.8).

3.8. Theorem ([L2], [E6]). Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7. Let us write $\Phi = \nu \circ \alpha^{-1} \circ T$, which is a normal, semi-finite faithful weight on M . Then

(i) there exists a $*$ -antiautomorphism R on M , such that $R^2 = id$, $R(\alpha(n)) = \beta(n)$ for all $n \in N$, and

$$\Gamma \circ R = \varsigma_{N^\circ} (R \underset{N}{\beta^* \alpha} R) \Gamma$$

R will be called the coinverse;

(ii) there exists a one-parameter group τ_t of automorphisms of M , such that $R \circ \tau_t = \tau_t \circ R$ for all $t \in \mathbb{R}$, and, for all $t \in \mathbb{R}$ and $n \in N$, $\tau_t(\alpha(n)) = \alpha(\sigma_t^\nu(n))$, $\tau_t(\beta(n)) = \beta(\sigma_t^\nu(n))$ and :

$$\Gamma \circ \tau_t = (\tau_t \underset{N}{\beta^* \alpha} \tau_t) \Gamma$$

$$\Gamma \circ \sigma_t^\Phi = (\tau_t \underset{N}{\beta^* \alpha} \sigma_t^\Phi) \Gamma$$

τ_t will be called the scaling group;

(iii) the weight ν is relatively invariant with respect to T and RTR ; moreover, R and τ_t are still the co-inverse and the scaling group of this new measured quantum groupoid, we shall denote

$$\underline{\mathfrak{G}} = (N, M, \alpha, \beta, \Gamma, T, RTR, \nu)$$

(iv) for any ξ, η in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^\circ)$, $(id * \omega_{\xi, \eta})(W)$ belongs to $D(\tau_{i/2})$, and, if we define $S = R\tau_{-i/2}$, we have :

$$S((id * \omega_{\xi, \eta})(W)) = (id * \omega_{\eta, \xi})(W)^*$$

More generally, for any x in $D(S) = D(\tau_{-i/2})$, we get that $S(x)^*$ belongs to $D(S)$ and $S(S(x)^*)^* = x$; S will be called the antipod of $\underline{\mathfrak{G}}$ (or $\underline{\mathfrak{G}}$), and, therefore, the co-inverse and the scaling group, given by polar decomposition of the antipod, rely only upon the pseudo-multiplicative W .

(v) there exists a one-parameter group γ_t of automorphisms of N such that, for all $t \in \mathbb{R}$ and $n \in N$, we have :

$$\sigma_t^T(\beta(n)) = \beta(\gamma_t(n))$$

Moreover, we get that $\nu \circ \gamma_t = \nu$.

(vi) there exists a positive non-singular operator λ affiliated to $Z(M)$, and a positive non singular operator δ affiliated to M , such that :

$$(D\Phi \circ R : D\Phi)_t = \lambda^{it^2/2} \delta^{it}$$

and, therefore, we have :

$$(D\Phi \circ \sigma_s^{\Phi \circ R} : D\Phi)_t = \lambda^{ist}$$

The operator λ will be called the scaling operator, and there exists a positive non-singular operator q affiliated to N such that $\lambda = \alpha(q) = \beta(q)$. We have $R(\lambda) = \lambda$.

The operator δ will be called the modulus; we have $R(\delta) = \delta^{-1}$, and $\tau_t(\delta) = \delta$, for all $t \in \mathbb{R}$, and we can define a one-parameter group of unitaries $\delta^{it} \underset{N}{\beta \otimes \alpha} \delta^{it}$ which acts naturally on elementary tensor products, which verifies, for all $t \in \mathbb{R}$:

$$\Gamma(\delta^{it}) = \delta^{it} \underset{N}{\beta \otimes \alpha} \delta^{it}$$

(vii) we have $(D\Phi \circ \tau_t : D\Phi)_s = \lambda^{-ist}$, which leads to define a one-parameter group of unitaries by, for any $x \in \mathfrak{N}_\Phi$:

$$P^{it} \Lambda_\Phi(x) = \lambda^{t/2} \Lambda_\Phi(\tau_t(x))$$

Moreover, for any y in M , we get :

$$\tau_t(y) = P^{it} y P^{-it}$$

and it is possible to define one parameter groups of unitaries $P_N^{it} \beta \otimes_\alpha P^{it}$ and $P_{N^o}^{it} \alpha \otimes_{\hat{\beta}} P^{it}$ such that :

$$W(P_N^{it} \beta \otimes_\alpha P^{it}) = (P_{N^o}^{it} \alpha \otimes_{\hat{\beta}} P^{it})W$$

Moreover, for all $v \in D(P^{-1/2})$, $w \in D(P^{1/2})$, p, q in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^o)$, we have:

$$(W^*(v_{\nu^o} \alpha \otimes_{\hat{\beta}} q)|w_{\nu} \beta \otimes_\alpha p) = (W(P^{-1/2} v_{\nu} \beta \otimes_\alpha J_\Phi p)|P^{1/2} w_{\nu^o} \alpha \otimes_{\hat{\beta}} J_\Phi q)$$

We shall say that the pseudo-multiplicative unitary W is "manageable", with "managing operator" P , which implies it is weakly regular in the sense of [E3], 4.1, which implies (with the notations of 3.3) that $A_w(W) = \mathcal{A}(W) = M$ and $\widehat{A_w(W)} = \widehat{\mathcal{A}(W)} = \widehat{M}$. As $\tau_t \circ \sigma_t^\Phi = \sigma_t^\Phi \circ \tau_t$, we get that $J_\Phi P J_\Phi = P$.

Proof. Result (i) is A.6, result (ii) is A.5, result (iii) is B.7(iv) and A.8; result (iv) is A.9, result (v) is B.2 and B.7(iii); result (vi) is given by 2.6 and (iii), B.7(iv), (v), and, using B.8, [L2], 5.6 and 5.20; result (vii), using B.8, is [L2] 7.3. \square

3.9. Notations. We shall use the notations of 2.6 about Vaes' Radon-Nykodym theorem for the weights Φ and $\Phi \circ R$. Let λ be the scaling operator of $\Phi \circ R$ with respect to Φ , and δ be the modulus of $\Phi \circ R$ with respect to Φ , defined in 3.8(vi). Then, the self-adjoint elements e_n of M defined in 2.6 verify $R(e_n) = e_n$.

If $x \in M$ is such that $x\delta^{1/2}$ is bounded and its closure $\overline{x\delta^{1/2}}$ belongs to \mathfrak{N}_Φ , then x belongs to $\mathfrak{N}_{\Phi \circ R}$ (2.6), and, then $R(x^*)$ belongs to \mathfrak{N}_Φ . In particular, if $x \in \mathfrak{N}_\Phi$, then xe_n belongs to $\mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$ (and $R(x^*)e_n$ belongs also to $\mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$), and $\lim_n \Lambda_\Phi(xe_n) = \Lambda_\Phi(x)$.

3.10. Theorem ([L2]). Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7. Let \mathcal{T} be the subset of $\mathfrak{N}_T \cap \mathfrak{N}_{RTR} \cap \mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$ made of elements x which are analytic with respect to both Φ and $\Phi \circ R$, and such that, for all z, z' in \mathbb{C} , $\sigma_z^\Phi \circ \sigma_{z'}^{\Phi \circ R}(x)$ belongs to $\mathfrak{N}_T \cap \mathfrak{N}_{RTR} \cap \mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$. Let E be the linear subspace generated by the elements of the form $e_n x$, where e_n had been defined in 3.9, and $x \in \mathcal{T}$, and let us denote \mathcal{E} the linear subspace $J_\Phi \Lambda_\Phi(E)$. Let us denote $M_*^{\alpha, \beta}$ the subspace of M_* spanned by the positive elements ω in M_*^+ such that there exists $k \in \mathbb{R}^+$ such that $\omega \circ \alpha \leq k\nu$ and $\omega \circ \beta \leq k\nu$. Then :

(i) \mathcal{T} and E are dense in M ; $\Lambda_\Phi(\mathcal{T})$ and \mathcal{E} are dense in H_Φ . As

$$\mathcal{E} \subset D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$$

we get that $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$ is dense in H_Φ ; therefore $M_*^{\alpha, \beta}$ is dense in M_* . Moreover, if η, η' are in \mathcal{E} , then $\langle \eta, \eta' \rangle_{\alpha, \nu}$ and $\langle \eta, \eta' \rangle_{\beta, \nu^o}$ are analytic with respect to both σ_t^ν and γ_t .

(ii) If ω_1, ω_2 are in $M_*^{\alpha, \beta}$, let us define $\omega_1 \omega_2 = (\omega_1 \beta *_{\alpha} \omega_2) \Gamma$. If $\omega \in M_*^{\alpha, \beta}$ and $x \in M$,

let us define $\omega^*(x) = \overline{\omega(R(x^*))}$; with such product and involution, $M_*^{\alpha, \beta}$ is an involutive algebra, and we can define a faithful representation of this involutive algebra by :

$$\hat{\pi}(\omega) = (\omega * id)(W)$$

Let \widehat{M} be the weak closure of the algebra generated (which is a von Neumann algebra, by 3.8 (vii)) and is included in $\beta(N)'$. More generally, for any v in $D({}_\alpha H_\Phi, \nu)$, w in

$D((H_\Phi)_\beta, \nu^o)$, let us write $\hat{\pi}(\omega_{v,w}) = (\omega_{v,w} * id)(W)$; then $\hat{\pi}(\omega_{v,w})$ belongs to \widehat{M} . If, moreover, v belongs also to $D((H_\Phi)_\beta, \nu^o)$, we have, for all $x \in \mathfrak{N}_\Phi$:

$$\Lambda_\Phi(\omega_{w,v} \underset{N}{\beta} *_{\alpha} id) \Gamma(x)) = \hat{\pi}(\omega_{v,w})^* \Lambda_\Phi(x)$$

(iii) If y is in \widehat{M} , let us define :

$$\widehat{\Gamma}(y) = \sigma_{\nu^o} W(y \underset{N}{\beta} \otimes_{\alpha} 1) W^* \sigma_{\nu}$$

with these notations, $(N, \widehat{M}, \alpha, \hat{\beta}, \widehat{\Gamma})$ is a Hopf-bimodule.

(iv) there exists a unique $*$ -anti-homomorphism \widehat{R} of \widehat{M} , such that $\widehat{R}\hat{\pi}(\omega) = \hat{\pi}(\omega \circ R)$, for all $\omega \in M_*^{\alpha, \beta}$. Moreover, we have $\widehat{R}(y) = J_\Phi y^* J_\Phi$ for all $y \in \widehat{M}$, $\widehat{R}(\alpha(n)) = \hat{\beta}(n)$, for all $n \in N$, and :

$$\widehat{\Gamma} \circ \widehat{R} = \varsigma_{N^o}(\widehat{R} \underset{N}{\beta} *_{\alpha} \widehat{R}) \widehat{\Gamma}$$

and, therefore, \widehat{R} is a co-inverse of the Hopf-bimodule $(N, \widehat{M}, \alpha, \hat{\beta}, \widehat{\Gamma})$.

(v) let $I_\Phi \subset M_*$ be defined as :

$$I_\Phi = \{\omega \in M_*; \exists k' \in \mathbb{R}^+, |\omega(x^*)| \leq k' \|\Lambda_\Phi(x)\|, \forall x \in \mathfrak{N}_\Phi\}$$

and, for any $\omega \in I_\Phi$, let $a_\Phi(\omega)$ be the unique vector in H_Φ such that, for all $x \in \mathfrak{N}_\Phi$:

$$(a_\Phi(\omega) | \Lambda_\Phi(x)) = \omega(x^*)$$

Then $I_\Phi \cap M_*^{\alpha, \beta}$ is a left ideal of $M_*^{\alpha, \beta}$, and there exists a unique normal semifinite faithful weight $\widehat{\Phi}$ on \widehat{M} , such that $\hat{\pi}(I_\Phi \cap M_*^{\alpha, \beta})$ is a core for $\Lambda_{\widehat{\Phi}}$. Moreover, we may identify $H_{\widehat{\Phi}}$ with H_Φ in such a way that we have, for any $\omega \in M_*^{\alpha, \beta} \cap I_\Phi$:

$$\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega)) = a_\Phi(\omega)$$

We have also, with the notations of 3.9, for all $x \in \mathfrak{N}_\Phi$ and $n \in N$:

$$J_{\widehat{\Phi}} \Lambda_\Phi(x e_n) = \Lambda_\Phi(R(x^*) \overline{e_n \delta^{1/2}})$$

and, for any $y \in M$, we get $R(y) = J_{\widehat{\Phi}} y^* J_{\widehat{\Phi}}$.

Moreover, there exists a unique left-invariant normal faithful semi-finite operator-valued weight \widehat{T} from \widehat{M} to $\alpha(N)$ such that

$$\widehat{\Phi} = \nu \circ \alpha^{-1} \circ \widehat{T}$$

(vi) the octuplet $(N, \widehat{M}, \alpha, \hat{\beta}, \widehat{\Gamma}, \widehat{T}, \widehat{R} \circ \widehat{T} \circ \widehat{R}, \nu)$ is a measured quantum groupoid, denoted by $\widehat{\mathfrak{G}}$ and called the dual of \mathfrak{G} (or of $\underline{\mathfrak{G}}$).

(vii) The pseudo-multiplicative unitary \widehat{W} constructed from $\widehat{\mathfrak{G}}$ is :

$$\widehat{W} = \sigma_{\nu} W^* \sigma_{\nu}$$

The scaling group $\hat{\tau}_t$ of $\widehat{\mathfrak{G}}$ is given, for all $y \in \widehat{M}$, by :

$$\hat{\tau}_t(y) = P^{it} y P^{-it}$$

The scaling operator $\hat{\lambda}$ of $\widehat{\mathfrak{G}}$ satisfies $\hat{\lambda} = \lambda^{-1}$. The managing operator \hat{P} of \widehat{W} is equal to P .

The modular operator $\Delta_{\widehat{\Phi}}$ is equal to the closure of $P J_\Phi \delta^{-1} J_\Phi$, and, therefore, for all $x \in M$, we have :

$$\tau_t(x) = \Delta_{\widehat{\Phi}}^{it} x \Delta_{\widehat{\Phi}}^{-it}$$

The automorphism group $\hat{\gamma}_t$ is equal to γ_{-t} .

Proof. Result (i) is [L2] 6.7, result (ii) is [L2] 8.4, 8.5, 3.18 and 8.1; result (iii) is [L2] 8.2, (iv) is [L2] 8.6, (v) is [L2] 8.7, 8.14, 8.18 and 8.28, and results (vi) and (vii) are [L2] 8.24. \square

3.11. Theorem ([L2]). *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7, $\widehat{\mathfrak{G}}$ its dual in the sense of 3.10; then :*

- (i) *the bidual $\widehat{\widehat{\mathfrak{G}}}$ is equal to \mathfrak{G} ;*
- (ii) *the modulus $\hat{\delta}$ is equal to the closure of $P^{-1}J_{\Phi}\delta J_{\Phi}\delta^{-1}\Delta_{\Phi}^{-1}$; therefore, we have $\overline{\delta\Delta_{\Phi}} = \overline{\hat{\delta}\Delta_{\Phi}}^{-1}$.*
- (iii) *it is possible to define one-parameter groups of unitaries $\Delta_{\Phi}^{it}{}_{\beta\otimes_{\nu}\alpha}\Delta_{\Phi}^{it}$ and $\Delta_{\Phi}^{it}{}_{\alpha\otimes_{\nu^o}\beta}\Delta_{\Phi}^{it}$ and we have :*

$$W(\Delta_{\Phi}^{it}{}_{\beta\otimes_{\nu}\alpha}\Delta_{\Phi}^{it}) = (\Delta_{\Phi}^{it}{}_{\alpha\otimes_{\nu^o}\beta}\Delta_{\Phi}^{it})W$$

$$W(J_{\Phi}{}_{\alpha\otimes_{\nu^o}\beta}J_{\Phi}) = (J_{\Phi}{}_{\beta\otimes_{\nu}\alpha}J_{\Phi})W^*$$

(iv) *for all s, t in \mathbb{R} , we have :*

$$\Delta_{\Phi}^{it}\Delta_{\Phi}^{is} = \lambda^{ist}\Delta_{\Phi}^{is}\Delta_{\Phi}^{it}$$

and we have $J_{\widehat{\Phi}}J_{\Phi} = \lambda^{i/4}J_{\Phi}J_{\widehat{\Phi}}$.

(v) *we have the following "Heisenberg-type" relations :*

$$M \cap \widehat{M} = \alpha(N)$$

$$M \cap \widehat{M}' = \beta(N)$$

$$M' \cap \widehat{M} = \hat{\beta}(N)$$

$$M' \cap \widehat{M}' = \hat{\alpha}(N)$$

where we put, for all $n \in N$, $\hat{\alpha}(n) = J_{\Phi}\beta(n)^*J_{\Phi} = J_{\widehat{\Phi}}\hat{\beta}(n^*)J_{\widehat{\Phi}}$.

Proof. Result (i) is [L2] 8.25, (ii) is [L2] 8.24 (iii); results (iii) and (iv) are [L2] 8.28, and (v) is [L2] 8.30. \square

3.12. Theorem([L2]). *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7, $\widehat{\mathfrak{G}}$ its dual in the sense of 3.10. Then :*

- (i) *The octuplet $(N^o, M, \beta, \alpha, \varsigma_N \circ \Gamma, RTR, T, \nu^o)$ is a measured quantum groupoid we shall write \mathfrak{G}^o and call the opposite measured quantum groupoid. We have $(\mathfrak{G}^o)^o = \mathfrak{G}$.*
- (ii) *For any x in M , let us write $j(x) = J_{\Phi}x^*J_{\Phi}$ the canonical $*$ -anti-homomorphism from M onto M' and let us define then $\Gamma^c = (j{}_{\beta*_{\alpha}}j)_{\Gamma} \circ j$, $R^c = j \circ R \circ j$, $T^c = j \circ T \circ j$.*

The octuplet $(N^o, M', \hat{\beta}, \hat{\alpha}, \Gamma^c, T^c, R^cT^cR^c, \nu^o)$ is a measured quantum groupoid we shall write \mathfrak{G}^c and call the commutant of \mathfrak{G} . We have $(\mathfrak{G}^c)^c = \mathfrak{G}$.

(iii) *we have $\mathfrak{G}^{oc} = \mathfrak{G}^{co}$ and this last measured quantum groupoid is isomorphic to \mathfrak{G} , via the isomorphism implemented by $J_{\Phi}J_{\widehat{\Phi}}$.*

(iv) *We have $\widehat{\mathfrak{G}^o} = (\widehat{\mathfrak{G}})^c$ and $\widehat{\mathfrak{G}^c} = (\widehat{\mathfrak{G}})^o$*

(v) *The pseudo-multiplicative unitaries W^o and W^c of \mathfrak{G}^o and \mathfrak{G}^c are given by :*

$$W^o = (J_{\widehat{\Phi}}{}_{\alpha\otimes_{\beta}J_{\widehat{\Phi}}})_{N^o}W(J_{\widehat{\Phi}}{}_{\alpha\otimes_{\beta}J_{\widehat{\Phi}}})_{N^o}$$

$$W^c = (J_{\Phi}{}_{\alpha\otimes_{\beta}J_{\Phi}})_{N^o}W(J_{\Phi}{}_{\alpha\otimes_{\beta}J_{\Phi}})_{N^o}$$

where W^o is a unitary from $H_{\Phi} \underset{\nu^o}{\alpha} \otimes_{\beta} H_{\Phi}$ onto $H_{\Phi} \underset{\nu}{\beta} \otimes_{\hat{\alpha}} H_{\Phi}$ and W^c is a unitary from $H_{\Phi} \underset{\nu^o}{\hat{\alpha}} \otimes_{\hat{\beta}} H_{\Phi}$ onto $H_{\Phi} \underset{\nu}{\beta} \otimes_{\alpha} H_{\Phi}$.

Moreover, for any $x \in M$, $y \in \widehat{M}$, we have :

$$\Gamma(x) = (\sigma_{\nu} W^o \sigma_{\nu})^* (x \underset{N^o}{\hat{\alpha}} \otimes_{\beta} 1) (\sigma_{\nu} W^o \sigma_{\nu})$$

$$\widehat{\Gamma}(y) = W^c (y \underset{N^o}{\hat{\alpha}} \otimes_{\hat{\beta}} 1) (W^c)^*$$

(vi) The pseudo-multiplicative unitary W^{oc} of \mathfrak{G}^{oc} is given by :

$$W^{oc} = (J_{\Phi} J_{\widehat{\Phi}} \underset{N^o}{\alpha} \otimes_{\hat{\beta}} J_{\Phi} J_{\widehat{\Phi}}) W (J_{\widehat{\Phi}} J_{\Phi} \underset{N}{\beta} \otimes_{\hat{\alpha}} J_{\widehat{\Phi}} J_{\Phi})$$

and is a unitary from $H_{\Phi} \underset{\nu}{\beta} \otimes_{\hat{\alpha}} H_{\Phi}$ onto $H_{\Phi} \underset{\nu^o}{\hat{\alpha}} \otimes_{\beta} H_{\Phi}$.

Let us denote $\mathcal{I}_{\mathfrak{G}}$ the isomorphism between M and M' given, for $x \in M$, by :

$$\mathcal{I}_{\mathfrak{G}}(x) = J_{\Phi} J_{\widehat{\Phi}} x J_{\widehat{\Phi}} J_{\Phi}$$

Then $\mathcal{I}_{\mathfrak{G}}$ is an isomorphism of measured quantum groupoids between \mathfrak{G} and \mathfrak{G}^{oc} .

Moreover, for any $y \in \widehat{M'}$, we have :

$$\widehat{\Gamma}^c(y) = W^{oc} (y \underset{N}{\beta} \otimes_{\hat{\alpha}} 1) (W^{oc})^*$$

Proof. Result (i) is [L2] 17.8, (ii) is [L2] 17.10, (iii) and (iv) are [L2] 17.14; result (v) is [L2] 17.11, and (vi) is a straightforward corollary of the preceeding statements. \square

3.13. Example. Let \mathcal{G} be a measured groupoid; let's use all notations introduced in 3.1 and 3.4. If $f \in L^{\infty}(\mathcal{G}, \mu)^+$, let's consider the function on $\mathcal{G}^{(0)}$, $u \mapsto \int_{\mathcal{G}} f d\lambda^u$, which belongs to $L^{\infty}(\mathcal{G}^{(0)}, \nu)$; the image of this function by the homomorphism $r_{\mathcal{G}}$ is the function on \mathcal{G} , $\gamma \mapsto \int_{\mathcal{G}} f d\lambda^{r(\gamma)}$; the application which sends f on this function can be considered as an operator-valued weight from $L^{\infty}(\mathcal{G}, \mu)$ to $r_{\mathcal{G}}(L^{\infty}(\mathcal{G}^{(0)}, \nu))$ which is normal, semi-finite, faithful; by definition of the Haar system $(\lambda^u)_{u \in \mathcal{G}^{(0)}}$, it is left-invariant, in the sense of 3.5; we shall denote by $T_{\mathcal{G}}$ this operator-valued weight from $L^{\infty}(\mathcal{G}, \mu)$ to $r_{\mathcal{G}}(L^{\infty}(\mathcal{G}^{(0)}, \nu))$. If we write λ_u for the image of λ^u under the inverse $x \mapsto x^{-1}$ of the groupoid \mathcal{G} , starting from the application which sends f to the function on $\mathcal{G}^{(0)}$ defined by $u \mapsto \int_{\mathcal{G}} f d\lambda_u$, we define a normal semifinite faithful operator-valued weight from $L^{\infty}(\mathcal{G}, \mu)$ to $s_{\mathcal{G}}(L^{\infty}(\mathcal{G}^{(0)}, \nu))$, which is right-invariant in the sense of 3.5, we shall denote by $T_{\mathcal{G}}^{(-1)}$.

We then get that :

$$(L^{\infty}(\mathcal{G}^{(0)}, \nu), L^{\infty}(\mathcal{G}, \mu), r_{\mathcal{G}}, s_{\mathcal{G}}, \Gamma_{\mathcal{G}}, T_{\mathcal{G}}, T_{\mathcal{G}}^{(-1)}, \nu)$$

is a measured quantum groupoid, we shall denote again \mathcal{G} .

It can be proved ([E5]) that any measured quantum groupoid, whose underlying von Neumann algebra is abelian, is of that type.

The constructions obtained by applying 3.8 to this measured quantum groupoid are the following : the pseudo-multiplicative unitary is the operator $W_{\mathcal{G}}$ defined in 3.4; the co-inverse is given by dualizing the inverse $x \mapsto x^{-1}$ of \mathcal{G} to $L^{\infty}(\mathcal{G}, \mu)$; the scaling group is trivial, and the co-inverse is equal to the antipod; the automorphism group γ_t is trivial, the scaling operator λ is 1, and the modulus δ is the usual Radon-Nikodym derivative between μ and its image under the inverse $x \mapsto x^{-1}$ of \mathcal{G} .

The underlying Hopf bimodule of the dual $\widehat{\mathcal{G}}$ constructed in 3.10 is the symmetric Hopf-bimodule $(L^{\infty}(\mathcal{G}^{(0)}, \nu), \mathcal{L}(\mathcal{G}), r_{\mathcal{G}}, r_{\mathcal{G}}, \widehat{\Gamma}_{\mathcal{G}})$ constructed in [Y3] (3.1).

The constructions obtained by applying 3.12 are the following : $(\mathcal{G})^o$ is given by the opposite groupoid (in which we return the product, and exchange source and range applications). As $L^\infty(\mathcal{G}, \mu)$ is abelian, we have clearly $(\mathcal{G})^c = \mathcal{G}$.

3.14. Example. Let $\mathbf{G} = (M, \Gamma, \Phi, \Psi)$ be a locally compact quantum group, in the sense of Kustermans and Vaes ([KV1], or, preferably, the von Neumann version given in [KV2]). Here M is a von Neumann algebra, Γ a coassociative coproduct $M \mapsto M \otimes M$ (where \otimes is the von Neumann tensor product), and Φ (resp. Ψ) is a left-invariant (resp. a right-invariant) normal semi-finite faithful weight on M^+ . Then, thanks to ([KV1], prop. 6.1), we see that :

$$(\mathbb{C}, M, id, id, \Gamma, \Phi, \Psi, \omega)$$

(where id is the canonical homomorphism from \mathbb{C} into M , and ω is the canonical state on \mathbb{C}), is a measured quantum groupoid, we shall still denote by \mathbf{G} .

Conversely, it is clear that any measured quantum groupoid whose basis N is equal to \mathbb{C} is of that type.

3.15. Theorem([EV], [E4] 8.2 and 8.3). *Let $M_0 \subset M_1$ be a depth 2 inclusion of σ -finite von Neumann algebras, equipped with a regular (in the sense of 2.2) normal semi-finite faithful operator-valued weight T_1 . Then :*

(i) *there exists an application $\tilde{\Gamma}$ from $M'_0 \cap M_2$ to*

$$(M'_0 \cap M_2)_{j_1} \underset{M'_0 \cap M_1}{*} id(M'_0 \cap M_2)$$

such that $(M'_0 \cap M_1, M'_0 \cap M_2, id, j_1, \tilde{\Gamma})$ is a Hopf-bimodule, (where id means here the injection of $M'_0 \cap M_1$ into $M'_0 \cap M_2$, and j_1 means here the restriction of j_1 to $M'_0 \cap M_1$, considered then as an anti-representation of $M'_0 \cap M_1$ into $M'_0 \cap M_2$). Moreover, the anti-automorphism j_1 of $M'_0 \cap M_2$ is a co-inverse for this Hopf-bimodule structure.

(ii) *Let us write \tilde{T}_2 for the restriction of T_2 to $M'_0 \cap M_2$ (which is semi-finite, by definition of the regularity of T_1 (2.2)); we then get that \tilde{T}_2 is a left-invariant operator-valued weight from $M'_0 \cap M_2$ to $M'_0 \cap M_1$, and, therefore, that $j_1 \circ \tilde{T}_2 \circ j_1$ is a right-invariant operator-valued weight from $M'_0 \cap M_2$ to $M'_1 \cap M_2$.*

(iii) *Let us suppose moreover that there exists on $M'_0 \cap M_1$ a normal faithful semi-finite weight χ invariant under the modular automorphism group of T_1 and let χ_2 be the weight $\chi \circ \tilde{T}_2$; the modular automorphism groups $\sigma_t^{\chi_2}$ and $\sigma_s^{\chi_2 \circ j}$ commute.*

Therefore, $(M'_0 \cap M_1, M'_0 \cap M_2, id, j_1, \tilde{\Gamma}, \tilde{T}_2, j_1 \circ \tilde{T}_2 \circ j_1, \chi)$ is a measured quantum groupoid \mathfrak{G}_1 in the sense of 3.7.

Moreover, the inclusion $M_1 \subset M_2$ satisfies the same hypothesis, and leads to another measured quantum groupoid \mathfrak{G}_2 , which is isomorphic to $\widehat{\mathfrak{G}_1}_o$.

Let us describe how the coproduct $\tilde{\Gamma}$ is constructed : let ψ_0 be a normal semi-finite faithful weight on M_0 , $\psi_1 = \psi_0 \circ T_1$ the normal semi-finite faithful weight constructed then on M_1 , $\psi_2 = \psi_1 \circ T_2$ the normal semi-finite faithful weight constructed then on M_2 ; let χ be a normal semi-finite faithful weight on $M'_0 \cap M_1$, $\chi_2 = \chi \circ \tilde{T}_2$ the normal semi-finite faithful weight constructed then on $M'_0 \cap M_2$.

Then, the application defined, for $x \in \mathfrak{N}_{\tilde{T}_2} \cap \mathfrak{N}_{\chi_2}$ and $y \in \mathfrak{N}_{\psi_1}$ by :

$$U(\Lambda_{\chi_2}(x) \otimes_{\chi} \Lambda_{\psi_1}(y)) = \Lambda_{\psi_2}(xy)$$

is a unitary from $H_{\chi_2} \otimes_{\chi} H_{\psi_1}$ onto H_{ψ_2} ([EV], 4.2 and 4.6).

Moreover, using the fact that the inclusion $M_0 \subset M_2 \subset M_4$ is standard ([EN], 4.5), we get ([EV], 4.6) :

$$U^*[M'_0 \cap M_4 \cap (M'_0 \cap M_2)']U = \pi_{\chi_2}(M'_0 \cap M_2)' *_{M'_0 \cap M_1} M'_0 \cap M_2$$

and, on the other hand, composing the mirrorings (with the notations of (2.2)), we get that :

$$j_2 \circ j_1(M'_0 \cap M_2) = M'_2 \cap M_4 \subset M'_0 \cap M_4 \cap (M'_0 \cap M_2)'$$

and, therefore, $x \mapsto U^* j_2 \circ j_1(x) U$ is an injective $*$ -homomorphism from $M'_0 \cap M_2$ into $\pi_{\chi_2}(M'_0 \cap M_2)' *_{M'_0 \cap M_1} M'_0 \cap M_2$.

On the other hand, using U , it is possible to define a pseudo-multiplicative unitary \mathcal{W} ([EV], 5.3), which leads to a coproduct Γ on $\pi_{\chi_2}(M'_0 \cap M_2)'$ by the formula ($y \in \pi_{\chi_2}(M'_0 \cap M_2)'$) :

$$\Gamma(y) = \mathcal{W}(y \otimes_{\chi} 1) \mathcal{W}^*$$

and we have ([E4], 5.1), for all $x \in M'_0 \cap M_2$:

$$\mathcal{W}^*(\pi_{\chi_2}(x) \otimes_{\chi} 1) \mathcal{W} = \varsigma_{M'_0 \cap M_1}(id *_{M'_0 \cap M_1} \pi_{\chi_2})(U^* j_2 \circ j_1(x) U)$$

which shows that, if we write $W = (J_{\chi_2} \hat{J} \otimes_{\chi} 1) \sigma_{\chi^{\circ}} \mathcal{W} \sigma_{\chi^{\circ}} (\hat{J} J_{\chi_2} \otimes_{\chi} 1)$ (where \hat{J} is the standard implementation of $j_1|_{M'_0 \cap M_2}$ on H_{χ_2}), or, equivalently $\mathcal{W} = \widehat{W}^c$, we obtain a coproduct $\tilde{\Gamma}$ on $M'_0 \cap M_2$ which satisfies

$$\tilde{\Gamma}(x) = W^*(1 \otimes_{\chi^{\circ}} x) W = (J_{\chi_2} \hat{J} \otimes_{\chi} 1) U^* j_2 \circ j_1(x) U (\hat{J} J_{\chi_2} \otimes_{\chi} 1)$$

(in which, for simplification, we have identified $M'_0 \cap M_2$ with $\pi_{\chi_2}(M'_0 \cap M_2)$.)

4. LEFT INVARIANCE REVISITED

In the definition of a left-invariant operator-valued weight, recalled in 3.5, we have, for any $x \in \mathfrak{M}_T^+$, $(id_{\beta *_{\alpha} T}) \Gamma(x) = T(x)_{\beta \otimes_{\alpha} 1}$ (or, equivalently, $T(x) = (id_{\beta *_{\alpha} \Phi}) \Gamma(x)$, where $\Phi = \nu \circ \alpha^{-1} \circ T$). In this chapter, we extend this formula to any positive x in M , and even, any positive x in a fiber product $\mathcal{L}(\mathfrak{H})_{\beta *_{\alpha} M}$, where b is a faithful antirepresentation of N on \mathfrak{H} (4.12(i) and (ii)). On our way, we prove a Plancherel-like theorem (4.8), which extends the construction of the dual left-invariant operator-valued weight recalled in 3.10(iv). The result 4.12(ii) will be used in 6.12, and I am indebted to S. Vaes who pointed in [V2] how this kind of result was necessary in order to get correct proofs.

4.1. Definitions and notations. Let N be a von Neumann algebra, ψ a normal semi-finite faithful weight on N ; following ([ES2], 2.1.6), we define, for all $\omega \in M_*$:

$$\|\omega\|_{\psi} = \sup\{|\omega(x^*)|, x \in \mathfrak{N}_{\psi}, \psi(x^*x) \leq 1\}$$

and $I_{\psi} = \{\omega \in M_*, \|\omega\|_{\psi} < \infty\}$. Then $\omega \in M_*$ belongs to I_{ψ} if and only if there exists a vector $a_{\psi}(\omega)$ in H_{ψ} such that :

$$\omega(x^*) = (a_{\psi}(\omega) | \Lambda_{\psi}(x))$$

and we have then $\|a_{\psi}(\omega)\| = \|\omega\|_{\psi}$. Moreover, the application a here defined is clearly linear from I_{ψ} into H_{ψ} , and, ψ being semi-finite, injective.

Let us identify N and $\pi_{\psi}(N)$; then, any element $\omega \in M_*$ is of the form $\omega_{\xi, \eta}$, where ξ, η belong to H_{ψ} . For any η in H_{ψ} , let us define the following operator $\pi'(\eta)$, with $\mathcal{D}(\pi'(\eta)) = \Lambda_{\psi}(\mathfrak{N}_{\psi})$, and, for any $x \in \mathfrak{N}_{\psi}$, by $\pi'(\eta) \Lambda_{\psi}(x) = x \eta$. We then easily get that $\omega_{\xi, \eta}$ belongs to I_{ψ} if and only if ξ belongs to $\mathcal{D}(\pi'(\eta)^*)$, and we have then :

$$a_{\psi}(\omega_{\xi, \eta}) = \pi'(\eta)^* \xi$$

Let us write $K_{\eta} = \overline{\mathcal{D}(\pi'(\eta)^*)}$; then $(\pi'(\eta)^*)|_{K_{\eta}}$ is a densely defined operator from K_{η} into H_{ψ} , and we may consider $((\pi'(\eta)^*)|_{K_{\eta}})^*(\pi'(\eta)^*)|_{K_{\eta}}$ which is a positive self-adjoint operator

on K_η ; we shall write, for simplicity, \mathcal{P}_η for $((\pi'(\eta)^*)|_{K_\eta})^*(\pi'(\eta)^*)|_{K_\eta} + \infty(1 - p_{K_\eta})$, which belongs to the positive extension of N' . We have then, for any ξ, η in H_ψ :

$$\|\omega_{\xi,\eta}\|_\psi^2 = \langle \mathcal{P}_\eta, \omega_\xi \rangle$$

4.2. Lemma. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid; let u, v in H_Φ such that $\omega_{u,v}$ belongs to I_Φ , and let e_n the self-adjoint elements introduced in 3.9. Then, we have, for all $x \in \mathfrak{N}_\Phi$:*

$$(J_{\widehat{\Phi}} a_\Phi(\omega_{u,v}) | \Lambda_\Phi(x e_n)) = (R(x^*) \overline{e_n \delta^{1/2} v} | u)$$

where $\overline{e_n \delta^{1/2}}$ is the closure of the bounded operator $e_n \delta^{1/2}$.

If v belongs to $D(\delta^{1/2})$ and $\omega_{u,v}$ belongs to I_Φ (for instance, if $v \in \mathcal{E}$, and any $u \in H_\Phi$), we have :

$$(J_{\widehat{\Phi}} a_\Phi(\omega_{u,v}) | \Lambda_\Phi(x)) = (R(x^*) \delta^{1/2} v | u) = \langle x^*, \omega_{\delta^{1/2} v, u} \circ R \rangle$$

and, therefore, $\omega_{\delta^{1/2} v, u} \circ R$ belongs to I_Φ , and $a_\Phi(\omega_{\delta^{1/2} v, u} \circ R) = J_{\widehat{\Phi}} a_\Phi(\omega_{u,v})$.

Proof. We have, using 3.10(v), 3.9 and 4.1 :

$$\begin{aligned} (J_{\widehat{\Phi}} a_\Phi(\omega_{u,v}) | \Lambda_\Phi(x e_n)) &= (J_{\widehat{\Phi}} \Lambda_\Phi(x e_n) | a_\Phi(\omega_{u,v})) \\ &= (\Lambda_\Phi(R(x^*) \overline{e_n \delta^{1/2}})) | a_\Phi(\omega_{u,v})) \\ &= (R(x^*) \overline{e_n \delta^{1/2} v} | u) \end{aligned}$$

which is the first result. The second result is clear by continuity. \square

4.3. Lemma. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let \mathcal{E} be the subspace introduced in 3.10; then :*

- (i) $\mathcal{E} \subset \{\xi \in D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}(\delta^{1/2}), \delta^{1/2} \xi \in D((H_\Phi)_\beta, \nu^o)\}$
- (ii) $\mathcal{E} \subset \{\xi \in D((H_\Phi)_\beta, \nu^o) \cap \mathcal{D}(\delta^{1/2}), \delta^{1/2} \xi \in D({}_\alpha H_\Phi, \nu)\}$

Proof. Let us recall that \mathcal{E} is the linear subspace generated by all the elements of the form $J_\Phi \Lambda_\Phi(e_n x)$, where the self-adjoint elements e_n had been introduced in 2.6, and x belongs to the subset \mathcal{T} of elements in $\mathfrak{N}_T \cap \mathfrak{N}_{RTR} \cap \mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$ which are analytic with respect both to Φ and $\Phi \circ R$, and such that, for any z, z' and \mathbb{C} , $\sigma_z^\Phi \circ \sigma_{z'}^{\Phi \circ R}(x)$ belongs to $\mathfrak{N}_T \cap \mathfrak{N}_{RTR} \cap \mathfrak{N}_\Phi \cap \mathfrak{N}_{\Phi \circ R}$. Using then the identifications made in 2.6, we have :

$$J_\Phi \Lambda_\Phi(e_n x) = \delta^{-1/2} J_{\Phi \circ R} \Lambda_{\Phi \circ R}(e_n x)$$

and, therefore, $J_\Phi \Lambda_\Phi(e_n x)$ belongs to $D(\delta^{1/2})$ and $\delta^{1/2} J_\Phi \Lambda_\Phi(e_n x) = J_{\Phi \circ R} \Lambda_{\Phi \circ R}(e_n x)$ which belongs to $D((H_\Phi)_\beta, \nu^o)$. So we have (i).

On the other hand, for all $t \in \mathbb{R}$, as $\delta^{-t} x \delta^t \subset \sigma_{-it}^\Phi \sigma_{it}^{\Phi \circ R}(x)$ is bounded, we get that $e_n x \delta^t \subset \overline{e_n \delta^t \delta^{-t} x \delta^t}$ is bounded also, and belongs to \mathfrak{N}_Φ , and that :

$$J_\Phi \Lambda_\Phi(\overline{e_n x \delta^t}) = \delta^t J_\Phi \Lambda_\Phi(\lambda^{-it/2} e_n x)$$

So, $\delta^t J_\Phi \Lambda_\Phi(e_n x) = J_\Phi \Lambda_\Phi(\lambda^{-it/2} \overline{e_n x \delta^t})$ which belongs to $D({}_\alpha H_\Phi, \nu)$. So, we get (ii). \square

4.4. Definitions and lemma. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let \mathcal{T} and \mathcal{E} be the subspaces introduced in 3.10; let us define \mathcal{T}_τ as the subspace of \mathcal{T} made of elements which are analytic with respect to τ_t , and such that, for any $z \in \mathbb{C}$, $\tau_z(x)$ belongs to \mathcal{T} , and \mathcal{E}_τ as the linear subspace of \mathcal{E} generated by the elements of the form $J_\Phi \Lambda_\Phi(e_n x)$, where the self-adjoint elements e_n had been introduced in 2.6, and x belongs to the subset \mathcal{T}_τ . Then :*

- (i) \mathcal{E}_τ is dense in H_Φ ; moreover, for any ξ in $D({}_\alpha H_\Phi, \nu)$, there exists ξ_n in \mathcal{E}_τ such that $R^{\alpha, \nu}(\xi_n)$ weakly converges to $R^{\alpha, \nu}(\xi)$.

(ii) $\mathcal{E}_\tau \subset \{\xi \in \mathcal{D}(\hat{\delta}^{1/2}), \hat{\delta}^{1/2}\xi \in D((H_\Phi)_{\hat{\beta}}, \nu^o)\}$.

(iii) for all s, t in \mathbb{R} , $\xi \in \mathcal{E}_\tau$, ξ belongs to $\mathcal{D}((\delta\Delta_{\hat{\Phi}})^t)$, and $(\delta\Delta_{\hat{\Phi}})^t\xi$ belongs to \mathcal{E}_τ ; moreover, ξ belongs also to $\mathcal{D}(P^t\delta^s)$, and $P^t\delta^s\xi$ belongs to \mathcal{E}_τ .

(iv) let us define $\hat{\mathcal{E}}_{\hat{\tau}}$ the linear subspace associated to the dual measured quantum groupoid $\widehat{\mathfrak{G}}$. Then :

$$\hat{\mathcal{E}}_{\hat{\tau}} \subset \{\xi \in D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}(\delta^{1/2}), \delta^{1/2}\xi \in D((H_\Phi)_\beta, \nu^o)\}$$

Proof. There exists an invertible positive operator q affiliated to $Z(N)$ such that $\lambda = \alpha(q) = \beta(q)$ (3.8(vi)). Let us write $q = \int_0^\infty \mu de_\mu$, and let $f_r = \int_{1/r}^r de_\mu$.

Let $x \in \mathcal{T}$; as the automorphisms groups τ_t , σ_t^Φ and $\sigma_t^{\Phi \circ R}$ are two by two commuting, we can check that, for any $x \in \mathcal{T}$, the operators :

$$x_{n,r} = \sqrt{\frac{n}{\pi}} \int_{-\infty}^\infty e^{-nt^2} f_r \tau_t(x) dt$$

belong to \mathcal{T}_τ , the strong limit $\lim_{n,r} x_{n,r} = x$, and $\lim_{n,r} \Lambda_\Phi(x_{n,r}) = \Lambda_\Phi(x)$. From which we get that \mathcal{E}_τ is dense in \mathcal{E} , and, therefore, in H_Φ ; moreover, as we can prove that all the operators $T(x_{n,r}^* e_p^2 x_{n,r})$ are uniformly bounded, we get also that $R^{\alpha,\nu}(J_\Phi \Lambda_\Phi(e_p x_{n,r}))$ is strongly converging to $R^{\alpha,\nu}(J_\Phi \Lambda_\Phi(e_p x))$, from which we get (i).

Let now x be in \mathcal{T}_τ . Then, using 3.11(ii), we get that :

$$\begin{aligned} \hat{\delta}^{1/2} J_\Phi \Lambda_\Phi(e_n x) &= P^{-1/2} J_\Phi \delta^{1/2} J_\Phi \delta^{-1/2} \Delta_\Phi^{-1/2} J_\Phi \Lambda_\Phi(e_n x) \\ &= P^{-1/2} \lambda^{i/4} J_\Phi \Lambda_\Phi(\sigma_{-i/2}(e_n) \sigma_{-i/2}^{\Phi \circ R}(x)) \\ &= \lambda^{i/4} J_\Phi \Lambda_\Phi(\sigma_{-i/2}(e_n) \tau_{-i/2}(\sigma_{-i/2}^{\Phi \circ R}(x))) \end{aligned}$$

We get also that $J_\Phi \delta^{1/2} J_\Phi \Lambda_\Phi(e_n x) = \lambda^{-i/4} J_\Phi \Lambda_\Phi(\sigma_{-i/2} \tau_{i/2} \sigma_{i/2}^{\Phi \circ R}(x^*) e_n)$, which belongs to $D({}_\alpha H_\Phi, \nu)$, and, therefore, $\hat{\delta}^{1/2} J_\Phi \Lambda_\Phi(e_n x)$ belongs to $D((H_\Phi)_{\hat{\beta}}, \nu^o)$, which is (ii).

If x belongs to \mathcal{T}_τ , and $t \in \mathbb{R}$, we get that :

$$P^t J_\Phi \Lambda_\Phi(e_n x) = J_\Phi(\lambda^{-t/2} e_n \tau_{-it}(x)) \in \mathcal{E}_\tau$$

and, using 3.8(vii) :

$$(\delta\Delta_{\hat{\Phi}})^t J_\Phi \Lambda_\Phi(e_n x) = J_\Phi \Lambda_\Phi(\lambda^{-it} e_n \sigma_{-it}^\Phi \sigma_{it}^{\Phi \circ R} \tau_{-it}(x))$$

and, therefore, $\mathcal{E}_\tau \subset D((\delta\Delta_{\hat{\Phi}})^t)$, and $(\delta\Delta_{\hat{\Phi}})^t \mathcal{E}_\tau \subset \mathcal{E}_\tau$

Moreover, we have also :

$$\delta^s J_\Phi \Lambda_\Phi(e_n x) = J_\Phi \Lambda_\Phi(\lambda^{-is/2} e_n x \delta^s) \in \mathcal{E}_\tau$$

from which we get (iii).

If we consider now the dual measured quantum groupoid $\widehat{\mathfrak{G}}$, and the linear subset $\widehat{\mathcal{E}}_{\hat{\tau}}$ associated with, we get, using (iii), that $\delta^{1/2} \widehat{\mathcal{E}}_{\hat{\tau}} \subset D((H_\Phi)_\beta, \nu^o)$, which is (iv). \square

4.5. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7, \mathfrak{G}^o the opposite measured quantum groupoid in the sense of 3.12(i), η in $D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}(\delta^{1/2})$ such that $\delta^{1/2} \eta$ belongs to $D((H_\Phi)_\beta, \nu^o)$, $(e_i)_{i \in I}$ an (α, ν) -orthonormal basis of H_Φ ; then, for all $x \in \mathfrak{N}_\Phi$, $\xi \in D({}_\alpha H_\Phi, \nu)$, $(id_{\beta * \alpha} \omega_{\eta, \xi}) \Gamma(x)$ belongs to \mathfrak{N}_Φ , and

the pseudo-multiplicative \widehat{W}^o satisfies :

$$\widehat{W}^o(\Lambda_\Phi(x) \underset{\nu^o}{\hat{\alpha} \otimes \beta} \delta^{1/2} \eta) = \sum_i \Lambda_\Phi[(id_{\beta * \alpha} \omega_{\eta, e_i}) \Gamma(x)] \underset{\nu}{\beta \otimes \alpha} e_i$$

$$\begin{aligned}
\Lambda_\Phi[(id \underset{N}{\beta^* \alpha} \omega_{\eta, \xi}) \Gamma(x)] &= (id * \omega_{\delta^{1/2} \eta, \xi})(\widehat{W}^o) \Lambda_\Phi(x) \\
&= J_{\widehat{\Phi}} \widehat{\pi}(\omega_{J_{\widehat{\Phi}} \xi, J_{\widehat{\Phi}} \delta^{1/2} \eta})^* J_{\widehat{\Phi}} \Lambda_\Phi(x)
\end{aligned}$$

If u belongs to $D((H_\Phi)_\beta, \nu^o)$ and v belongs to $D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}(\delta^{1/2})$ such that $\delta^{1/2}v$ belongs to $D((H_\Phi)_\beta, \nu^o)$, we get that $(id \underset{N}{\beta^* \alpha} \omega_{\delta^{1/2}u, v} \circ R) \Gamma(x)$ belongs to \mathfrak{N}_Φ , and :

$$\Lambda_\Phi((id \underset{N}{\beta^* \alpha} \omega_{u, \delta^{1/2}v} \circ R) \Gamma(x)) = J_{\widehat{\Phi}} \widehat{\pi}(\omega_{u, v})^* J_{\widehat{\Phi}} \Lambda_\Phi(x)$$

Proof. Let's apply the definition of W^* (3.6) to the measured quantum groupoid \mathfrak{G}^o , with the identification made in 3.9 of $\Lambda_\Phi(xe_n)$ with $\Lambda_{\Phi \circ R}(xe_n \delta^{-1/2})$ (where e_n has been defined in 2.6) :

$$\widehat{W}^o(\Lambda_\Phi(xe_n) \underset{\nu^o}{\hat{\alpha} \otimes \beta} \delta^{1/2} \eta) = \sum_i \Lambda_\Phi[(id \underset{N}{\beta^* \alpha} \omega_{\eta, e_i}) \Gamma(xe_n)] \underset{\nu}{\beta \otimes \alpha} e_i$$

from which we get :

$$\Lambda_\Phi[(id \underset{N}{\beta^* \alpha} \omega_{\eta, \xi}) \Gamma(xe_n)] = (id * \omega_{\delta^{1/2} \eta, \xi})(\widehat{W}^o) \Lambda_\Phi(xe_n)$$

As Λ_Φ is σ -strong $*$ -norm closed, we get that $(id \underset{N}{\beta^* \alpha} \omega_{\eta, \xi}) \Gamma(x)$ belongs to \mathfrak{N}_Φ , and that :

$$\Lambda_\Phi[(id \underset{N}{\beta^* \alpha} \omega_{\eta, \xi}) \Gamma(x)] = (id * \omega_{\delta^{1/2} \eta, \xi})(\widehat{W}^o) \Lambda_\Phi(x)$$

Now, we get easily the first formula. The second equality is just an application of 3.10(ii) and 3.12(v). Then, writing $u = J_{\widehat{\Phi}} \delta^{1/2} \eta$, and $v = J_{\widehat{\Phi}} \xi$, and using 3.10(v) and 3.8(vi), we get the last formula. \square

4.6. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid; let ξ in $D((H_\Phi)_\beta, \nu^o)$, η in $D({}_\alpha H_\Phi, \nu)$ such that $\omega_{\xi, \eta}$ belongs to I_Φ ; then, the operator $\widehat{\pi}(\omega_{\xi, \eta})$ belongs to $\mathfrak{N}_{\widehat{\Phi}}$, and we have then :

$$\Lambda_{\widehat{\Phi}}(\widehat{\pi}(\omega_{\xi, \eta})) = a(\omega_{\xi, \eta})$$

Proof. Let $\xi \in D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$, $\eta \in D({}_\alpha H_\Phi, \nu)$ such that $\omega_{\xi, \eta}$ belongs to I_Φ ; let $u \in D({}_\alpha H_\Phi, \nu)$, $v \in \mathcal{D}(\delta^{1/2})$ such that $\delta^{1/2}v$ is in $D((H_\Phi)_\beta, \nu)$ and $\omega_{u, v}$ is in I_Φ ; we have then, for any $x \in \mathfrak{N}_\Phi$, using successively 3.10(ii), 4.2 and 4.5 :

$$\begin{aligned}
(\widehat{\pi}(\omega_{\xi, \eta}) J_{\widehat{\Phi}} a_\Phi(\omega_{u, v}) | \Lambda_\Phi(x)) &= (J_{\widehat{\Phi}} a_\Phi(\omega_{u, v}) | \widehat{\pi}(\omega_{\xi, \eta})^* \Lambda_\Phi(x)) \\
&= (J_{\widehat{\Phi}} a_\Phi(\omega_{u, v}) | \Lambda_\Phi((\omega_{\eta, \xi} \underset{N}{\beta^* \alpha} id) \Gamma(x))) \\
&= (a_\Phi(\omega_{\delta^{1/2}v, u} \circ R) | \Lambda_\Phi((\omega_{\eta, \xi} \underset{N}{\beta^* \alpha} id) \Gamma(x))) \\
&= \langle [(\omega_{\eta, \xi} \underset{N}{\beta^* \alpha} id) \Gamma(x)]^*, \omega_{\delta^{1/2}v, u} \circ R \rangle \\
&= \langle (\omega_{\xi, \eta} \underset{N}{\beta^* \alpha} id) \Gamma(x^*), \omega_{\delta^{1/2}v, u} \circ R \rangle \\
&= \langle [(id \underset{N}{\beta^* \alpha} \omega_{u, \delta^{1/2}v} \circ R) \Gamma(x)]^*, \omega_{\xi, \eta} \rangle \\
&= (a_\Phi(\omega_{\xi, \eta}) | \Lambda_\Phi((id \underset{N}{\beta^* \alpha} \omega_{u, \delta^{1/2}v} \circ R) \Gamma(x))) \\
&= (a_\Phi(\omega_{\xi, \eta}) | J_{\widehat{\Phi}} \widehat{\pi}(\omega_{u, v})^* J_{\widehat{\Phi}} \Lambda_\Phi(x)) \\
&= (J_{\widehat{\Phi}} \widehat{\pi}(\omega_{u, v}) J_{\widehat{\Phi}} a_\Phi(\omega_{\xi, \eta}) | \Lambda_\Phi(x))
\end{aligned}$$

From which we get, by density :

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}a_{\Phi}(\omega_{u,v}) = J_{\widehat{\Phi}}\hat{\pi}(\omega_{u,v})J_{\widehat{\Phi}}a_{\Phi}(\omega_{\xi,\eta})$$

Let us take $v = J_{\Phi}\Lambda_{\Phi}(e_n x)$, with e_n defined in 3.9 and $x \in \mathcal{T}$ (3.10). Then, this formula gives :

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}J_{\Phi}(e_n x)^*J_{\Phi}u = J_{\widehat{\Phi}}\hat{\pi}(\omega_{u,J_{\Phi}\Lambda_{\Phi}(e_n x)})J_{\widehat{\Phi}}a_{\Phi}(\omega_{\xi,\eta})$$

which, by continuity, gives, for any $u \in D({}_{\alpha}H_{\Phi}, \nu)$, $v \in J_{\Phi}\Lambda_{\Phi}(\mathfrak{N}_{\Phi}) \cap D((H_{\Phi})_{\beta}, \nu^o)$:

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}\pi'(v)^*u = J_{\widehat{\Phi}}\hat{\pi}(\omega_{u,v})J_{\widehat{\Phi}}a_{\Phi}(\omega_{\xi,\eta})$$

Let us suppose now that u, v belong to $D({}_{\alpha}H_{\Phi}, \nu) \cap D((H_{\Phi})_{\beta}, \nu^o)$ and are such that $\omega_{u,v}$ belongs to I_{Φ} ; with the notations of 4.1, let us write :

$$\mathcal{P}_v = \int_0^{\infty} \lambda de_{\lambda} + (1-p)\infty$$

As $\omega_{u,v}$ belongs to I_{Φ} , we get that $pu = u$. Let us define $v_{\mu} = e_{\mu}v$ which belongs to $J_{\Phi}\Lambda_{\Phi}(\mathfrak{N}_{\Phi}) \cap D((H_{\Phi})_{\beta}, \nu^o)$. We have :

$$\lim_{\mu} \pi'(v_{\mu})^*u = \lim_{\mu} \pi'(v)^*e_{\mu}u = \pi'(v)^*pu = \pi'(v)^*u = a_{\Phi}(\omega_{u,v})$$

and we get also, for the weak limits :

$$\lim_{\mu} R^{\beta, \nu^o}(v_{\mu}) = \lim_{\mu} e_{\mu} R^{\beta, \nu^o}(v) = p R^{\beta, \nu^o}(v) = R^{\beta, \nu^o}(pv)$$

and, as $p \in M'$, we get for the weak limits :

$$\begin{aligned} \lim_{\mu} \hat{\pi}(\omega_{u,v_{\mu}}) &= \lim_{\mu} (\omega_{u,v_{\mu}} * id)(W) = (\omega_{u,pv} * id)(W) = \\ &= (\omega_{pv,v} * id)(W) = (\omega_{u,v} * id)(W) = \hat{\pi}(\omega_{u,v}) \end{aligned}$$

and, taking the limits of the equality :

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}\pi'(v_{\mu})^*u = J_{\widehat{\Phi}}\hat{\pi}(\omega_{u,v_{\mu}})J_{\widehat{\Phi}}a_{\Phi}(\omega_{\xi,\eta})$$

we get :

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}a_{\Phi}(\omega_{u,v}) = J_{\widehat{\Phi}}\hat{\pi}(\omega_{u,v})J_{\widehat{\Phi}}a_{\Phi}(\omega_{\xi,\eta})$$

for all u, v in $D({}_{\alpha}H_{\Phi}, \nu) \cap D((H_{\Phi})_{\beta}, \nu^o)$ and are such that $\omega_{u,v}$ belongs to I_{Φ} ; therefore, by linearity, we get, for all $\omega \in M_{*}^{\alpha, \beta} \cap I_{\Phi}$, that :

$$\hat{\pi}(\omega_{\xi,\eta})J_{\widehat{\Phi}}a(\omega) = J_{\widehat{\Phi}}\hat{\pi}(\omega)J_{\widehat{\Phi}}a(\omega_{\xi,\eta}) = J_{\widehat{\Phi}}\hat{\pi}(\omega)J_{\widehat{\Phi}}\pi'(\eta)^*\xi$$

By the closedness of $\pi'(\eta)_{|K_{\eta}}^*$ (4.1), this formula remains true for ξ in $D((H_{\Phi})_{\beta}, \nu^o)$ such that $\omega_{\xi,\eta}$ belongs to I_{Φ} . From which we get that $a(\omega_{\xi,\eta})$ is left-bounded with respect to $\widehat{\Phi}$, and the result, by standard arguments of Tomita-Takesaki's theory. \square

4.7. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid; let's take ξ in $D((H_{\Phi})_{\beta}, \nu^o)$ and η in $D({}_{\alpha}H_{\Phi}, \nu)$, and let us suppose that $\hat{\pi}(\omega_{\xi,\eta})$ belongs to $\mathfrak{N}_{\widehat{\Phi}}$. Then $\omega_{\xi,\eta}$ belongs to I_{Φ} and :*

$$\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi,\eta})) = a_{\Phi}(\omega_{\xi,\eta})$$

Proof. Let u, v be in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$; then the linear form $\hat{\omega}_{u,v}$ on \widehat{M} defined, for y in \widehat{M} by $\hat{\omega}_{u,v}(y) = (yu|v)$ belongs to $\widehat{M}_*^{\alpha, \beta}$ (with the notations of 3.10(i) applied to $\widehat{\mathfrak{G}}$). Let us suppose that $\hat{\omega}_{u,v}$ belongs to $I_{\widehat{\Phi}}$. Using 3.10(v) and (ii), applied to $\widehat{\mathfrak{G}}$, we get :

$$\begin{aligned} (a_{\widehat{\Phi}}(\hat{\omega}_{u,v})|\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi, \eta}))) &= \langle \hat{\pi}(\omega_{\xi, \eta})^*, \hat{\omega}_{u,v} \rangle \\ &= ((\omega_{\xi, \eta} * id)(W)^* u|v) \\ &= (\eta \otimes_{\beta} u | W(\xi \otimes_{\alpha} v)) \\ &= (\widehat{W}(u \otimes_{\beta} \eta) | v \otimes_{\alpha} \xi) \\ &= (\hat{\pi}(\hat{\omega}_{u,v})\eta | \xi) \end{aligned}$$

and, by linearity, we obtain, for all $\hat{\omega}$ in $\widehat{M}_*^{\alpha, \beta} \cap I_{\widehat{\Phi}}$:

$$(\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi, \eta}))|a_{\widehat{\Phi}}(\hat{\omega})) = \langle \hat{\pi}(\hat{\omega})^*, \omega_{\xi, \eta} \rangle$$

But, by 3.10(v) applied to $\widehat{\mathfrak{G}}$, and 3.11(i), we have $a_{\widehat{\Phi}}(\hat{\omega}) = \Lambda_{\widehat{\Phi}}(\hat{\pi}(\hat{\omega}))$, and, by 3.10(v) applied to $\widehat{\mathfrak{G}}$ and again 3.11(i), we know that the algebra $\hat{\pi}(\widehat{M}_*^{\alpha, \beta} \cap I_{\widehat{\Phi}})$ is a core for $\Lambda_{\widehat{\Phi}}$; so, by the closedness of $\Lambda_{\widehat{\Phi}}$, we get, for all $x \in \mathfrak{N}_{\widehat{\Phi}}$:

$$(\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi, \eta}))|\Lambda_{\widehat{\Phi}}(x)) = \langle x^*, \omega_{\xi, \eta} \rangle$$

from which we get that $|\langle x^*, \omega_{\xi, \eta} \rangle| \leq \|\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi, \eta}))\| \|\Lambda_{\widehat{\Phi}}(x)\|$, and that $\omega_{\xi, \eta}$ belongs to $I_{\widehat{\Phi}}$. Then, using 3.10(v), the proof is finished. \square

4.8. Theorem. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid; let's take ξ in $D((H_\Phi)_\beta, \nu^o)$ and η in $D({}_\alpha H_\Phi, \nu)$; then, are equivalent :

(i) $\hat{\pi}(\omega_{\xi, \eta})$ belongs to $\mathfrak{N}_{\widehat{\Phi}}$;

(ii) $\omega_{\xi, \eta}$ belongs to I_Φ .

We have then $\Lambda_{\widehat{\Phi}}(\hat{\pi}(\omega_{\xi, \eta})) = a_\Phi(\omega_{\xi, \eta})$.

Proof. This is just 4.6 and 4.7. \square

4.9. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid; for any $X \in \hat{\alpha}(N)'$, let us define :

$$\mathfrak{S}(X) = (id \otimes_{\beta} \alpha \otimes_{\widehat{\Phi}})(W^c(X \otimes_{\alpha} 1)W^{c*})$$

Then :

(i) $\mathfrak{S}(X)$ belongs to the positive extended part of $\hat{\alpha}(N)'$, and, for any $\xi \in D((H_\Phi)_\beta, \nu^o)$, $\eta \in D({}_\alpha H_\Phi, \nu)$, we have :

$$\langle \mathfrak{S}(\theta^{\hat{\alpha}}(J_{\widehat{\Phi}} J_\Phi \eta, J_{\widehat{\Phi}} J_\Phi \eta)), \omega_{J_{\widehat{\Phi}} J_\Phi \xi} \rangle = \widehat{\Phi}(\hat{\pi}(\omega_{\xi, \eta})^* \hat{\pi}(\omega_{\xi, \eta})) = \langle \mathcal{P}_\eta, \omega_\xi \rangle$$

(i) for all $\eta \in D({}_\alpha H_\Phi, \nu)$, we have :

$$\mathfrak{S}(\theta^{\hat{\alpha}}(J_{\widehat{\Phi}} J_\Phi \eta, J_{\widehat{\Phi}} J_\Phi \eta)) = J_{\widehat{\Phi}} J_\Phi \mathcal{P}_\eta J_\Phi J_{\widehat{\Phi}}$$

(ii) $\mathfrak{S}(X)$ belongs to the positive extended part of M .

Proof. Using 3.12(v), we get that $W^{c*} = (J_\Phi J_{\widehat{\Phi}} \otimes_{\beta} 1)W(J_{\widehat{\Phi}} J_\Phi \otimes_{\alpha} 1)$; so, with the intertwining properties of W , we easily get that $\mathfrak{S}(X)$ is an element of the extended positive part of $\hat{\alpha}(N)'$.

Let us notice that $J_{\widehat{\Phi}}J_{\Phi}\xi$ belongs to $D((H_{\Phi})_{\widehat{\beta}}, \nu^o)$, and $J_{\widehat{\Phi}}J_{\Phi}\eta$ to $D(\widehat{\alpha}H_{\Phi}, \nu)$. We then get that $\langle \mathcal{S}(\theta^{\widehat{\alpha}}(J_{\widehat{\Phi}}J_{\Phi}\eta, J_{\widehat{\Phi}}J_{\Phi}\eta)), \omega_{J_{\widehat{\Phi}}J_{\Phi}\xi} \rangle$ is equal to :

$$\begin{aligned} \widehat{\Phi}((\omega_{J_{\widehat{\Phi}}J_{\Phi}\xi, J_{\widehat{\Phi}}J_{\Phi}\eta} * id)(W^{c*})^*(\omega_{J_{\widehat{\Phi}}J_{\Phi}\xi, J_{\widehat{\Phi}}J_{\Phi}\eta} * id)(W^{c*})) &= \\ &= \widehat{\Phi}((\omega_{\xi, \eta} * id)(W)^*(\omega_{\xi, \eta} * id)(W)) = \widehat{\Phi}(\widehat{\pi}(\omega_{\xi, \eta})^* \widehat{\pi}(\omega_{\xi, \eta})) \end{aligned}$$

If it is finite, it is equal to $\|\Lambda_{\widehat{\Phi}}(\widehat{\pi}(\omega_{\xi, \eta}))\|^2 = \|a_{\Phi}(\omega_{\xi, \eta})\|^2 = \|\omega_{\xi, \eta}\|_{\Phi}^2$, thanks to 4.8. If it is infinite, using 4.8 again, $\|\omega_{\xi, \eta}\|_{\Phi}$ also is infinite. So, in both cases, it is equal to $\|\omega_{\xi, \eta}\|_{\Phi}^2$, which is equal, using 4.1, to $\langle \mathcal{P}_{\eta}, \omega_{\xi} \rangle$, which finishes the proof of (i). Then (ii) (resp. (iii)) is a straightforward corollary of (i) (resp. (ii)). \square

4.10. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid. Let X be a positive element in \widehat{M} ; then, we have:*

$$(id_{\beta * \alpha} \widehat{\Phi}) \widehat{\Gamma}(X) = \widehat{T}(X)$$

Proof. Using 3.12(v), we get that :

$$\widehat{\Gamma}(X) = W^c(X \underset{N^o}{\widehat{\alpha} \otimes \widehat{\beta}} 1) W^{c*}$$

and, therefore, with the notations of 4.9, $(id_{\beta * \alpha} \widehat{\Phi}) \widehat{\Gamma}(X) = \mathcal{S}(X)$. Using 4.9(iii), it belongs to the positive extended part of M , and, by construction, it belongs to the positive extended part of \widehat{M} . So, it belongs to the positive extended part of $\alpha(N)$. If u is a unitary in N , we easily get that $\mathcal{S}(\alpha(u^*)X\alpha(u)) = \alpha(u^*)\mathcal{S}(X)\alpha(u)$, and, therefore, the application $X \rightarrow \mathcal{S}(X)$ is an operator-valued weight from \widehat{M} on $\alpha(N)$. It is clearly normal and faithful, and, as $\mathcal{S}(X) = \widehat{T}(X)$ for any $X \in \mathfrak{M}_{\widehat{T}}^+$, we get it is also semi-finite, and that, for any positive X in \widehat{M} , we have $\mathcal{S}(X) \leq \widehat{T}(X)$.

Let us define $\tilde{\Phi}(X) = \nu \circ \alpha^{-1} \circ \mathcal{S}(X)$; we have $\tilde{\Phi} \leq \widehat{\Phi}$, and these two weights are equal on $\mathfrak{M}_{\widehat{T}}$; moreover, as, for all $t \in \mathbb{R}$, we have :

$$\widehat{\Gamma}\sigma_t^{\widehat{\Phi}} = (\widehat{\tau}_t \underset{N}{\beta * \alpha} \sigma_t^{\widehat{\Phi}}) \widehat{\Gamma}$$

we get that $\mathcal{S}\sigma_t^{\widehat{\Phi}} = \widehat{\tau}_t \mathcal{S}$, and that $\tilde{\Phi}$ is invariant under $\sigma_t^{\widehat{\Phi}}$, by 3.8(ii) applied to $\widehat{\mathfrak{G}}$; therefore, we have $\tilde{\Phi} = \widehat{\Phi}$, from which we deduce that $\mathcal{S} = \widehat{T}$, which gives the result. \square

4.11. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid, \mathfrak{H} an Hilbert space with b a faithful normal anti-representation of N on \mathfrak{H} . Let X be a positive element in $\mathcal{L}(\mathfrak{H})_{b * \alpha} \widehat{M}$; then, we have:*

$$(id_{b * \alpha} id_{\beta * \alpha} \widehat{\Phi})(id_{b * \alpha} \widehat{\Gamma})(X) = (id_{b * \alpha} \widehat{\Phi})(X) \underset{N}{b \otimes \alpha} 1$$

Proof. Using 3.12(v), we get that, for $X \in \mathcal{L}(\mathfrak{H})_{b * \alpha} \widehat{M}$:

$$(id_{b * \alpha} \widehat{\Gamma})(X) = (1 \underset{N}{b \otimes \alpha} W^c)(X \underset{N^o}{\widehat{\alpha} \otimes \widehat{\beta}} 1)(1 \underset{N}{b \otimes \alpha} W^c)^*$$

For any Y in $\mathcal{L}(\mathfrak{H})_{b \otimes \alpha} H_{\Phi}^+$, commuting with $1 \underset{N}{b \otimes \alpha} \widehat{\alpha}(N)$, let us write

$$\tilde{\mathcal{S}}(Y) = (id_{b * \alpha} id_{\beta * \alpha} \widehat{\Phi})[(1 \underset{N}{b \otimes \alpha} W^c)(Y \underset{N^o}{\widehat{\alpha} \otimes \widehat{\beta}} 1)(1 \underset{N}{b \otimes \alpha} W^c)^*]$$

which is an element of the positive extended part of $\mathcal{L}(\mathfrak{H}_{b \otimes_\alpha H_\Phi})$.

Let's take ξ in $D((\mathfrak{H}_{b \otimes_\alpha H_\Phi})_{1_{b \otimes_\alpha \hat{\beta}}}, \nu^o)$ and η in $D(1_{b \otimes_\alpha \hat{\alpha}}(\mathfrak{H}_{b \otimes_\alpha H_\Phi})_{\nu}, \nu)$. Then, we get that $\langle S(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta)), \omega_\xi \rangle$ is equal to :

$$\widehat{\Phi}[(\omega_{\xi, \hat{\beta} * \alpha} id)((1_{b \otimes_\alpha} W^c)(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta)_{\hat{\alpha} \otimes_{\hat{\beta}} 1})(1_{b \otimes_\alpha} W^c)^*])]$$

which can be written also :

$$\widehat{\Phi}[(\omega_{\xi, \eta} * id)((1_{b \otimes_\alpha} W^{c*}))^*(\omega_{\xi, \eta} * id)((1_{b \otimes_\alpha} W^{c*}))]$$

Using now 3.12(v), we get that $W^{c*} = (J_\Phi J_{\hat{\Phi}} \alpha \otimes_{\hat{\beta}} 1)W(J_{\hat{\Phi}} J_\Phi \hat{\beta} \otimes_\alpha 1)$ and, therefore,

$\langle S(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta)), \omega_\xi \rangle$ is equal to $\widehat{\Phi}(A^*A)$, where :

$$A = (\omega_{(1_{b \otimes_\alpha} J_{\hat{\Phi}} J_\Phi) \xi, (1_{b \otimes_\alpha} J_{\hat{\Phi}} J_\Phi) \eta} * id)(1_{b \otimes_\alpha} W)$$

Let $(e_i)_{i \in I}$ be a (b, ν^o) -orthogonal basis of \mathfrak{H} . Using ([E3], 2.3(ii)), there exist ξ_i in $D((H_\Phi)_{\hat{\beta}}, \nu^o)$ and η_i in $D(\hat{\alpha} H_\Phi, \nu)$ such that :

$$\xi = \sum_i e_i b \otimes_\alpha \xi_i$$

$$\eta = \sum_i e_i b \otimes_\alpha \eta_i$$

and we get that $A = \sum_i \alpha(\langle e_i, e_i \rangle_{b, \nu^o}) (\omega_{J_{\hat{\Phi}} J_\Phi \xi_i, J_{\hat{\Phi}} J_\Phi \eta_i} * id)(W)$ because $J_{\hat{\Phi}} J_\Phi \xi_i$ belongs to $D((H_\Phi)_{\hat{\beta}}, \nu^o)$, and $J_{\hat{\Phi}} J_\Phi \eta_i$ belongs to $D(\alpha H_\Phi, \nu)$; so, $\langle S(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta)), \omega_\xi \rangle$ is equal to :

$$\sum_{i,j} \widehat{\Phi}(\hat{\pi}(\omega_{J_{\hat{\Phi}} J_\Phi \xi_i, J_{\hat{\Phi}} J_\Phi \eta_i})^* \alpha(\langle e_i, e_i \rangle_{b, \nu^o} \langle e_j, e_j \rangle_{b, \nu^o}) \hat{\pi}(\omega_{J_{\hat{\Phi}} J_\Phi \xi_j, J_{\hat{\Phi}} J_\Phi \eta_j}))$$

If it is bounded, then so are, for all $i \in I$:

$$\widehat{\Phi}(\hat{\pi}(\omega_{J_{\hat{\Phi}} J_\Phi \xi_i, J_{\hat{\Phi}} J_\Phi \eta_i})^* \alpha(\langle e_i, e_i \rangle_{b, \nu^o}) \hat{\pi}(\omega_{J_{\hat{\Phi}} J_\Phi \xi_i, J_{\hat{\Phi}} J_\Phi \eta_i}))$$

which implies (4.7) that, for all $i \in I$, $\omega_{J_{\hat{\Phi}} J_\Phi \xi_i, J_{\hat{\Phi}} J_\Phi \eta_i}$ belongs to I_Φ , and, therefore, using 4.1, that $J_{\hat{\Phi}} J_\Phi \xi_i$ belongs to $D(\pi'(J_{\hat{\Phi}} J_\Phi \eta_i)^*)$; moreover, we then get :

$$\begin{aligned} \langle S(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta)), \omega_\xi \rangle &= \left\| \sum_i \alpha(\langle e_i, e_i \rangle_{b, \nu^o}) \pi'(J_{\hat{\Phi}} J_\Phi \eta_i)^* J_{\hat{\Phi}} J_\Phi \xi_i \right\|^2 \\ &= \left\| \sum_i e_i b \otimes_\alpha \pi'(J_{\hat{\Phi}} J_\Phi \eta_i)^* J_{\hat{\Phi}} J_\Phi \xi_i \right\|^2 \end{aligned}$$

By definition of the extended positive part, there exists a closed subspace $K \subset \mathfrak{H}_{b \otimes_\alpha H_\Phi}$

on which $S(\theta^{1_{b \otimes_\alpha \hat{\alpha}}}(\eta, \eta))$ is a positive self-adjoint operator, and it is ∞ on the orthogonal of K .

So, $K = \oplus_i K_i$, where the K_i are the closed orthogonal subspaces $e_i b \otimes_\alpha J_\Phi J_{\hat{\Phi}} K_{J_{\hat{\Phi}} J_\Phi \eta_i}$ (with the notations of 4.1).

Let u be a unitary in M' ; we get that each K_i is invariant under $1_{b \otimes_\alpha} u$, and so, we get

that p_K belongs to $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} M}$; as $(1_{b \otimes_{\alpha} u})\xi$ belongs to $D((\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}})_{1_{b \otimes_{\alpha} \hat{\beta}}, \nu^o})$, we get that :

$$\begin{aligned} \langle S(\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(\eta, \eta)), \omega_{u\xi} \rangle &= \left\| \sum_i e_i \otimes_{\nu} \pi'(J_{\hat{\Phi}} J_{\Phi} \eta_i)^* J_{\hat{\Phi}} J_{\Phi} u \xi_i \right\|^2 \\ &= \left\| \sum_i e_i \otimes_{\nu} R(J_{\Phi} u^* J_{\Phi}) \pi'(J_{\hat{\Phi}} J_{\Phi} \eta_i)^* J_{\hat{\Phi}} J_{\Phi} \xi_i \right\|^2 \\ &= \langle S(\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(\eta, \eta)), \omega_{\xi} \rangle \end{aligned}$$

So, we get that $(1_{b \otimes_{\alpha} u^*})S(\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(\eta, \eta))(1_{b \otimes_{\alpha} u})$ is equal to $\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(\eta, \eta)$ from which we get that $S(\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(\eta, \eta))$ belongs to the positive extended part of $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} M}$.

Let $(\eta_j)_{j \in J}$ be an $((1_{b \otimes_{\alpha} \hat{\alpha}}, \nu)$ orthogonal basis of $\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}}$; for any $Y \in \mathcal{L}(\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}})^+$, commuting with $1_{b \otimes_{\alpha} \hat{\alpha}}(N)$. We have :

$$Y = \sum_j \theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(Y^{1/2} \eta_j, Y^{1/2} \eta_j)$$

and, therefore :

$$S(Y) = \sum_j S(\theta^{1_{b \otimes_{\alpha} \hat{\alpha}}}(Y^{1/2} \eta_j, Y^{1/2} \eta_j))$$

from which we get that $S(Y)$ belongs to the positive extended part of $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} M}$.

In particular, if X is a positive element in $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} \widehat{M}}$; then

$$(id_{b^*_{\alpha}} id_{\hat{\beta}^*_{\alpha}} \widehat{\Phi})(id_{b^*_{\alpha}} \widehat{\Gamma})(X) = S(X)$$

belongs to the positive extended part of $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} M}$. As, by definition, it belongs to the positive extended part of $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} \widehat{M}}$, we get that it belongs to the positive extended part of $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha} \alpha(N)}$, from which we get that there exists X' in the positive extended part of $b(N)'$ such that :

$$(id_{b^*_{\alpha}} id_{\hat{\beta}^*_{\alpha}} \widehat{\Phi})(id_{b^*_{\alpha}} \widehat{\Gamma})(X) = X' \otimes_{\alpha} 1$$

Let ξ be in $D(\mathfrak{H}_b, \nu^o)$, η in $D(\alpha H_{\Phi}, \nu) \cap D((H_{\Phi})_{\hat{\beta}}, \nu^o)$; we get :

$$\begin{aligned} \langle b(\langle \eta, \eta \rangle_{\alpha, \nu}) X', \omega_{\xi} \rangle &= (\omega_{\xi} \otimes_{\nu} \omega_{\eta} \otimes_{\nu} \widehat{\Phi})(id_{b^*_{\alpha}} \widehat{\Gamma})(X) \\ &= (\omega_{\eta} \otimes_{\nu} \widehat{\Phi}) \widehat{\Gamma}[(\omega_{\xi} \otimes_{\nu} id)(X)] \end{aligned}$$

which, using 4.10, is equal to :

$$\begin{aligned} \langle \widehat{T}(\omega_{\xi} \otimes_{\nu} id)(X), \omega_{\eta} \rangle &= (\omega_{\xi} \otimes_{\nu} \omega_{\eta})(id_{b^*_{\alpha}} \widehat{T})(X) \\ &= \langle b(\langle \eta, \eta \rangle_{\alpha, \nu})(id_{b^*_{\alpha}} \widehat{\Phi})(X), \omega_{\xi} \rangle \end{aligned}$$

from which we infer that $X' = (id_{b^*_{\alpha}} \widehat{\Phi})(X)$, which gives the result. \square

4.12. **Theorem.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let \mathfrak{H} be an Hilbert space with b a faithful normal anti-representation of N on \mathfrak{H} . Let X be a positive element in M , \tilde{X} a positive element in $\mathcal{L}(\mathfrak{H})_{b^* \alpha_N M}$; then :

- (i) we have $(id_{\beta^* \alpha_N} T) \Gamma(X) = T(X)_{\beta \otimes \alpha_N} 1$;
- (ii) we have $(id_{b^* \alpha_N} id_{\beta^* \alpha_N} T)(id_{b^* \alpha_N} \Gamma)(\tilde{X}) = (id_{b^* \alpha_N} T)(\tilde{X})_{\beta \otimes \alpha_N} 1$

Proof. Let us apply 4.10 and 4.11 to the dual measured quantum groupoid $\widehat{\mathfrak{G}}$. \square

5. COREPRESENTATIONS OF MEASURED QUANTUM GROUPOIDS

In this chapter, we define corepresentations of measured quantum groupoids (5.1), and we prove a fundamental property of these with respect to the antipod S (5.10).

5.1. **Definition.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; we shall use all the notations of chapter 3; let \mathfrak{H} be a $N - N$ -bimodule, i.e. an Hilbert space equipped with a normal faithful representation a of N on \mathfrak{H} and a normal faithful anti-representation b on \mathfrak{H} , such that $b(N) \subset a(N)'$. Let now V be a unitary from $\mathfrak{H}_{a \otimes \beta_{N^o}} H_\Phi$ onto $\mathfrak{H}_{b \otimes \alpha_N} H_\Phi$, satisfying all the following properties, for all $x \in N$:

$$V(b(x)_{a \otimes \beta_{N^o}} 1) = (1_{b \otimes \alpha_N} \beta(x))V$$

$$V(1_{a \otimes \beta_{N^o}} \alpha(x)) = (a(x)_{b \otimes \alpha_N} 1)V$$

$$V(1_{a \otimes \beta_{N^o}} \hat{\beta}(x)) = (1_{b \otimes \alpha_N} \hat{\beta}(x))V$$

Thanks to these intertwining properties, the following operator has a meaning :

$$(1_{b \otimes \alpha_N} W^*)(1_{b \otimes \alpha_N} \sigma_\nu)(V_{\hat{\beta} \otimes \alpha_{N^o}} 1)(1_{a \otimes \beta_{N^o}} \sigma_{\nu^o})(1_{a \otimes \beta_{N^o}} W)$$

and is a unitary from $\mathfrak{H}_{a \otimes \beta_{N^o}} (H_\Phi \beta \otimes \alpha_N H_\Phi)$ onto $\mathfrak{H}_{b \otimes \alpha_N} H_\Phi \beta \otimes \alpha_N H_\Phi$. Let us recall that the parenthesis in the first Hilbert space means that the antirepresentation β is here acting on the second leg of $H_\Phi \beta \otimes \alpha_N H_\Phi$. Moreover, here, σ_ν is the flip from $H_\Phi \hat{\beta} \otimes \alpha_N H_\Phi$ onto $H_\Phi \alpha \otimes \hat{\beta} H_\Phi$, and σ_{ν^o} is the flip from $H_\Phi \beta \otimes \alpha_N H_\Phi$ onto $H_\Phi \alpha \otimes \beta_{N^o} H_\Phi$.

We may consider also the following unitary from $\mathfrak{H}_{a \otimes \beta_{N^o}} (H_\Phi \beta \otimes \alpha_N H_\Phi)$ onto $\mathfrak{H}_{b \otimes \alpha_N} H_\Phi \beta \otimes \alpha_N H_\Phi$:

$$(V_{b \otimes \alpha_N} 1) \sigma_{b,a}^{2,3} (V_{\alpha \otimes \beta_{N^o}} 1) (1_{a \otimes \beta_{N^o}} \sigma_\nu)$$

Now, in that formula, σ_ν is the flip from $H_\Phi \beta \otimes \alpha_N H_\Phi$ onto $H_\Phi \alpha \otimes \beta_{N^o} H_\Phi$, and $\sigma_{b,a}^{2,3}$ is the flip from $(\mathfrak{H}_{b \otimes \alpha_N} H_\Phi)_{a \otimes \beta_{N^o}} H_\Phi$ onto $(\mathfrak{H}_{a \otimes \beta_{N^o}} H_\Phi)_{b \otimes \alpha_N} H_\Phi$.

We shall say that V is a corepresentation of \mathfrak{G} on the bimodule ${}_a \mathfrak{H}_b$ if we have :

$$\begin{aligned} (1_{b \otimes \alpha_N} W^*)(1_{b \otimes \alpha_N} \sigma_\nu)(V_{\hat{\beta} \otimes \alpha_{N^o}} 1)(1_{a \otimes \beta_{N^o}} \sigma_{\nu^o})(1_{a \otimes \beta_{N^o}} W) \\ = (V_{b \otimes \alpha_N} 1) \sigma_{b,a}^{2,3} (V_{\alpha \otimes \beta_{N^o}} 1) (1_{a \otimes \beta_{N^o}} \sigma_\nu) \end{aligned}$$

5.2. Theorem. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a unitary from $\mathfrak{H}_{a \otimes_{\nu^o} \beta} H_\Phi$ onto $\mathfrak{H}_{b \otimes_\nu \alpha} H_\Phi$, satisfying all the following properties, for all $x \in N$:

$$V(b(x) {}_{a \otimes_{\nu^o} \beta} 1) = (1 {}_{b \otimes_\nu \alpha} \beta(x))V$$

$$V(1 {}_{a \otimes_{\nu^o} \beta} \alpha(x)) = (a(x) {}_{b \otimes_\nu \alpha} 1)V$$

$$V(1 {}_{a \otimes_{\nu^o} \beta} \hat{\beta}(x)) = (1 {}_{b \otimes_\nu \alpha} \hat{\beta}(x))V$$

Then, are equivalent :

(i) V is a corepresentation of \mathfrak{G} on the bimodule ${}_a\mathfrak{H}_b$;

(ii) for any $\xi \in D({}_a\mathfrak{H}, \nu)$ and $\eta \in D(\mathfrak{H}_b, \nu^o)$, the operator $(\omega_{\xi, \eta} * id)(V)$ belongs to M and is such that :

$$\Gamma((\omega_{\xi, \eta} * id)(V)) = (\omega_{\xi, \eta} * id * id)[(V {}_{b \otimes_\nu \alpha} 1) \sigma_{b, a}^{2, 3}(V {}_{a \otimes_{\nu^o} \beta} 1) (1 {}_{a \otimes_{\nu^o} \beta} \sigma_\nu)]$$

Proof. Let V be a corepresentation on ${}_a\mathfrak{H}_b$; let us take now a unitary u in M' ; as we have then :

$$(u^* {}_{\alpha \otimes_{\nu^o} \beta} 1)W(u {}_{\beta \otimes_\nu \alpha} 1) = W$$

it is easy to get, because V is a unitary, that :

$$(1 {}_{b \otimes_\nu \alpha} u^*)V(1 {}_{a \otimes_{\nu^o} \beta} u) = V$$

from which we get, for any $\xi \in D({}_a\mathfrak{H}, \nu)$ and $\eta \in D(\mathfrak{H}_b, \nu^o)$, that the operator $(\omega_{\xi, \eta} * id)(V)$ belongs to M ; from the definition of a corepresentation, we then get :

$$\Gamma((\omega_{\xi, \eta} * id)(V)) = (\omega_{\xi, \eta} * id * id)[(V {}_{b \otimes_\nu \alpha} 1) \sigma_{b, a}^{2, 3}(V {}_{a \otimes_{\nu^o} \beta} 1) (1 {}_{a \otimes_{\nu^o} \beta} \sigma_\nu)]$$

The converse is trivial. □

5.3. Corollary. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ be a $N - N$ -bimodule; let V be unitary from $\mathfrak{H}_{a \otimes_{\nu^o} \beta} H_\Phi$ onto $\mathfrak{H}_{b \otimes_\nu \alpha} H_\Phi$; then, are equivalent :

(i) V is a corepresentation of \mathfrak{G} on the $N - N$ -bimodule ${}_a\mathfrak{H}_b$;

(ii) V^* is a corepresentation of \mathfrak{G}^o on the $N^o - N^o$ -bimodule ${}_b\mathfrak{H}_a$.

Proof. Let us suppose that V is a corepresentation of \mathfrak{G} on the $N - N$ -bimodule ${}_a\mathfrak{H}_b$; then, for any $\xi \in D({}_a\mathfrak{H}, \nu)$ and $\eta \in D(\mathfrak{H}_b, \nu^o)$, the operator $(\omega_{\xi, \eta} * id)(V)$ belongs to M and we have :

$$\Gamma((\omega_{\xi, \eta} * id)(V)) = (\omega_{\xi, \eta} * id * id)[(V {}_{b \otimes_\nu \alpha} 1) \sigma_{b, a}^{2, 3}(V {}_{a \otimes_{\nu^o} \beta} 1) (1 {}_{a \otimes_{\nu^o} \beta} \sigma_\nu)]$$

Therefore, taking the adjoint, we get :

$$\varsigma_\nu \Gamma((\omega_{\eta, \xi} * id)(V^*)) = (\omega_{\eta, \xi} * id * id)[(V^* {}_{a \otimes_{\nu^o} \beta} 1) \sigma_{a, b}^{2, 3}(V^* {}_{b \otimes_\nu \alpha} 1) (1 {}_{a \otimes_{\nu^o} \beta} \sigma_\nu)]$$

which gives, thanks to 5.2, that V^* is a corepresentation of \mathfrak{G}^o on the $N^o - N^o$ -bimodule ${}_b\mathfrak{H}_a$. The proof of the converse is the same, applied to \mathfrak{G}^o . □

5.4. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be an isometry from ${}_a\mathfrak{H}_\beta H_\Phi$ to ${}_b\mathfrak{H}_\alpha H_\Phi$, satisfying all the following properties, for all $x \in N$:

$$V(b(x) {}_a\mathfrak{H}_\beta 1) = (1 {}_b\mathfrak{H}_\alpha \beta(x))V$$

$$V(1 {}_a\mathfrak{H}_\beta \alpha(x)) = (a(x) {}_b\mathfrak{H}_\alpha 1)V$$

$$V(1 {}_a\mathfrak{H}_\beta \hat{\beta}(x)) = (1 {}_b\mathfrak{H}_\alpha \hat{\beta}(x))V$$

and let us suppose that for any $\xi \in D({}_a\mathfrak{H}, \nu)$ and $\eta \in D({}_b\mathfrak{H}, \nu^o)$, that the operator $(\omega_{\xi, \eta} * id)(V)$ belongs to M and is such that :

$$\Gamma((\omega_{\xi, \eta} * id)(V)) = (\omega_{\xi, \eta} * id * id)[(V {}_b\mathfrak{H}_\alpha 1)\sigma_{b,a}^{2,3}(V {}_a\mathfrak{H}_\beta 1)(1 {}_a\mathfrak{H}_\beta \sigma_\nu)]$$

then V is a unitary and a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$.

Proof. We easily have :

$$(1 {}_b\mathfrak{H}_\alpha \sigma_{\nu^o} W^* \sigma_\nu)(V {}_b\mathfrak{H}_\alpha 1)(1 {}_a\mathfrak{H}_\beta \sigma_{\nu^o} W \sigma_\nu) = (1 {}_b\mathfrak{H}_\alpha \sigma_{\nu^o})(V {}_b\mathfrak{H}_\alpha 1)\sigma_{b,a}^{2,3}(V {}_a\mathfrak{H}_\beta 1)$$

Then, taking the adjoints, we get :

$$(1 {}_a\mathfrak{H}_\beta \sigma_{\nu^o} W^* \sigma_\nu)(V^* {}_b\mathfrak{H}_\alpha 1)(1 {}_b\mathfrak{H}_\alpha \sigma_{\nu^o} W \sigma_\nu) = (V^* {}_a\mathfrak{H}_\beta 1)\sigma_{a,b}^{2,3}(V^* {}_b\mathfrak{H}_\alpha 1)(1 {}_b\mathfrak{H}_\alpha \sigma_\nu)$$

and, by multiplication, we get, because $Q = VV^*$ is a projection in $\mathcal{L}(\mathfrak{H}) {}_b^* {}_a M$:

$$\begin{aligned} (1 {}_b\mathfrak{H}_\alpha \sigma_{\nu^o} W^* \sigma_\nu)(Q {}_b\mathfrak{H}_\alpha 1)(1 {}_a\mathfrak{H}_\beta \sigma_{\nu^o} W \sigma_\nu) \\ = (1 {}_b\mathfrak{H}_\alpha \sigma_{\nu^o})(V {}_b\mathfrak{H}_\alpha 1)\sigma_{b,a}^{2,3}(Q {}_b\mathfrak{H}_\alpha 1)\sigma_{a,b}^{2,3}(V^* {}_a\mathfrak{H}_\beta 1)(1 {}_a\mathfrak{H}_\beta \sigma_\nu) \end{aligned}$$

which can be written :

$$(id {}_b^* {}_a \Gamma)(Q) = (V {}_b\mathfrak{H}_\alpha 1)\sigma_{b,a}^{2,3}(Q {}_b\mathfrak{H}_\alpha 1)\sigma_{a,b}^{2,3}(V^* {}_a\mathfrak{H}_\beta 1)$$

from which we get that $(id {}_b^* {}_a \Gamma)(Q) = (Q {}_b\mathfrak{H}_\alpha 1)(id {}_b^* {}_a \Gamma)(Q)$.

But, using 3.12(v), we know that $(id {}_b^* {}_a \Gamma)(Q)$ is equal to

$$(1 {}_b\mathfrak{H}_\alpha \sigma_\nu W^o \sigma_\nu)^*(Q {}_a\mathfrak{H}_\beta 1)(1 {}_b\mathfrak{H}_\alpha \sigma_\nu W^o \sigma_\nu)$$

from which we infer that :

$$(1 {}_b\mathfrak{H}_\alpha \sigma_\nu W^o \sigma_\nu)^*(Q {}_a\mathfrak{H}_\beta 1) = (Q {}_b\mathfrak{H}_\alpha 1)(1 {}_b\mathfrak{H}_\alpha \sigma_\nu W^o \sigma_\nu)^*(Q {}_a\mathfrak{H}_\beta 1)$$

Let's take now $\xi \in D((H_\Phi)_\beta, \nu^o)$ and $\eta \in D({}_a H_\Phi, \nu)$, and let us apply $(id * id * \omega_{\xi, \eta})$ to this last equality. We find :

$$(1 {}_b\mathfrak{H}_\alpha (\omega_{\eta, \xi} * id)(W^o)^*)Q = Q(1 {}_b\mathfrak{H}_\alpha (\omega_{\eta, \xi} * id)(W^o)^*)Q$$

and, by density, we get $(1 {}_b\mathfrak{H}_\alpha y)Q = Q(1 {}_b\mathfrak{H}_\alpha y)Q$ for all $y \in \widehat{M'}$, from which we get that Q belongs also to $\mathcal{L}(\mathfrak{H}) {}_b^* {}_a \widehat{M}$.

Therefore Q belongs to $\mathcal{L}(\mathfrak{H}) {}_b^* {}_a \alpha(N) = b(N)' {}_b\mathfrak{H}_\alpha 1$. So, let $q \in b(N)'$ such that

$Q = q \underset{N}{b} \otimes_{\alpha} 1$, and $(id \underset{N}{b} \ast_{\alpha} \Gamma)(Q) = q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1$

So, we get :

$$(q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1) = (V \underset{N}{b} \otimes_{\alpha} 1)(q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1)(V^* \underset{N^o}{a} \otimes_{\beta} 1)$$

and, taking the adjoints, we get :

$$(V^* \underset{N}{\beta} \otimes_{\alpha} 1)(q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1)(V \underset{N^o}{\alpha} \otimes_{\beta} 1) = (q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1)$$

but, as, by definition of q , we have $(q \underset{N}{b} \otimes_{\alpha} 1 \underset{N}{\beta} \otimes_{\alpha} 1)(V \underset{N^o}{\alpha} \otimes_{\beta} 1) = V \underset{N^o}{\alpha} \otimes_{\beta} 1$, and, as V is an isometry, we finally get that $q = 1$; so, V is a unitary, which finishes the proof. \square

5.5. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a unitary from $\mathfrak{H}_{\underset{N^o}{a} \otimes_{\beta} H_{\Phi}}$ onto $\mathfrak{H}_{\underset{\nu}{b} \otimes_{\alpha} H_{\Phi}}$, satisfying all the following properties, for all $x \in N$:*

$$V(b(x) \underset{N^o}{a} \otimes_{\beta} 1) = (1 \underset{N}{b} \otimes_{\alpha} \beta(x))V$$

$$V(1 \underset{N^o}{a} \otimes_{\beta} \alpha(x)) = (a(x) \underset{N}{b} \otimes_{\alpha} 1)V$$

$$V(1 \underset{N^o}{a} \otimes_{\beta} \hat{\beta}(x)) = (1 \underset{N}{b} \otimes_{\alpha} \hat{\beta}(x))V$$

and let us suppose that, for any $\xi \in D({}_a\mathfrak{H}, \nu)$ and $\eta \in D(\mathfrak{H}_b, \nu^o)$, the operator $(\omega_{\xi, \eta} * id)(V)$ belongs to M ; therefore, it is possible to define $(id * \theta)(V)$, for any θ in $M_*^{\alpha, \beta}$. Moreover, are equivalent :

- (i) V is a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$;
- (ii) the application $\theta \rightarrow (id * \theta)(V)$ from $M_*^{\alpha, \beta}$ into $\mathcal{L}(\mathfrak{H})$ is multiplicative.

Proof. Clear. \square

5.6. Example. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; then the pseudo-multiplicative unitary W verifies :

$$(1_{\mathfrak{H}} \underset{N^o}{\alpha} \otimes_{\hat{\beta}} W)(W \underset{N}{\beta} \otimes_{\alpha} 1_{\mathfrak{H}})(1_{\mathfrak{H}} \underset{N}{\beta} \otimes_{\alpha} W^*) = (W \underset{N^o}{\alpha} \otimes_{\hat{\beta}} 1_{\mathfrak{H}})\sigma_{\alpha, \beta}^{2,3}(W \underset{N}{\beta} \otimes_{\alpha} 1)(1_{\mathfrak{H}} \underset{N}{\beta} \otimes_{\alpha} \sigma_{\nu^o})$$

from which one gets, using the definition of $\hat{\Gamma}$, that we have, for any $\xi \in D((H_{\Phi})_{\beta}, \nu^o)$ and $\eta \in D({}_{\alpha}H_{\Phi}, \nu)$:

$$\varsigma_N \hat{\Gamma}((\omega_{\xi, \eta} * id)(W)) = (\omega_{\xi, \eta} * id * id)[(W \underset{N^o}{\alpha} \otimes_{\hat{\beta}} 1_{\mathfrak{H}})\sigma_{\alpha, \beta}^{2,3}(W \underset{N}{\beta} \otimes_{\alpha} 1)(1_{\mathfrak{H}} \underset{N}{\beta} \otimes_{\alpha} \sigma_{\nu^o})]$$

which means, thanks to 5.4, that W is a corepresentation of $(\hat{\mathfrak{G}})^o$ on the $N^o - N^o$ bimodule ${}_{\beta}(H_{\Phi})_{\alpha}$. So, $(\sigma_{\nu^o} W \sigma_{\nu^o})^*$ is a corepresentation of \mathfrak{G}^o on the $N^o - N^o$ bimodule ${}_{\hat{\beta}}(H_{\Phi})_{\alpha}$, and, using 5.3, $\sigma_{\nu^o} W \sigma_{\nu^o}$ is a corepresentation of \mathfrak{G} on the $N - N$ bimodule ${}_{\alpha}(H_{\Phi})_{\hat{\beta}}$. We get also that $(\sigma_{\nu} W^o \sigma_{\nu})^*$ is a corepresentation of \mathfrak{G} on the $N - N$ bimodule ${}_{\hat{\alpha}}(H_{\Phi})_{\beta}$ and that W^{oc} is a corepresentation of $\hat{\mathfrak{G}}^c$ on the $N^o - N^o$ bimodule ${}_{\hat{\beta}}(H_{\Phi})_{\hat{\alpha}}$.

5.7. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$. Then, we have :

$$(V_{\alpha \otimes_{N^o} \beta} 1)(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W^* \sigma_{\nu})(V^*_{\beta \otimes_{N^o} \alpha} 1) = \sigma_{a,b}^{2,3}(V^*_{\beta \otimes_{N^o} \alpha} 1)(1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu})(1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu^o} W^* \sigma_{\nu})$$

Proof. From 5.1, we easily have :

$$(1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu^o} W^* \sigma_{\nu})(V_{\beta \otimes_{N^o} \alpha} 1)(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W \sigma_{\nu}) = (1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu^o})(V_{\beta \otimes_{N^o} \alpha} 1) \sigma_{b,a}^{2,3}(V_{\alpha \otimes_{N^o} \beta} 1)$$

Then, taking the adjoints, we get :

$$(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W^* \sigma_{\nu})(V^*_{\beta \otimes_{N^o} \alpha} 1)(1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu^o} W \sigma_{\nu}) = (V^*_{\alpha \otimes_{N^o} \beta} 1) \sigma_{a,b}^{2,3}(V^*_{\beta \otimes_{N^o} \alpha} 1)(1_{\beta \otimes_{N^o} \alpha} \sigma_{\nu})$$

from which we get the result, because V is an unitary. \square

5.8. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$. Let ξ_1, ξ_2 be in $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$, η_1, η_2 be in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$. Then, the element :

$$(\omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1), V^*(\xi_{2b} \otimes_{\nu} \eta_2)} * id)(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W \sigma_{\nu})$$

belongs to $D(S)$, and :

$$S((\omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1), V^*(\xi_{2b} \otimes_{\nu} \eta_2)} * id)(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W \sigma_{\nu})) = (\omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1), V^*(\xi_{2b} \otimes_{\nu} \eta_2)} * id)(1_{\alpha \otimes_{N^o} \beta} \sigma_{\nu^o} W^* \sigma_{\nu})$$

Proof. By linearity, it suffices to prove that the element of \widehat{M}_* defined, for y in \widehat{M} by :

$$y \mapsto \omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1)}(1_{\alpha \otimes_{N^o} \beta} y)$$

belongs to $\widehat{M}_*^{\alpha, \hat{\beta}}$. For $n \in N$, we have :

$$(1_{\alpha \otimes_{N^o} \beta} \alpha(n)) V^*(\xi_{1b} \otimes_{\nu} \eta_1) = V^*(a(n) \xi_{1b} \otimes_{\nu} \eta_1)$$

$$(1_{\alpha \otimes_{N^o} \beta} \hat{\beta}(n)) V^*(\xi_{1b} \otimes_{\nu} \eta_1) = V^*(\xi_{1b} \otimes_{\nu} \hat{\beta}(n) \eta_1)$$

and, therefore, if $n \in \mathfrak{N}_\nu$,

$$\|(1_{\alpha \otimes_{N^o} \beta} \alpha(n)) V^*(\xi_{1b} \otimes_{\nu} \eta_1)\| \leq \|R^\alpha(\eta_1)\| \|R^a(\xi_1)\| \|\Lambda_\nu(n)\|$$

$$\|(1_{\alpha \otimes_{N^o} \beta} \hat{\beta}(n)) V^*(\xi_{1b} \otimes_{\nu} \eta_1)\| \leq \|R^b(\xi_1)\| \|R^\alpha(\eta_1)\| \|\Lambda_\nu(n)\|$$

which, therefore, gives the result. \square

5.9. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$. Let ξ_1, ξ_2 be in $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$, η_1, η_2 be in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^o)$. Then, the element :

$$(id * \omega_{\eta_1, \eta_2})(W)(\omega_{\xi_1, \xi_2} * id)(V)$$

belongs to $D(S)$, and :

$$S[(id * \omega_{\eta_1, \eta_2})(W)(\omega_{\xi_1, \xi_2} * id)(V)] = (\omega_{\xi_1, \xi_2} * id)(V^*)(id * \omega_{\eta_1, \eta_2})(W^*)$$

Proof. Using 5.7, we get that :

$$(id * \omega_{\eta_1, \eta_2})(W)(\omega_{\xi_1, \xi_2} * id)(V) = (\omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1), V^*(\xi_{2b} \otimes_{\nu} \eta_2)} * id)(1_{\underset{N^o}{a} \otimes_{\beta} \sigma_{\nu^o}} W \sigma_{\nu})$$

$$(\omega_{\xi_1, \xi_2} * id)(V^*)(id * \omega_{\eta_1, \eta_2})(W^*) = (\omega_{V^*(\xi_{1b} \otimes_{\nu} \eta_1), V^*(\xi_{2b} \otimes_{\nu} \eta_2)} * id)(1_{\underset{N^o}{a} \otimes_{\beta} \sigma_{\nu^o}} W^* \sigma_{\nu})$$

We then get the result by 5.8. \square

5.10. Theorem. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule; let V be a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$. Let ξ_1, ξ_2 in $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$; then the element $(\omega_{\xi_1, \xi_2} * id)(V^*)$ belongs to $D(S)$, and :

$$S((\omega_{\xi_1, \xi_2} * id)(V)) = (\omega_{\xi_1, \xi_2} * id)(V^*)$$

Proof. Using 5.9, we get that, for any x of the form $(id * \omega_{\eta_1, \eta_2})(W^*)$, where η_1, η_2 are in $D({}_\alpha H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^o)$, the operator $x(\omega_{\xi_1, \xi_2} * id)(V)$ belongs to $D(S)$, and $S(x(\omega_{\xi_1, \xi_2} * id)(V^*)) = (\omega_{\xi_1, \xi_2} * id)(V^*)S(x)$,

If, moreover, η_1 belongs to $D(P^{-1/2})$ and is such that $P^{-1/2}\eta_1$ belongs to $D((H_\Phi)_{\hat{\beta}}, \nu^o)$, and if η_2 belongs to $D(P^{1/2})$ and is such that $P^{1/2}\eta_2$ belongs to $D({}_\alpha(H_\Phi, \nu))$, then using ([E5], 3.8), we know that :

$$(id * \omega_{J_\Phi \eta_1, J_\Phi \eta_2})(W^*) = (id * \omega_{P^{-1/2} \eta_1, P^{1/2} \eta_2})(W^*)^*$$

and we know ([E5], 3.9) that the linear space generated by such elements is weakly dense in M . Let \mathcal{A} be the algebra generated by these elements, which is therefore involutive, and dense in M .

Therefore, we obtain that $S(x(\omega_{\xi_1, \xi_2} * id)(V^*)) = (\omega_{\xi_1, \xi_2} * id)(V^*)S(x)$ for any $x \in \mathcal{A}$. Then, by Kaplanski's theorem, it is therefore possible to find a sequence a_n in $D(S)$ increasing to 1 such that :

$$S(a_n(\omega_{\xi_1, \xi_2} * id)(V^*)) = (\omega_{\xi_1, \xi_2} * id)(V^*)S(a_n)$$

Let's take now

$$e_n = 1/\sqrt{\pi} \int_{-\infty}^{\infty} e^{-t^2} \tau_t(a_n) dt$$

As S is closed, we have again

$$S(e_n(\omega_{\xi_1, \xi_2} * id)(V^*)) = (\omega_{\xi_1, \xi_2} * id)(V^*)S(e_n)$$

Moreover, e_n is increasing to 1, and, as $\tau_{-i/2}(e_n)$ is bounded, $\tau_{-i/2}(e_n)$ converges also to 1, and so does $S(e_n)$. So, we get the result, using again the closedness of S . \square

5.11. **Theorem.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and ${}_a\mathfrak{H}_b$ a $N - N$ -bimodule, and let us suppose that $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$ is dense in \mathfrak{H} . Let V be a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$. Then, for any ζ_1, ζ_2 in \mathfrak{H} , for any $\xi \in D((H_\Phi)_\beta, \nu^o) \cap \mathcal{D}(P^{1/2})$ such that $P^{1/2}\xi$ belongs to $D({}_\alpha H_\Phi, \nu)$, and for any $\eta \in D({}_\alpha H_\Phi, \nu) \cap D(P^{-1/2})$ such that $P^{-1/2}\eta$ belongs to $D((H_\Phi)_\beta, \nu^o)$, we have :

$$(V(\zeta_1 {}_a\otimes_\beta \xi)|\zeta_2 {}_b\otimes_\alpha \eta) = (V^*(\zeta_1 {}_b\otimes_\alpha J_{\widehat{\Phi}}P^{-1/2}\eta)|\zeta_2 {}_a\otimes_\beta J_{\widehat{\Phi}}P^{1/2}\xi)$$

Proof. It is clear that $J_{\widehat{\Phi}}P^{-1/2}\eta$ belongs to $D({}_\alpha H_\Phi, \nu)$ and that $J_{\widehat{\Phi}}P^{1/2}\xi$ belongs to $D((H_\Phi)_\beta, \nu^o)$. Moreover, we have, if we take ζ_1, ζ_2 in $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$:

$$\begin{aligned} (V(\zeta_1 {}_a\otimes_\beta \xi)|\zeta_2 {}_b\otimes_\alpha \eta) &= (\zeta_1 {}_a\otimes_\beta \xi|V^*(\zeta_2 {}_b\otimes_\alpha \eta)) \\ &= (\xi|(\omega_{\zeta_2, \zeta_1} * id)(V^*)\eta) \end{aligned}$$

which, using 5.10, and the fact that $J_{\widehat{\Phi}}P^{1/2} = P^{-1/2}J_{\widehat{\Phi}}$, is equal to :

$$\begin{aligned} (\xi|S[(\omega_{\zeta_2, \zeta_1} * id)(V)]\eta) &= (\xi|J_{\widehat{\Phi}}P^{-1/2}(\omega_{\zeta_2, \zeta_1} * id)(V)^*P^{1/2}J_{\widehat{\Phi}}\eta) \\ &= ((\omega_{\zeta_2, \zeta_1} * id)(V)^*J_{\widehat{\Phi}}P^{-1/2}\eta|J_{\widehat{\Phi}}P^{1/2}\xi) \\ &= (J_{\widehat{\Phi}}P^{-1/2}\eta|(\omega_{\zeta_2, \zeta_1} * id)(V)J_{\widehat{\Phi}}P^{1/2}\xi) \\ &= (\zeta_1 {}_b\otimes_\alpha J_{\widehat{\Phi}}P^{-1/2}\eta|V(\zeta_2 {}_a\otimes_\beta J_{\widehat{\Phi}}P^{1/2}\xi)) \\ &= (V^*(\zeta_1 {}_b\otimes_\alpha J_{\widehat{\Phi}}P^{-1/2}\eta)|\zeta_2 {}_a\otimes_\beta J_{\widehat{\Phi}}P^{1/2}\xi) \end{aligned}$$

Therefore, the result is proved for ζ_1, ζ_2 in $D({}_a\mathfrak{H}, \nu) \cap D(\mathfrak{H}_b, \nu^o)$, and it remains true, by continuity, for any ζ_1, ζ_2 in \mathfrak{H} . \square

6. ACTIONS OF MEASURED QUANTUM GROUPOIDS

In this chapter, we define actions of measured quantum groupoids on von Neumann algebras (6.1) and actions implemented by a corepresentation (6.6). We define also invariant elements by an action (6.11), the canonical operator-valued weight on the invariants (6.13), and integrable actions (6.14)

6.1. **Definition.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and let A be a von Neumann algebra.

An action of \mathfrak{G} on A is a couple (b, \mathfrak{a}) , where :

- (i) b is an injective $*$ -antihomomorphism from N into A ;
- (ii) \mathfrak{a} is an injective $*$ -homomorphism from A into $A {}_b\alpha_N M$;
- (iii) b and \mathfrak{a} are such that, for all n in N :

$$\mathfrak{a}(b(n)) = 1 {}_b\otimes_\alpha \beta(n)$$

(which allow us to define $\mathfrak{a} {}_b\alpha_N id$ from $A {}_b\alpha_N M$ into $A {}_b\alpha_N M {}_\beta\alpha_N M$) and such that :

$$(\mathfrak{a} {}_b\alpha_N id)\mathfrak{a} = (id {}_b\alpha_N \Gamma)\mathfrak{a}$$

If there is no ambiguity, we shall say that \mathfrak{a} is the action.

So, a measured quantum groupoid \mathfrak{G} can act only on a von Neumann algebra A which is a right module over the basis N .

Moreover, if M is abelian, then, as, for all $n \in N$, $\mathfrak{a}(b(n)) = 1 {}_b\otimes_\alpha \beta(n)$ commutes with $\mathfrak{a}(x)$, for all $x \in A$, we see that $b(N)$ is in the center of A . As in that case

(3.13) the measured quantum groupoid comes then from a measured groupoid \mathcal{G} , we have $N = L^\infty(\mathcal{G}^{(0)}, \nu)$, and A can be decomposed as $A = \int_{\mathcal{G}^{(0)}}^\oplus A^x d\nu(x)$.

6.2. **Example.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; then (id, β) is an action of \mathfrak{G} on N^o , we shall call the trivial action of \mathfrak{G} on N^o .

6.3. **Example.** Let \mathcal{G} be a measured groupoid, and let \mathfrak{a} be an action of \mathcal{G} on a von Neumann algebra $A = \int_{\mathcal{G}^{(0)}}^\oplus A^x d\nu(x)$, as defined in ([Y3], def. 3.1(i)); let us define then b as the homomorphism which sends $L^\infty(\mathcal{G}^{(0)})$ into the diagonalizable operators in A . Then, (b, \mathfrak{a}) is an action (in the sense of 6.1) of the measured quantum groupoid \mathcal{G} defined in 3.13 on A ([OR], Th. 3.2), i.e. for all $g \in \mathcal{G}$, there exists a family of $*$ -isomorphisms \mathfrak{a}_g from $A^{s(g)}$ onto $A^{r(g)}$, such that, if $(g_1, g_2) \in \mathcal{G}^{(2)}$, we have $\mathfrak{a}_{g_1 g_2} = \mathfrak{a}_{g_1} \mathfrak{a}_{g_2}$, and such that, for any normal positive functional $\omega = \int_{\mathcal{G}^{(0)}}^\oplus \omega^x d\nu(x)$, and any $y = \int_{\mathcal{G}^{(0)}}^\oplus y^x d\nu(x)$, the function $g \mapsto \omega^{r(g)}(\mathfrak{a}_g(y^{s(g)}))$ is μ -measurable. We then get :

$$\mathfrak{a}\left(\int_{\mathcal{G}^{(0)}}^\oplus y^x d\nu(x)\right) = \int_{\mathcal{G}}^\oplus \mathfrak{a}_g(y^{s(g)}) d\mu(g)$$

Let now \mathfrak{b} be a coaction of \mathcal{G} on a von Neumann algebra B which is a $L^\infty(\mathcal{G}^{(0)}, \nu)$ -module, as defined in ([Y3], def. 3.1(ii)); let us define then b as the homomorphism which sends $L^\infty(\mathcal{G}^{(0)}, \nu)$ into B ; then, (b, \mathfrak{b}) is an action of $\widehat{\mathcal{G}}$ (as defined in 3.13) on B . ([Y3], 3.1 (ii) and [OR], 2.3).

6.4. **Example.** Let $\mathbf{G} = (M, \Gamma, \Phi, \Psi)$ be a locally compact quantum group; an action of \mathbf{G} on a von Neumann algebra A is an injective $*$ -homomorphism \mathfrak{a} from M into $A \otimes M$, such that

$$(\mathfrak{a} \otimes id)\mathfrak{a} = (id \otimes \Gamma)\mathfrak{a}$$

(where id means the identity of A or of M). ([V2], 1.1)

Writing now id for the canonical $*$ -homomorphism from \mathbb{C} into A , we get that (id, \mathfrak{a}) is an action of the measured quantum groupoid \mathbf{G} (3.14) on A .

6.5. **Example.** Let $M_0 \subset M_1$ a depth 2 inclusion of von Neumann algebras, with a regular operator-valued weight T_1 from M_1 to M_0 , as defined in 2.2. Let \mathfrak{G}_1 be the measured quantum groupoid constructed (3.15) from this inclusion, and \mathfrak{G}_2 the measured quantum groupoid constructed from the inclusion $M_1 \subset M_2$, which is isomorphic to $\widehat{\mathfrak{G}_1}^o$; we shall use the notations introduced in 2.2 and 3.15. Then, there exists ([EV], 7.3) a canonical action of \mathfrak{G}_2 on M_1 , which can be described as follows : the anti-representation of the basis $M'_1 \cap M_2$ (which, using j_1 , is anti-isomorphic to $M'_0 \cap M_1$), is given by the natural inclusion of $M'_0 \cap M_1$ into M_1 , and the homomorphism from M_1 is given by the natural inclusion of M_1 into M_3 (which is, thanks to ([EV], 4.6), isomorphic to $M_1 * \mathcal{L}(H_{\chi_2})$).

6.6. **Proposition.** Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a\mathfrak{H}_b$ be a $N - N$ bimodule, and V be a corepresentation of \mathfrak{G} on \mathfrak{H} ; for any $x \in a(N)'$, let us write $\mathfrak{a}(x) = V(x {}_a\otimes_\beta 1)V^*$ and let A be a von Neumann algebra on \mathfrak{H} , such that

$b(N) \subset A \subset a(N)'$, and $\mathfrak{a}(A) \subset A {}_b\otimes_\alpha M$; then :

(i) (b, \mathfrak{a}) is an action of \mathfrak{G} on $a(N)'$;

(ii) $(b, \mathfrak{a}|_A)$ is an action of \mathfrak{G} on A .

In such a situation, we shall say that \mathfrak{a} and $\mathfrak{a}|_A$ are implemented by V .

Proof. The intertwining properties of V imply that, for all $n \in N$:

$$\mathbf{a}(b(n)) = 1_{b \otimes_{\alpha} N} \beta(n)$$

$$(a(n)_{b \otimes_{\alpha} 1}) \mathbf{a}(x) = \mathbf{a}(x)(a(n)_{b \otimes_{\alpha} 1})$$

and this last property gives that $\mathbf{a}(a(N)') \subset a(N)'_{b^* \otimes_{\alpha} N} M$. Therefore, we can consider $(\mathbf{a}_{b^* \otimes_{\alpha} N} id) \mathbf{a}(x)$, which will be equal to :

$$(V_{b \otimes_{\alpha} N} 1) \sigma_{b,a}^{2,3} (V_{\alpha \otimes_{\beta} N^o} 1) (x_{a \otimes_{\beta} N^o} 1_{\alpha \otimes_{\beta} N^o}) (V^*_{\alpha \otimes_{\beta} N^o} 1) \sigma_{a,b}^{2,3} (V^*_{b \otimes_{\alpha} N} 1)$$

which, thanks to 5.1, is equal to :

$$(1_{b \otimes_{\alpha} N} W^*) (1_{b \otimes_{\alpha} N} \sigma_{\nu}) (\mathbf{a}(x)_{\beta \otimes_{\alpha} N} 1) (1_{b \otimes_{\alpha} N} \sigma_{\nu^o}) (1_{b \otimes_{\alpha} N} W)$$

which is equal to $(id_{b^* \otimes_{\alpha} N} \Gamma) \mathbf{a}(x)$. \square

6.7. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, A a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} ; then, there exist an Hilbert space \mathfrak{H} , a faithful normal representation π of A on \mathfrak{H} , a normal faithful representation a of N on \mathfrak{H} , such that $\pi(A) \subset a(N)'$, a corepresentation V of \mathfrak{G} on the $N - N$ bimodule ${}_a \mathfrak{H}_{\pi \circ b}$, such that, for all x in A :*

$$(\pi_{b^* \otimes_{\alpha} N} id) \mathbf{a}(x) = V(\pi(x)_{a \otimes_{\beta} N^o} 1) V^*$$

Proof. Let's take $\mathfrak{H} = L^2(A)_{b \otimes_{\alpha} N} H_{\Phi}$, let us write $a(n) = 1_{L^2(A)}_{b \otimes_{\alpha} N} \hat{a}(n)$, for all $n \in N$, and take $V = 1_{L^2(A)}_{b \otimes_{\alpha} N^o} (\sigma_{\nu} W^o \sigma_{\nu})^*$ which is (5.6) a corepresentation of \mathfrak{G} on ${}_a \mathfrak{H}_{(1_{L^2(A)}_{b \otimes_{\alpha} N^o} \beta)}$ and choose $\pi = \mathbf{a}$ to obtain the result. \square

6.8. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a \mathfrak{H}_b$ be a $N - N$ bimodule, and V be a corepresentation of \mathfrak{G} on \mathfrak{H} ; let A be a von Neumann algebra on \mathfrak{H} , such that $b(N) \subset A \subset a(N)'$; then, are equivalent :*

- (i) *the corepresentation V of \mathfrak{G} implements an action \mathbf{a} of \mathfrak{G} on A ;*
- (ii) *the corepresentation V^* of \mathfrak{G}^o implements an action \mathbf{a}' of \mathfrak{G}^o on A' .*

Proof. Let us suppose (i); then, we have $V(A_{a \otimes_{\beta} N^o} 1) V^* \subset A_{b^* \otimes_{\alpha} N} M$; taking the commutants, we get :

$$A'_{b \otimes_{\alpha} N} M' \subset V(A'_{a^* \otimes_{\beta} N^o} \mathcal{L}(H_{\Phi})) V^*$$

and, therefore, $A'_{b \otimes_{\alpha} N} 1 \subset V(A'_{a^* \otimes_{\beta} N^o} \mathcal{L}(H_{\Phi})) V^*$, from which have :

$$V^*(A'_{b \otimes_{\alpha} N} 1) V \subset A'_{a^* \otimes_{\beta} N^o} \mathcal{L}(H_{\Phi})$$

but, from 5.3 and 6.6 applied to V^* , we get that :

$$V^*(b(N)'_{b \otimes_{\alpha} N} 1) V \subset b(N)'_{a^* \otimes_{\beta} N^o} M$$

and, therefore, we have :

$$V^*(A'_{b \otimes_{\alpha} N} 1) V \subset A'_{a^* \otimes_{\beta} N^o} M$$

from which we get (ii), using 6.6 again. If we apply this proof to \mathfrak{G}^o , we obtain that (ii) implies (i). \square

6.9. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, A a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A ; let ψ a normal semi-finite faithful weight on A , and let us define, for all $n \in N$, $a(n) = J_\psi b(n^*) J_\psi$, which gives to ${}_a(H_\psi)_b$ a canonical structure of $N - N$ bimodule; let V be a corepresentation of \mathfrak{G} on H_ψ ; let us suppose that V implements \mathfrak{a} , i.e. that, for any $x \in A$, we have :

$$\mathfrak{a}(x) = V(x {}_a \otimes_{\beta} 1) V^*$$

We shall say that V is a standard implementation of \mathfrak{a} if, moreover, we have :

$$V^* = (J_\psi {}_\alpha \otimes_{\beta} J_{\widehat{\Phi}}) V (J_\psi {}_b \otimes_{\alpha} J_{\widehat{\Phi}})$$

6.10. Example. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; then (β, Γ) is an action of \mathfrak{G} on M .

Moreover, $(\sigma_\nu W^o \sigma_\nu)^*$ is a corepresentation of \mathfrak{G} on ${}_{\widehat{\alpha}}(H_{\Phi})_{\beta}$ which is a standard implementation of Γ .

6.11. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A ; we shall call $A^{\mathfrak{a}}$ the invariant subalgebra of A defined by :

$$A^{\mathfrak{a}} = \{x \in A \cap b(N)'; \mathfrak{a}(x) = x {}_b \otimes_{\alpha} 1\}$$

In the example 6.2, the invariant subalgebra is $Z(N)$; in the example 6.5, the invariant subalgebra is M_0 ([EV], 7.5); in the example 6.10, the invariant subalgebra is $\alpha(N)$ (3.11(v)).

6.12. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A . Let $\Phi = \nu \circ \alpha^{-1} \circ T$; for any $x \in A^+$, the extended positive element of A

$$T_{\mathfrak{a}}(x) = (id {}_b {}^*_{\alpha} \Phi) \mathfrak{a}(x)$$

is an extended positive element of $A^{\mathfrak{a}}$.

Proof. Let $\omega \in (A {}_b {}^*_{\alpha} M)_*^+$; then, we have :

$$\begin{aligned} \omega \circ \mathfrak{a}(T_{\mathfrak{a}}(x)) &= \omega(id {}_b {}^*_{\alpha} id {}_{\beta} {}^*_{\alpha} \Phi) (\mathfrak{a} {}_b {}^*_{\alpha} id) \mathfrak{a}(x) \\ &= \omega(id {}_b {}^*_{\alpha} id {}_{\beta} {}^*_{\alpha} \Phi) (id {}_b {}^*_{\alpha} \Gamma) \mathfrak{a}(x) \end{aligned}$$

which, thanks to 4.12(ii), is equal to :

$$\omega((id {}_b {}^*_{\alpha} \Phi) \mathfrak{a}(x) {}_{\beta} \otimes_{\alpha} 1)$$

from which we get that $\mathfrak{a}(T_{\mathfrak{a}}(x)) = T_{\mathfrak{a}}(x) {}_b \otimes_{\alpha} 1$, which is the result. \square

6.13. Proposition. Let $(N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and let A be a von Neumann algebra, (b, \mathfrak{a}) an action of $(N, M, \alpha, \beta, \Gamma, T, T', \nu)$ on A . Let $\Phi = \nu \circ \alpha^{-1} \circ T$, and, for any $x \in A^+$, let $T_{\mathfrak{a}}(x) = (id {}_b {}^*_{\alpha} \Phi) \mathfrak{a}(x)$ be the extended positive element of $A^{\mathfrak{a}}$ defined in 6.12. Then $T_{\mathfrak{a}}$ is a normal faithful operator-valued weight from A on $A^{\mathfrak{a}}$.

Proof. Let y in A^a ; let $\omega \in A_*^+$, such that there exists $k > 0$ for which $\omega \circ b \leq k\nu$; as $y \in b(N)'$, the element ω_y of A_*^+ defined, for all $x \in A$ by $\omega_y(x) = \omega(y^*xy)$ satisfies the same property, and we have, for $x \in A^+$:

$$\begin{aligned} \langle T_a(y^*xy), \omega \rangle &= \omega((id_{b^*_\alpha} \Phi) \mathbf{a}(y^*xy)) \\ &= \omega((id_{b^*_\alpha} \Phi)(y^*_{b \otimes_\alpha 1} \mathbf{a}(x)(y_{b \otimes_\alpha 1}))) \\ &= \omega_y(T_a(x)) \\ &= \langle y^*T_a(x)y, \omega \rangle \end{aligned}$$

which proves that T_a is indeed an operator-valued weight. Normality and faithfulness are trivial. \square

6.14. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid-id, and let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A ; we shall say that the action is integrable if the operator-valued weight T_a introduced in 6.12 and 6.13 is semi-finite.

7. SOME TECHNICAL PROPERTIES OF ACTIONS

In this chapter, we define technical properties (property A in 7.1, saturation property in 7.3) and prove technical results (7.6, 7.7) about these properties which will be necessary in the sequel. We shall prove in chapter 11 that these properties are always fulfilled.

7.1. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, (b, \mathbf{a}) an action of \mathfrak{G} on a von Neumann algebra A ; following ([E1], III2), we shall say that the action \mathbf{a} satisfies the property (A) if :

$$(\mathbf{a}(A) \cup 1_{b \otimes_\alpha} \alpha(N)')'' = A_{b^*_\alpha} \mathcal{L}(H_\Phi)$$

7.2. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; then, we have :

$$(\Gamma(M) \cup 1_{\beta \otimes_\alpha} \alpha(N)')'' = M_{\beta^*_\alpha} \mathcal{L}(H_\Phi)$$

In other words, the action (β, Γ) of \mathfrak{G} on M (6.10) satisfies the property (A) (7.1).

Proof. Let y be in $\beta(N)'$ such that $W(y_{\beta \otimes_\alpha} 1)W^*$ belongs to $\mathcal{L}(H_\Phi)_{\alpha^*_\beta} M'$; then, for any u unitary in M , we have :

$$(1_{\alpha \otimes_\beta} u)_{N^\circ} W(y_{\beta \otimes_\alpha} 1)_{N^\circ} W^*(1_{\alpha \otimes_\beta} u^*)_{N^\circ} = W(y_{\beta \otimes_\alpha} 1)_{N^\circ} W^*$$

which can be written :

$$\Gamma(u)(y_{\beta \otimes_\alpha} 1)_{N^\circ} \Gamma(u^*)_{N^\circ} = y_{\beta \otimes_\alpha} 1_{N^\circ}$$

and, therefore, we have $\Gamma(u)(y_{\beta \otimes_\alpha} 1)_{N^\circ} = (y_{\beta \otimes_\alpha} 1)_{N^\circ} \Gamma(u)$; therefore, y commutes with all elements of the form $(id_{\alpha^*_\beta} \omega)_{N^\circ} \Gamma(u)$, for all u unitary in M and all $\omega \in M_*$ for which there exists $k > 0$ such that $\omega \circ \alpha \leq k\nu$, and, using A.7, we get that y belongs to M' . So, we have :

$$(\beta(N)'_{\beta \otimes_\alpha} 1)_{N^\circ} \cap W^*(\mathcal{L}(H_\Phi)_{\alpha^*_\beta} M')_{N^\circ} W = M'_{\beta \otimes_\alpha} 1_{N^\circ}$$

and, as $(\beta(N)'_{\beta \otimes_\alpha} 1)_{N^\circ} = \mathcal{L}(H_\Phi)_{\beta^*_\alpha} \alpha(N)_{N^\circ} = (1_{\beta \otimes_\alpha} \alpha(N)')'_{N^\circ}$, we get the result, taking the commutants. \square

7.3. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A ; following ([ES1], II8), we shall denote:

$$Sata = \{X \in A_{b^* \alpha} M, (\mathfrak{a}_{b^* \alpha} id)(X) = (id_{b^* \alpha} \Gamma)(X)\}$$

By definition, we have $\mathfrak{a}(A) \subset Sata$, and we shall say that \mathfrak{a} is saturated if $\mathfrak{a}(A) = Sata$

7.4. Lemma. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a \mathfrak{H}_b$ a $N - N$ bimodule, V a corepresentation of \mathfrak{G} on ${}_a \mathfrak{H}_b$; let $SatV$ be the set of elements X in $a(N)'_{b^* \alpha} \hat{\beta}(N)'$ such that :

$$\begin{aligned} (V_{b \otimes \alpha} 1) \sigma_{b,a}^{2,3} (X_{a \otimes \beta} 1) \sigma_{a,b}^{2,3} (V_{\beta \otimes \alpha}^* 1) \\ = (1_{b \otimes \alpha} W^*) (1_{b \otimes \alpha} \sigma_\nu) (X_{\beta \otimes \alpha} 1) (1_{b \otimes \alpha} \sigma_\nu) (1_{b \otimes \alpha} W) \end{aligned}$$

Then, we have :

$$\begin{aligned} (i) \quad SatV &= V(\mathcal{L}(\mathfrak{H})_{a^* \beta} \widehat{M}') V^*; \\ (ii) \quad V^* (1_{b \otimes \alpha} \widehat{M}') V &\subset \mathcal{L}(\mathfrak{H})_{a^* \beta} \widehat{M}' \end{aligned}$$

Proof. As V is a corepresentation, we have, by definition :

$$\begin{aligned} (1_{b \otimes \alpha} W^*) (1_{b \otimes \alpha} \sigma_\nu) (V_{\beta \otimes \alpha} 1) (1_{a \otimes \beta} \sigma_{\nu^o}) (1_{a \otimes \beta} W) \\ = (V_{b \otimes \alpha} 1) \sigma_{b,a}^{2,3} (V_{\alpha \otimes \beta} 1) (1_{a \otimes \beta} \sigma_{\nu^o}) \end{aligned}$$

and, therefore :

$$\begin{aligned} (1_{b \otimes \alpha} \sigma_{\nu^o}) (1_{b \otimes \alpha} W) (V_{b \otimes \alpha} 1) \sigma_{b,a}^{2,3} \\ = (V_{\beta \otimes \alpha} 1) (1_{a \otimes \beta} \sigma_{\nu^o}) (1_{a \otimes \beta} W) (1_{\alpha \otimes \beta} \sigma_{\nu^o}) (V_{\alpha \otimes \beta}^* 1) \end{aligned}$$

So, X belongs to $SatV$ if and only if we have :

$$(1_{\alpha \otimes \beta} \sigma_{\nu^o} W \sigma_{\nu^o}) (V^* X V_{\alpha \otimes \beta} 1) = (V^* X V_{\beta \otimes \alpha} 1) (1_{\alpha \otimes \beta} \sigma_{\nu^o} W \sigma_{\nu^o})$$

from which we get that X belongs to $SatV$ if and only if $V^* X V$ belongs to $\mathcal{L}(\mathfrak{H})_{a^* \beta} \widehat{M}'$,

which finishes the proof of (i).

As it is clear that $1_{b \otimes \alpha} \widehat{M}' \subset SatV$, we get (ii) from (i). \square

7.5. Lemma. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a \mathfrak{H}_b$ a $N - N$ bimodule, A a von Neumann algebra such that $b(N) \subset A \subset a(N)'$, V a corepresentation of \mathfrak{G} on ${}_a \mathfrak{H}_b$ which implements an action (b, \mathfrak{a}) on A ; then we have :

$$Sata = A_{b^* \alpha} M \cap V(\mathcal{L}(\mathfrak{H})_{a^* \beta} \widehat{M}') V^*$$

and also :

$$Sata = A_{b^* \alpha} \mathcal{L}(H_\Phi) \cap V(a(N)'_{a \otimes \beta} 1) V^*$$

Proof. Let $X \in A_{b^*_{\alpha} M}$; as we have :

$$(\mathfrak{a}_{b^*_{\alpha} N} id)(X) = (V_{b \otimes_{\alpha} N} 1) \sigma_{b,a}^{2,3}(X_{a \otimes_{\beta} N^o} 1) \sigma_{a,b}^{2,3}(V_{\beta \otimes_{\alpha} N}^* 1)$$

and :

$$(id_{b^*_{\alpha} N} \Gamma)(X) = (1_{b \otimes_{\alpha} N} W^*)(1_{b \otimes_{\alpha} N} \sigma_{\nu})(X_{\beta \otimes_{\alpha} N} 1)(1_{b \otimes_{\alpha} N} \sigma_{\nu})(1_{b \otimes_{\alpha} N} W)$$

we get that $Sat \mathfrak{a} = A_{b^*_{\alpha} N} M \cap Sat V$, which, thanks to 7.4(i), gives the first formula.

Starting from this formula, we get, because V is a corepresentation of \mathfrak{G} :

$$\begin{aligned} Sat \mathfrak{a} &= V(V^*(A_{b^*_{\alpha} N} M) V \cap \mathcal{L}(\mathfrak{H})_{a^*_{\beta} N^o} \widehat{M}') V^* \\ &\subset V(\mathcal{L}(\mathfrak{H})_{a^*_{\beta} N} M \cap \mathcal{L}(\mathfrak{H})_{a^*_{\beta} N^o} \widehat{M}') V^* \\ &= V(\mathcal{L}(\mathfrak{H})_{a^*_{\beta} N} \beta(N)) V^* \\ &= V(a(N)'_{a \otimes_{\beta} N^o} 1) V^* \end{aligned}$$

and, therefore, $Sat \mathfrak{a} \subset A_{b^*_{\alpha} N} M \cap V(a(N)'_{a \otimes_{\beta} N^o} 1) V^*$.

As $a(N)'_{a \otimes_{\beta} N^o} 1 \subset \mathcal{L}(\mathfrak{H})_{a^*_{\beta} N^o} \widehat{M}'$, using the first formula, we get :

$$Sat \mathfrak{a} = A_{b^*_{\alpha} N} M \cap V(a(N)'_{a \otimes_{\beta} N^o} 1) V^*$$

And, as $V(a(N)'_{a \otimes_{\beta} N^o} 1) V^* \subset \mathcal{L}(\mathfrak{H})_{b^*_{\alpha} N} M$, we get the second formula. \square

7.6. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a \mathfrak{H}_b$ a $N - N$ bimodule, A a von Neumann algebra such that $b(N) \subset A \subset a(N)'$, V a corepresentation of \mathfrak{G} on ${}_a \mathfrak{H}_b$ which implements an action (b, \mathfrak{a}) of \mathfrak{G} on A and an action (a, \mathfrak{a}') of \mathfrak{G}^o on A' . Then, are equivalent :

- (i) \mathfrak{a} is saturated;
- (ii) \mathfrak{a}' satisfies property (A).

Proof. Using 7.5, we get that \mathfrak{a} is saturated if and only if :

$$V(A_{a \otimes_{\beta} N^o} 1) V^* = A_{b^*_{\alpha} N} \mathcal{L}(H_{\Phi}) \cap V(a(N)'_{a \otimes_{\beta} N^o} 1) V^*$$

which, taking the commutants, is equivalent to :

$$V(A'_{a^*_{\beta} N^o} \mathcal{L}(H_{\Phi})) V^* = (A'_{b \otimes_{\alpha} N} 1 \cup V(a(N)_{a^*_{\beta} N^o} \mathcal{L}(H_{\Phi})) V^*)''$$

or, to :

$$A'_{a^*_{\beta} N^o} \mathcal{L}(H_{\Phi}) = (V^*(A'_{b \otimes_{\alpha} N} 1) V \cup a(N)_{a^*_{\beta} N^o} \mathcal{L}(H_{\Phi}))''$$

which is :

$$A'_{a^*_{\beta} N^o} \mathcal{L}(H_{\Phi}) = (\mathfrak{a}'(A') \cup (1_{a \otimes_{\beta} N^o} \beta(N)'))''$$

which means that \mathfrak{a}' satisfies property (A). The converse is proved the same way. \square

7.7. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, A a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A ; let ψ be a normal semi-finite faithful weight on A , and let us suppose that there exists a corepresentation V on H_ψ which is a standard implementation for \mathfrak{a} in the sense of 6.9. Then, are equivalent :

- (i) \mathfrak{a} is saturated;
- (ii) \mathfrak{a} satisfies property (A).

Proof. By 6.9, for all $n \in N$, we write $a(n) = J_\psi b(n^*) J_\psi$, and V is a corepresentation of \mathfrak{G} on ${}_a(H_\psi)_b$, which implements \mathfrak{a} , and verifies :

$$V^* = (J_\psi \alpha \otimes_{\beta} J_{\widehat{\Phi}}) V (J_\psi b \otimes_{\alpha} J_{\widehat{\Phi}})$$

from which we get that the action \mathfrak{a}' implemented by V^* satisfies :

$$(J_\psi \alpha \otimes_{\beta} J_{\widehat{\Phi}}) \mathfrak{a}'(A') (J_\psi b \otimes_{\alpha} J_{\widehat{\Phi}}) = \mathfrak{a}(A)$$

As $(J_\psi \alpha \otimes_{\beta} J_{\widehat{\Phi}})(A \otimes_{\beta} 1)(J_\psi b \otimes_{\alpha} J_{\widehat{\Phi}}) = A' \otimes_{\alpha} 1$, we get that :

$$(J_\psi \alpha \otimes_{\beta} J_{\widehat{\Phi}})(A' \otimes_{\beta} \mathcal{L}(H_\Phi))(J_\psi b \otimes_{\alpha} J_{\widehat{\Phi}}) = A \otimes_{\alpha} \mathcal{L}(H_\Phi)$$

Thanks to 7.6, \mathfrak{a} is saturated if and only if \mathfrak{a}' satisfies property (A), which means that :

$$A' \otimes_{\beta} \mathcal{L}(H_\Phi) = (\mathfrak{a}'(A') \cup (1 \otimes_{\beta} \beta(N'))''$$

from which we get that :

$$A \otimes_{\alpha} \mathcal{L}(H_\Phi) = (\mathfrak{a}(A) \cup (1 \otimes_{\alpha} \alpha(N'))''$$

which means that \mathfrak{a} satisfies property (A). □

7.8. Corollary. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; then, we have :

$$\Gamma(M) = \{X \in M \otimes_{\beta} M; (\Gamma \otimes_{\alpha} id)(X) = (id \otimes_{\beta} \Gamma)(X)\}$$

Proof. We have seen in 6.10 that Γ can be considered as an action of \mathfrak{G} on M , which has a standard implementation; moreover, by 7.2, it satisfies property (A), and, therefore, by 7.7, this action is saturated, which finishes the proof. □

8. THE STANDARD IMPLEMENTATION OF AN ACTION : THE CASE OF A δ -INVARIANT WEIGHT

In this chapter, we define δ -invariant weights on a von Neumann algebra on which is acting a measured quantum groupoid (8.1); in that situation, we construct (8.8) a standard implementation of the action. As a corollary, we obtain in that case that the property A and the saturation property introduced in chapter 7 are equivalent (8.10).

8.1. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A and let ψ be a normal faithful semi-finite weight on A . We shall identify A and $\pi_\psi(A)$, for simplification. Let a be the representation of N on H_ψ given by :

$$a(n) = J_\psi b(n^*) J_\psi$$

We shall say that ψ is δ -invariant if, for all $\eta \in D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}(\delta^{1/2})$, such that $\delta^{1/2}\eta$ belongs to $D((H_\Phi)_\beta, \nu^o)$, and for all $x \in \mathfrak{N}_\psi$, we have:

$$\psi((id_{b^*_\alpha \omega_\eta})\mathbf{a}(x^*x)) = \|\Lambda_\psi(x)_{\alpha \otimes_\beta \delta^{1/2}\eta}\|_{\nu^o}^2$$

One should remark that $(id_{b^*_\alpha \omega_\eta})\mathbf{a}(x^*x)$ belongs then to \mathfrak{M}_ψ^+ .

Moreover, if the subset $D((H_\psi)_b, \nu^o) \cap D({}_\alpha H_\psi, \nu)$ is dense in H_ψ , we shall say that ψ bears the density property,

8.2. Example. Let us consider (6.10) the action (β, Γ) of \mathfrak{G} on M ; then, by 4.5, we get that Φ is δ -invariant. Namely, taking $(e_i)_{i \in I}$ an orthonormal (α, ν) basis of H_Φ , we get, for $x \in \mathfrak{N}_\Phi$ and convenient η , that :

$$\begin{aligned} \Phi[(id_{\beta^*_\alpha \omega_\eta})\Gamma(x^*x)] &= \sum_i \Phi((id_{\beta^*_\alpha \omega_{\eta, e_i}})\Gamma(x)^*(id_{\beta^*_\alpha \omega_{\eta, e_i}})\Gamma(x)) \\ &= \sum_i \|(id * \omega_{\delta^{1/2}\eta, e_i})(\widehat{W}^o)\Lambda_\Phi(x)\|^2 \\ &= \|\sum_i (1_{\beta \otimes_\alpha \theta^{\alpha, \nu}(e_i, e_i)})\widehat{W}^o(\Lambda_\Phi(x)_{\alpha \otimes_\beta \delta^{1/2}\eta})\|^2 \\ &= \|\Lambda_\Phi(x)_{\alpha \otimes_\beta \delta^{1/2}\eta}\|_{\nu^o}^2 \end{aligned}$$

Moreover, as $D((H_\Phi)_\beta, \nu^o) \cap D({}_\alpha H_\Phi, \nu)$ is dense in H_Φ , Φ bears the density property.

8.3. Lemma. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let $(e_i)_{i \in I}$ be an (α, ν) -orthogonal basis of H_Φ . Let $x \in \mathfrak{N}_\psi$, and η as in 8.1. Then:

- (i) for all $\xi \in D({}_\alpha H_\Phi, \nu)$, $(id_{b^*_\alpha \omega_{\eta, \xi}})\mathbf{a}(x)$ belongs to \mathfrak{N}_ψ ;
- (ii) the sum $\sum_i \Lambda_\psi((id_{b^*_\alpha \omega_{\eta, e_i}})\mathbf{a}(x))_{b \otimes_\alpha e_i}$ is strongly converging, the limit does not depend upon the choice of the (α, ν) -orthogonal basis of H_Φ , and :

$$\|\sum_i \Lambda_\psi((id_{b^*_\alpha \omega_{\eta, e_i}})\mathbf{a}(x))_{b \otimes_\alpha e_i}\|^2 = \psi((id_{b^*_\alpha \omega_\eta})\mathbf{a}(x^*x))$$

Proof. We have :

$$(id_{b^*_\alpha \omega_{\eta, \xi}})\mathbf{a}(x)^*(id_{b^*_\alpha \omega_{\eta, \xi}})\mathbf{a}(x) = (id_{b^*_\alpha \omega_\eta})(\mathbf{a}(x^*)(1_{b \otimes_\alpha \theta^{\alpha, \nu}(\xi, \xi)})\mathbf{a}(x))$$

which is less than $\|R^{\alpha, \nu}(\xi)\|^2(id_{b^*_\alpha \omega_\eta})\mathbf{a}(x^*x)$, which belongs to \mathfrak{M}_ψ^+ , by hypothesis; so we get (i).

As the vectors $\Lambda_\psi((id_{b^*_\alpha \omega_{\eta, e_i}})\mathbf{a}(x))_{b \otimes_\alpha e_i}$ are two by two orthogonal, for any finite $J \subset I$, we have :

$$\begin{aligned} \|\sum_{i \in J} \Lambda_\psi((id_{b^*_\alpha \omega_{\eta, e_i}})\mathbf{a}(x))_{b \otimes_\alpha e_i}\|^2 &= \sum_{i \in J} \|\Lambda_\psi((id_{b^*_\alpha \omega_{\eta, e_i}})\mathbf{a}(x))_{b \otimes_\alpha e_i}\|^2 \\ &= \psi(id_{b^*_\alpha \omega_\eta})(\mathbf{a}(x^*)(1_{b \otimes_\alpha \sum_{i \in J} \theta^{\alpha, \nu}(e_i, e_i)})\mathbf{a}(x)) \end{aligned}$$

which is less than (and uprising to) $\psi(id_{b^*_\alpha \omega_\eta} \mathbf{a}(x^*x))$. If $\xi \in D({}_\alpha H_\Phi, \nu)$, we have :

$$(\rho_\xi^{b,\alpha})^* \left(\sum_i \Lambda_\psi((id_{b^*_\alpha \omega_{\eta,e_i}}) \mathbf{a}(x))_{b \otimes_\alpha e_i} \right) = \Lambda_\psi((id_{b^*_\alpha \omega_{\eta,\xi}}) \mathbf{a}(x))$$

which does not depend upon the choice of the basis, and finishes the proof of (ii). \square

8.4. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_ψ by, for $n \in N$:*

$$a(n) = J_\psi b(n^*) J_\psi$$

Then :

(i) *there exists an isometry V_ψ from $H_\psi \otimes_{\nu^o} H_\Phi$ to $H_\psi \otimes_\alpha H_\Phi$ such that:*

$$V_\psi(\Lambda_\psi(x) \otimes_{\nu^o} \delta^{1/2} \eta) = \sum_i \Lambda_\psi((id_{b^*_\alpha \omega_{\eta,e_i}}) \mathbf{a}(x)) \otimes_\alpha e_i$$

for all $x \in \mathfrak{N}_\psi$, η as in 8.1 and for all (α, ν) -orthogonal basis $(e_i)_{i \in I}$ of H_Φ .

(ii) *for all $e \in \mathfrak{N}_\psi$, we have :*

$$(J_\psi e J_\psi \otimes_\nu 1) V_\psi(\Lambda_\psi(x) \otimes_{\nu^o} \delta^{1/2} \eta) = \mathbf{a}(x) (J_\psi \Lambda_\psi(e) \otimes_\nu \eta)$$

(iii) *for all $\xi \in D({}_\alpha H_\Phi, \nu)$, we have :*

$$\Lambda_\psi((id_{b^*_\alpha \omega_{\eta,\xi}}) \mathbf{a}(x)) = (id * \omega_{\delta^{1/2} \eta, \xi})(V_\psi) \Lambda_\psi(x)$$

(iv) *for all $y \in A$, $z \in M'$, we have :*

$$\mathbf{a}(y) V_\psi = V_\psi (y \otimes_{\nu^o} 1)$$

$$(1 \otimes_N z) V_\psi = V_\psi (1 \otimes_{N^o} z)$$

(v) *for all $n \in N$; we have :*

$$(a(n) \otimes_N 1) V_\psi = V_\psi (1 \otimes_{N^o} \alpha(n))$$

$$(1 \otimes_N \beta(n)) V_\psi = V_\psi (b(n) \otimes_{N^o} 1)$$

$$(1 \otimes_N \hat{\beta}(n)) V_\psi = V_\psi (1 \otimes_{N^o} \hat{\beta}(n))$$

Proof. Defining V_ψ by the formula given in (i), we easily obtain, using 8.3(ii), that for all $x' \in \mathfrak{N}_\psi$, η' as in 8.1, we get :

$$(V_\psi(\Lambda_\psi(x) \otimes_{\nu^o} \delta^{1/2} \eta) | V_\psi(\Lambda_\psi(x') \otimes_{\nu^o} \delta^{1/2} \eta')) = \psi((id_{b^*_\alpha \omega_{\eta,\eta'}}) \mathbf{a}(x'^* x))$$

which, by polarization of the definition of δ -invariance, is equal to:

$$(\Lambda_\psi(x) \otimes_{\nu^o} \delta^{1/2} \eta | \Lambda_\psi(x') \otimes_{\nu^o} \delta^{1/2} \eta')$$

which implies that this formula defines an isometry which can be extended by continuity to $H_{\psi} \otimes_{\beta} H_{\Phi}$, and does not depend upon the choice of the basis, which is (i). We then get :

$$\begin{aligned}
(J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) &= \sum_i J_{\psi} e J_{\psi} \Lambda_{\psi}((id \otimes_{N} \omega_{\eta, e_i}) \mathbf{a}(x)) \otimes_{\alpha} e_i \\
&= \sum_i (id \otimes_{N} \omega_{\eta, e_i}) \mathbf{a}(x) J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} e_i \\
&= \mathbf{a}(x) (J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} \eta)
\end{aligned}$$

which is (ii). Using (ii), we then get :

$$\begin{aligned}
J_{\psi} e J_{\psi} \Lambda_{\psi}((id \otimes_{N} \omega_{\eta, \xi}) \mathbf{a}(x)) &= (id \otimes_{N} \omega_{\eta, \xi}) \mathbf{a}(x) J_{\psi} \Lambda_{\psi}(e) \\
&= J_{\psi} e J_{\psi} (id * \omega_{\delta^{1/2} \eta, \xi}) (V_{\psi}) \Lambda_{\psi}(x)
\end{aligned}$$

from which we get (iii), by taking the limit when e goes to 1. Using (ii) again, we have also :

$$\begin{aligned}
(J_{\psi} e J_{\psi} \otimes_{\alpha} 1) \mathbf{a}(y) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) &= \mathbf{a}(y) (J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) \\
&= \mathbf{a}(yx) (J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} \eta)
\end{aligned}$$

which, thanks to (ii) again, is equal to :

$$(J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(yx) \otimes_{\beta} \delta^{1/2} \eta) = (J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(y \otimes_{\beta} 1) (\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta)$$

from which we get the first formula of (iv), by taking the limit when e goes to 1. We have also :

$$\begin{aligned}
(J_{\psi} e J_{\psi} \otimes_{\alpha} z) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) &= (1 \otimes_{\alpha} z) \mathbf{a}(x) (J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} \eta) \\
&= \mathbf{a}(x) (J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} z \eta)
\end{aligned}$$

and, thanks to (ii) again, we get :

$$\begin{aligned}
(J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} z \eta) &= (J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} z \delta^{1/2} \eta) \\
&= (J_{\psi} e J_{\psi} \otimes_{\alpha} 1) V_{\psi}(1 \otimes_{\beta} z) (\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta)
\end{aligned}$$

from which we get the second formula of (iv), by taking the limit when e goes to 1.

Let's take now $n \in N$, analytic with respect both to σ^{ν} and γ . We have :

$$\begin{aligned}
(J_{\psi} e J_{\psi} a(n) \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) &= (J_{\psi} e b(n^*) J_{\psi} \otimes_{\alpha} 1) V_{\psi}(\Lambda_{\psi}(x) \otimes_{\beta} \delta^{1/2} \eta) \\
&= \mathbf{a}(x) (J_{\psi} \Lambda_{\psi}(e b(n^*)) \otimes_{\alpha} \eta)
\end{aligned}$$

It is equal to :

$$\mathbf{a}(x) (\sigma_{-i/2}^{\psi}(b(n)) J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} \eta) = \mathbf{a}(x) (b(\gamma_{-i/2}(n)) J_{\psi} \Lambda_{\psi}(e) \otimes_{\alpha} \eta)$$

which is equal to $\mathbf{a}(x)(J_\psi \Lambda_\psi(e) {}_{b \otimes_\alpha} \alpha(\sigma_{-i/2}^\nu \gamma_{-i/2}(n))\eta)$, and, thanks to (ii) again, to :

$$\begin{aligned} & (J_\psi e J_\psi {}_{b \otimes_\alpha} 1) V_\psi(\Lambda_\psi(x) {}_{a \otimes_\beta} \delta^{1/2} \alpha(\sigma_{-i/2}^\nu \gamma_{-i/2}(n))\eta) \\ &= (J_\psi e J_\psi {}_{b \otimes_\alpha} 1) V_\psi(\Lambda_\psi(x) {}_{a \otimes_\beta} \alpha(n) \delta^{1/2} \eta) \\ &= (J_\psi e J_\psi {}_{b \otimes_\alpha} 1) V_\psi(1 {}_{a \otimes_\beta} \alpha(n)) (\Lambda_\psi(x) {}_{a \otimes_\beta} \delta^{1/2} \eta) \end{aligned}$$

from which we get the first formula of (v), by taking the limit when e goes to 1.

The other formulae of (v) are special cases of (iv), taking $y = b(n)$, and $z = \hat{\beta}(n)$. \square

8.5. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_ψ by, for $n \in N$:*

$$a(n) = J_\psi b(n^*) J_\psi$$

and let V_ψ the isometry from $H_\psi {}_{a \otimes_\beta} H_\Phi$ to $H_\psi {}_{b \otimes_\alpha} H_\Phi$ constructed in 8.4, which satisfies the intertwining properties proved in 8.4(v). Then we have :

$$(1 {}_{b \otimes_\alpha} \sigma_{\nu^\circ} W^* \sigma_\nu) (V_\psi {}_{\beta \otimes_\alpha} 1) (1 {}_{a \otimes_\beta} \sigma_{\nu^\circ} W \sigma_\nu) = (1 {}_{b \otimes_\alpha} \sigma_{\nu^\circ}) (V_\psi {}_{b \otimes_\alpha} 1) \sigma_{b,a}^{2,3} (V_\psi {}_{\alpha \otimes_\beta} 1)$$

Proof. Let x_1, x_2 in \mathfrak{N}_ψ , ξ_1, ξ_2 in $D({}_\alpha H_\Phi, \nu)$, and η_1, η_2 as in 8.1 ($i = 1, 2$); then, the scalar product of the vector :

$$(V_\psi {}_{b \otimes_\alpha} 1) (1 {}_{a \otimes_\beta} \sigma_{\nu^\circ}) (V_\psi {}_{\alpha \otimes_\beta} 1) (1 {}_{\alpha \otimes_\beta} \sigma_\nu) (\Lambda_\psi(x_1) {}_{a \otimes_\beta} (\delta^{1/2} \eta_1 {}_{\beta \otimes_\alpha} \delta^{1/2} \eta_2))$$

with $\Lambda_\psi(x_2) {}_{b \otimes_\alpha} \xi_1 {}_{\beta \otimes_\alpha} \xi_2$ is equal to :

$$(V_\psi((id * \omega_{\delta^{1/2} \eta_2, \xi_2})) (V_\psi) \Lambda_\psi(x_1) {}_{a \otimes_\beta} \delta^{1/2} \eta_1) | \Lambda_\psi(x_2) {}_{b \otimes_\alpha} \xi_1)$$

which, thanks to 8.4(iii), is equal to :

$$(V_\psi(\Lambda_\psi[(id {}_{b \otimes_\alpha} \omega_{\eta_2, \xi_2}) \mathbf{a}(x_1)] {}_{a \otimes_\beta} \delta^{1/2} \eta_1) | \Lambda_\psi(x_2) {}_{b \otimes_\alpha} \xi_1)$$

and, using 8.4(iii) again, is equal to :

$$(\Lambda_\psi[(id {}_{b \otimes_\alpha} \omega_{\eta_1, \xi_1}) \mathbf{a}((id {}_{b \otimes_\alpha} \omega_{\eta_2, \xi_2}) \mathbf{a}(x_1))] | \Lambda_\psi(x_2))$$

which, using 6.1, is equal to :

$$(\Lambda_\psi((id {}_{b \otimes_\alpha} \omega_{\eta_1, \xi_1} {}_{\beta \otimes_\alpha} \omega_{\eta_2, \xi_2}) (id {}_{\beta \otimes_\alpha} \Gamma) \mathbf{a}(x_1)) | \Lambda_\psi(x_2))$$

Let us define $\Omega \in M_*$ such that, for any $x \in M$, $\Omega(x) = (\omega_{\eta_1, \xi_1} {}_{\beta \otimes_\alpha} \omega_{\eta_2, \xi_2}) \Gamma(x)$.

Let $(e_i)_{i \in I}$ be an (α, ν) -orthogonal basis of H_Φ . There exists a family of vectors ξ_i in H , such that $\xi_i = \hat{\beta}(\langle e_i, e_i \rangle_{\alpha, \nu}) \xi_i$, and :

$$W(\xi_1 {}_{\beta \otimes_\alpha} \xi_2) = \sum_i e_i {}_{\alpha \otimes_\beta} \xi_i$$

Moreover, as, for $n \in \mathfrak{N}_\nu$:

$$\begin{aligned} \sum_i \|\alpha(n)\xi_i\|^2 &= \|(1_{\alpha \otimes_{\hat{\beta}} \alpha(n)})W(\xi_{1 \beta \otimes_\alpha \xi_2})\|^2 \\ &= \|W(\alpha(n)\xi_{1 \beta \otimes_\alpha \xi_2})\|^2 \\ &\leq \|R^{\alpha, \nu}(\xi_2)\|^2 \|R^{\alpha, \nu}(\xi_1)\|^2 \|\Lambda_\nu(n)\|^2 \end{aligned}$$

we get that the vectors ξ_i belong to $D({}_\alpha H_\Phi, \nu)$. The same way, there exist vectors $\eta_i = \hat{\beta}(< e_i, e_i >_{\alpha, \nu})\eta_i \in D({}_\alpha H_\Phi, \nu)$, such that :

$$W(\eta_{1 \beta \otimes_\alpha \eta_2}) = \sum_i e_i \alpha \otimes_{\hat{\beta}} \eta_i$$

Moreover, as we have $\Gamma(\delta^{it}) = \delta^{it} \beta \otimes_\alpha \delta^{it}$, we get that the vectors η_i belongs to $\mathcal{D}(\delta^{1/2})$, and that :

$$W(\delta^{1/2} \eta_{1 \beta \otimes_\alpha \delta^{1/2} \eta_2}) = \sum_i e_i \alpha \otimes_{\hat{\beta}} \delta^{1/2} \eta_i$$

We have also :

$$\begin{aligned} \sum_i \|\beta(n)\delta^{1/2}\eta_i\|^2 &= \|(1_{\alpha \otimes_{\hat{\beta}} \beta(n)})W(\delta^{1/2} \eta_{1 \beta \otimes_\alpha \delta^{1/2} \eta_2})\|^2 \\ &\leq \|R^{\beta, \nu^o}(\delta^{1/2} \eta_1)\|^2 \|R^{\beta, \nu^o}(\delta^{1/2} \eta_2)\|^2 \|\Lambda_\nu(n)\|^2 \end{aligned}$$

from which we get that the vectors $\delta^{1/2}\eta_i$ belong to $D((H_\Phi)_\beta, \nu^o)$. From these results, we get that, for any $x \in M$, we have $\Omega(x) = \sum_i \omega_{\eta_i, \xi_i}(x)$, and, therefore, we have :

$$(\Lambda_\psi((id_{b^* \alpha} \omega_{\eta_1, \xi_1} \beta^* \alpha \omega_{\eta_2, \xi_2})(id_{\beta^* \alpha} \Gamma) \mathbf{a}(x_1)) | \Lambda_\psi(x_2)) = \sum_i (\Lambda_\psi((id_{b^* \alpha} \omega_{\eta_i, \xi_i}) \mathbf{a}(x_1)) | \Lambda_\psi(x_2))$$

The properties of the ξ_i s and the η_i s allow us to use again 8.5(iii), and this is equal to :

$$\sum_i ((id * \omega_{\delta^{-1/2} \eta_i, \xi_i})(V_\psi) \Lambda_\psi(x_1) | \Lambda_\psi(x_2))$$

On the other hand, if ζ_1 belongs to $D({}_\alpha H_\psi, \nu)$ and ζ_2 to $D((H_\psi)_b, \nu^o)$, we get, by 8.5(iv), that $(\omega_{\zeta_1, \zeta_2} * id)(V)$ belongs to M , and that :

$$\begin{aligned} \sum_i ((id * \omega_{\delta^{1/2} \eta_i, \xi_i})(V_\psi) \zeta_1 | \zeta_2) &= \sum_i \omega_{\delta^{1/2} \eta_i, \xi_i}(\omega_{\zeta_1, \zeta_2} * id)(V_\psi) \\ &= (\omega_{\delta^{1/2} \eta_1, \xi_1} \beta^* \alpha \omega_{\delta^{1/2} \eta_2, \xi_2}) \Gamma[(\omega_{\zeta_1, \zeta_2} * id)(V_\psi)] \end{aligned}$$

is equal to :

$$(W^*((1_{\alpha \otimes_{\hat{\beta}}} (\omega_{\zeta_1, \zeta_2} * id)(V_\psi))W(\delta^{1/2} \eta_{1 \beta \otimes_\alpha \delta^{1/2} \eta_2}) | \xi_{1 \beta \otimes_\alpha \xi_2}))$$

which is equal to :

$$((1_{b \otimes_\alpha \sigma_{\nu^o}} W^* \sigma_\nu)(V_\psi \beta \otimes_\alpha 1)(1_{a \otimes_\beta \sigma_{\nu^o}} W \sigma_\nu)(\zeta_{1 a \otimes_\beta \delta^{1/2} \eta_2 \alpha \otimes_\beta \delta^{1/2} \eta_1} | \zeta_{2 a \otimes_\beta \xi_2 \alpha \otimes_\beta \xi_1}))$$

From which, by continuity, we get that the scalar product of the vector :

$$(V_\psi b \otimes_\alpha 1) \sigma_{b,a}^{2,3} (V_\psi \alpha \otimes_\beta 1) (1_{\alpha \otimes_\beta \sigma_\nu}) (\Lambda_\psi(x_1) \alpha \otimes_\beta (\delta^{1/2} \eta_{1 \beta \otimes_\alpha \delta^{1/2} \eta_2}))$$

with $\Lambda_\psi(x_2) {}_{b \otimes_\alpha} \xi_1 {}_{\beta \otimes_\alpha} \xi_2$ is equal to the scalar product of the vector :

$$(1 {}_{b \otimes_\alpha} \sigma_{\nu^o} W^* \sigma_\nu) (V_\psi {}_{\hat{\beta} \otimes_\alpha} 1) (1 {}_{a \otimes_\beta} \sigma_{\nu^o} W \sigma_\nu) (\Lambda_\psi(x_1) {}_{a \otimes_\beta} \delta^{1/2} \eta_2 {}_{\alpha \otimes_\beta} \delta^{1/2} \eta_1)$$

with $\Lambda_\psi(x_2) {}_{b \otimes_\alpha} \xi_1 {}_{\beta \otimes_\alpha} \xi_2$, from which, by continuity, we get the result. \square

8.6. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_ψ by, for $n \in N$:*

$$a(n) = J_\psi b(n^*) J_\psi$$

and let V_ψ the isometry from $H_\psi {}_{a \otimes_\beta} H_\Phi$ to $H_\psi {}_{b \otimes_\alpha} H_\Phi$ constructed in 8.4; then :

(i) V_ψ is a unitary, and, for all x in A , we have $\mathfrak{a}(x) = V_\psi(x {}_{a \otimes_\beta} 1) V_\psi^*$;

(ii) V_ψ is a corepresentation of \mathfrak{G} on the bimodule ${}_a(H_\psi)_b$ which implements \mathfrak{a} .

Proof. Thanks to 8.4(i), (v), (iv) and 8.5, we can apply 5.4, which says that V is a unitary and a corepresentation, which gives (ii), and (i) by 8.4(iv) again. \square

8.7. Lemma. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_ψ such that, for $n \in N$:*

$$a(n) = J_\psi b(n^*) J_\psi$$

and let us suppose that ψ bears the density property, as defined in 8.1. Let V_ψ be the corepresentation of \mathfrak{G} on the bimodule ${}_a(H_\psi)_b$ constructed in 8.6. Then, for all x, y in $\mathfrak{N}_\psi \cap \mathfrak{N}_\psi^*$, analytic with respect to ψ , and ξ in $D((H_\Phi)_\beta, \nu^o) \cap \mathcal{D}((P\delta)^{-1})$ such that $(P\delta)^{-1}\xi$ belongs to $D((H_\Phi)_\beta, \nu^o)$ and η in $D({}_\alpha H_\Phi, \nu) \cap \mathcal{D}((P\delta)^{-1})$ such that $(P\delta)^{-1}\eta$ belongs to $D({}_\alpha H_\Phi, \nu)$, we have :

$$(V_\psi(\Delta_\psi \Lambda_\psi(x) {}_{a \otimes_\beta} (P\delta)^{-1}\xi) | \Lambda_\psi(y) {}_{b \otimes_\alpha} \eta) = (V_\psi(\Lambda_\psi(x) {}_{a \otimes_\beta} \xi) | \Delta_\psi \Lambda_\psi(y) {}_{b \otimes_\alpha} (P\delta)^{-1}\eta)$$

Proof. Thanks to the density property, we can use 5.11; therefore, with ξ, η in \mathcal{E}_τ , using 4.4(iii) :

$$\begin{aligned} & (V_\psi(\Delta_\psi \Lambda_\psi(x) {}_{a \otimes_\beta} P^{-1}\delta^{-1}\xi) | \Lambda_\psi(y) {}_{b \otimes_\alpha} \eta) \\ &= (V_\psi^*(\Delta_\psi \Lambda_\psi(x) {}_{b \otimes_\alpha} J_{\hat{\Phi}} P^{-1/2}\eta) | \Lambda_\psi(y) {}_{a \otimes_\beta} J_{\hat{\Phi}} P^{-1/2}\delta^{-1}\xi) \\ &= (\Delta_\psi \Lambda_\psi(x) {}_{b \otimes_\alpha} J_{\hat{\Phi}} P^{-1/2}\eta | V_\psi(\Lambda_\psi(y) {}_{a \otimes_\beta} J_{\hat{\Phi}} P^{-1/2}\delta^{-1}\xi)) \end{aligned}$$

which, thanks to 8.4(iii), is equal to :

$$\begin{aligned} & (\Delta_\psi \Lambda_\psi(x) | (id * \omega_{J_{\hat{\Phi}} P^{-1/2}\delta^{-1}\xi, J_{\hat{\Phi}} P^{-1/2}\eta})(V_\psi) \Lambda_\psi(y)) \\ &= (\Delta_\psi \Lambda_\psi(x) | \Lambda_\psi[(id * \omega_{J_{\hat{\Phi}} P^{-1/2}\delta^{-1/2}\xi, J_{\hat{\Phi}} P^{-1/2}\eta}) \mathfrak{a}(y)]) \\ &= (\Lambda_\psi[(id * \omega_{J_{\hat{\Phi}} P^{-1/2}\eta, J_{\hat{\Phi}} P^{-1/2}\delta^{-1/2}\xi}) \mathfrak{a}(y^*)] | \Lambda_\psi(x^*)) \end{aligned}$$

which, using again 8.4(iii), is equal to :

$$\begin{aligned}
& ((id * \omega_{J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \eta, J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \xi})(V_\psi) \Lambda_\psi(y^*) | \Lambda_\psi(x^*)) \\
&= (V_\psi(\Lambda_\psi(y^*)_{a \otimes_{\beta} J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \eta}) | \Lambda_\psi(x^*)_{b \otimes_{\alpha} J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \xi}) \\
&= (\Lambda_\psi(y^*)_{a \otimes_{\beta} J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \eta} | V_\psi^*(\Lambda_\psi(x^*)_{b \otimes_{\alpha} J_{\widehat{\mathfrak{F}}} P^{-1/2} \delta^{-1/2} \xi}))
\end{aligned}$$

which, using again 5.11, is equal to :

$$(\Lambda_\psi(y^*)_{b \otimes_{\alpha} \delta^{-1/2} \xi} | V_\psi(\Lambda_\psi(x^*)_{a \otimes_{\beta} P^{-1} \delta^{-1/2} \eta})) = (\Lambda_\psi(y^*) | (id * \omega_{P^{-1} \delta^{-1/2} \eta, \delta^{-1/2} \xi})(V_\psi) \Lambda_\psi(x^*))$$

which, using again twice 8.4(iii), is equal to :

$$\begin{aligned}
(\Lambda_\psi(y^*) | \Lambda_\psi[(id * \omega_{(P\delta)^{-1} \eta, \delta^{-1/2} \xi}) \mathbf{a}(x^*)]) &= (S_\psi \Lambda_\psi(y) | S_\psi \Lambda_\psi[(id * \omega_{\delta^{-1/2} \xi, (P\delta)^{-1} \eta}) \mathbf{a}(x)]) \\
&= (\Delta_\psi \Lambda_\psi[(id * \omega_{\delta^{-1/2} \xi, (P\delta)^{-1} \eta}) \mathbf{a}(x)] | \Lambda_\psi(y)) \\
&= (\Delta_\psi(id * \omega_{\xi, (P\delta)^{-1} \eta})(V_\psi) \Lambda_\psi(x) | \Lambda_\psi(y)) \\
&= (V_\psi(\Lambda_\psi(x)_{a \otimes_{\beta} \xi}) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_{\alpha} (P\delta)^{-1} \eta})
\end{aligned}$$

from which we get the result, thanks to 4.4(i). \square

8.8. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_ψ such that, for $n \in N$:*

$$a(n) = J_\psi b(n^*) J_\psi$$

and let us suppose that ψ bears the density property, as defined in 8.1. Let V_ψ the corepresentation of \mathfrak{G} on the bimodule ${}_a(H_\psi)_b$ constructed in 8.6. Then :

- (i) for all $t \in \mathbb{R}$ and $n \in N$, we have $\sigma_t^\psi(b(n)) = b(\gamma_t(n))$.
- (ii) it is possible to define two one parameter groups of unitaries $\Delta_\psi^{it} {}_{a \otimes_{\beta} N^o} \delta^{-it} P^{-it}$ and $\Delta_\psi^{it} {}_{b \otimes_{\alpha} N} \delta^{-it} P^{-it}$ with natural values on elementary tensor products, and we have :

$$V_\psi(\Delta_\psi^{it} {}_{a \otimes_{\beta} N^o} \delta^{-it} P^{-it}) = (\Delta_\psi^{it} {}_{b \otimes_{\alpha} N} \delta^{-it} P^{-it}) V_\psi$$

- (iii) for every x in A , $t \in \mathbb{R}$, we have :

$$\mathbf{a}(\sigma_t^\psi(x)) = (\sigma_t^\psi {}_{b \otimes_{\alpha} N} \tau_{-t} \sigma_{-t}^{\Phi \circ R} \sigma_t^\Phi) \mathbf{a}(x)$$

- (iv) we have :

$$V_\psi = (J_\psi {}_{b \otimes_{\alpha} N} J_{\widehat{\mathfrak{F}}}) V_\psi^* (J_\psi {}_{b \otimes_{\alpha} N} J_{\widehat{\mathfrak{F}}})$$

and, therefore, V_ψ is a standard implementation of \mathbf{a} .

Proof. Let x, y in $\mathfrak{N}_\psi \cap \mathfrak{N}_\psi^*$, analytic with respect to ψ ; let ξ, η as in 8.7, n in N analytic with respect both to σ_t^ν and γ_t ; we have :

$$\begin{aligned}
& (V_\psi(\Delta_\psi \Lambda_\psi(b(n)x)_{a \otimes_{\beta} (P\delta)^{-1} \xi}) | \Lambda_\psi(y)_{b \otimes_{\alpha} \eta}) \\
&= (V_\psi(\Lambda_\psi(b(n)x)_{a \otimes_{\beta} \xi}) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_{\alpha} (P\delta)^{-1} \eta}) \\
&= (V_\psi(b(n) \Lambda_\psi(x)_{a \otimes_{\beta} \xi}) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_{\alpha} (P\delta)^{-1} \eta})
\end{aligned}$$

which, using 8.4(v), is equal to :

$$\begin{aligned} ((1_{b \otimes_{N^o} \alpha} \beta(n)) V_\psi(\Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} \xi) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_\nu \alpha} (P\delta)^{-1} \eta) \\ = (V_\psi(\Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} \xi) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_\nu \alpha} \beta(n^*) (P\delta)^{-1} \eta) \end{aligned}$$

but, we know that $\beta(n^*)(P\delta)^{-1}\eta = (P\delta)^{-1}\beta(\gamma_i(n^*))\eta$, and $\beta(\gamma_i(n^*))\eta$ bears the property necessary to apply 8.7; therefore, it is equal to :

$$\begin{aligned} (V_\psi(\Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} \xi) | \Delta_\psi \Lambda_\psi(y)_{b \otimes_\nu \alpha} (P\delta)^{-1} \beta(\gamma_i(n^*))\eta) \\ = (V_\psi(\Delta_\psi \Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} (P\delta)^{-1} \xi) | \Lambda_\psi(y)_{b \otimes_\nu \alpha} \beta(\gamma_i(n^*))\eta) \\ = ((1_{b \otimes_{N^o} \alpha} \beta(\gamma_{-i}(n)) V_\psi(\Delta_\psi \Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} (P\delta)^{-1} \xi) | \Lambda_\psi(y)_{b \otimes_\nu \alpha} \eta) \end{aligned}$$

which, using again 8.4(v) again, is equal to :

$$(V_\psi(b(\gamma_{-i}(n)) \Delta_\psi \Lambda_\psi(x)_{a \otimes_{\nu^o} \beta} (P\delta)^{-1} \xi) | \Lambda_\psi(y)_{b \otimes_\nu \alpha} \eta)$$

from which we get that :

$$\Delta_\psi \Lambda_\psi(b(n)x) = b(\gamma_{-i}(n)) \Delta_\psi \Lambda_\psi(x)$$

and we see that $b(n)$ belongs to $\mathcal{D}(\sigma_{-i}^\psi)$, and that $\sigma_{-i}^\psi(b(n)) = b(\gamma_{-i}(n))$. By classical approximations, we can prove this result for any n in $\mathcal{D}(\gamma_{-i})$, from which, using ([H], 4.3 and 4.4), one get (i).

Let $\xi \in D({}_a H_\psi)$; we have, for all $n \in \mathfrak{N}_\nu$, $t \in \mathbb{R}$:

$$\begin{aligned} a(n) \Delta_\psi^{it} \xi &= J_\psi b(n^*) J_\psi \Delta_\psi^{it} \xi \\ &= J_\psi b(n^*) \Delta_\psi^{it} J_\psi \xi \\ &= J_\psi \Delta_\psi^{it} \sigma_{-t}^\psi(b(n^*)) J_\psi \xi \\ &= J_\psi \Delta_\psi^{it} b(\gamma_{-t}(n^*)) J_\psi \xi \\ &= \Delta_\psi^{it} a(\gamma_{-t}(n)) \xi \\ &= \Delta_\psi^{it} R^{\alpha, \nu}(\xi) \Lambda_\nu(\gamma_{-t}(n)) \end{aligned}$$

If we define H^{it} by $H^{it} \Lambda_\nu(n) = \Lambda_\nu(\gamma_t(n))$, we get that $\Delta_\psi^{it} \xi$ belongs to $D({}_a H_\psi, \nu)$, and that $R^{\alpha, \nu}(\Delta_\psi^{it} \xi) = \Delta_\psi^{it} R^{\alpha, \nu}(\xi) H^{-it}$; then, we get, if $\xi' \in D({}_a H_\psi, \nu)$, that :

$$\langle \Delta_\psi^{it} \xi, \Delta_\psi^{it} \xi' \rangle_{\alpha, \nu^o}^o = \gamma_t(\langle \xi, \xi' \rangle_{\alpha, \nu^o}^o)$$

But, for all $n \in N$, we have:

$$\begin{aligned} \beta(\gamma_t(n)) &= \sigma_t^\Phi(\beta(n)) \\ &= \delta^{-it} \sigma_t^{\Phi \circ R}(\beta(n)) \delta^{it} \\ &= \delta^{-it} \beta(\sigma_{-t}^\nu(n)) \delta^{it} \\ &= \delta^{-it} \tau_{-t}(\beta(n)) \delta^{it} \\ &= (\delta P)^{-it} \beta(n) (\delta P)^{it} \end{aligned}$$

Therefore, it is possible to define a one-parameter group of unitaries $\Delta_\psi^{it} a \otimes_{N^o} \beta \delta^{-it} P^{-it}$ on $H_\psi a \otimes_{N^o} \beta H_\Phi$, with natural values on elementary tensor products.

The construction of $\Delta_\psi^{it} b \otimes_\alpha \delta^{-it} P^{-it}$ is done in a similar way.

Let $\Delta_{\psi \underset{\nu^o}{a} \otimes_{\beta} (P\delta)^{-1}}$ and $\Delta_{\psi \underset{\nu}{b} \otimes_{\alpha} (P\delta)^{-1}}$ be the analytic generators of these two one parameter automorphisms groups. We had got in 8.7 that :

$$V_{\psi}(\Delta_{\psi \underset{\nu^o}{a} \otimes_{\beta} (P\delta)^{-1}) \subset (\Delta_{\psi \underset{\nu}{b} \otimes_{\alpha} (P\delta)^{-1})V_{\psi}$$

from which we finish the proof of (ii). Then, we get (iii) from (ii) and 8.6(i).

We can verify that it is possible to define an antilinear unitary $J_{\psi \underset{\nu}{b} \otimes_{\alpha} J_{\widehat{\Phi}}}$ from $H_{\psi \underset{\nu}{b} \otimes_{\alpha} H_{\widehat{\Phi}}}$ onto $H_{\psi \underset{\nu^o}{a} \otimes_{\beta} H_{\widehat{\Phi}}}$, with natural values on elementary tensors, whose inverse is $J_{\psi \underset{\nu^o}{a} \otimes_{\beta} J_{\widehat{\Phi}}}$; moreover, we have, using 5.11 :

$$\begin{aligned} & ((J_{\psi \underset{\nu^o}{a} \otimes_{\beta} J_{\widehat{\Phi}}})V_{\psi}^*(J_{\psi \underset{\nu^o}{a} \otimes_{\beta} J_{\widehat{\Phi}}})(\Lambda_{\psi}(x) \underset{\nu^o}{a} \otimes_{\beta} \xi) | \Lambda_{\psi}(y) \underset{\nu}{b} \otimes_{\alpha} \eta) \\ &= \overline{(V_{\psi}^*(J_{\psi} \Lambda_{\psi}(x) \underset{\nu}{b} \otimes_{\alpha} J_{\widehat{\Phi}} \xi) | J_{\psi} \Lambda_{\psi}(y) \underset{\nu^o}{a} \otimes_{\beta} J_{\widehat{\Phi}} \eta)} \\ &= \overline{(V_{\psi}(J_{\psi} \Lambda_{\psi}(x) \underset{\nu^o}{a} \otimes_{\beta} P^{-1/2} \eta) | J_{\psi} \Lambda_{\psi}(y) \underset{\nu}{b} \otimes_{\alpha} P^{1/2} \xi)} \end{aligned}$$

which, using (ii), is equal to the scalar product of $J_{\psi} \Lambda_{\psi}(y) \underset{\nu}{b} \otimes_{\alpha} P^{1/2} \xi$ with :

$$(\Delta_{\psi \underset{\nu}{b} \otimes_{\alpha} \delta^{-1/2} P^{-1/2})V_{\psi}(\Delta_{\psi \underset{\nu^o}{a} \otimes_{\beta} \delta^{1/2} P^{1/2})(J_{\psi} \Lambda_{\psi}(x) \underset{\nu^o}{a} \otimes_{\beta} P^{-1/2} \eta)$$

which is :

$$(F_{\psi} \Lambda_{\psi}(y) \underset{\nu}{b} \otimes_{\alpha} \delta^{-1/2} \xi | V_{\psi}(\Lambda_{\psi}(x^*) \underset{\nu^o}{a} \otimes_{\beta} \delta^{1/2} \eta)) = (F_{\psi} \Lambda_{\psi}(y) | (id * \omega_{\delta^{1/2} \eta, \delta^{-1/2} \xi})(V_{\psi}) \Lambda_{\psi}(x))$$

and, using twice 8.4(iii), is equal to :

$$\begin{aligned} & (F_{\psi} \Lambda_{\psi}(y) | \Lambda_{\psi}((id \underset{N}{b} *_{\alpha} \omega_{\eta, \delta^{-1/2} \xi}) \mathbf{a}(x^*)) \\ &= (\Lambda_{\psi}((id \underset{N}{b} *_{\alpha} \omega_{\delta^{-1/2} \xi, \eta}) \mathbf{a}(x) | \Lambda_{\psi}(y)) \\ &= ((id * \omega_{\xi, \eta})(V_{\psi}) \Lambda_{\psi}(x) | \Lambda_{\psi}(y)) \end{aligned}$$

from which we get (iv). □

8.9. Corollary. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A and let ψ be a δ -invariant normal faithful semi-finite weight on A . Let us define the representation a of N on H_{ψ} such that, for $n \in N$:*

$$a(n) = J_{\psi} b(n^*) J_{\psi}$$

and let us suppose that ψ bears the density property, as defined in 8.1. Let V_{ψ} be the corepresentation of \mathfrak{G} on the bimodule ${}_a(H_{\psi})_b$ constructed in 8.6. Then :

(i) it is possible to define the one parameter group of unitaries $\Delta_{\psi \underset{N^o}{a} \otimes_{\beta} \delta^{-it} \Delta_{\widehat{\Phi}}^{-it}}$ and another one $\Delta_{\psi \underset{N}{b} \otimes_{\alpha} \delta^{-it} \Delta_{\widehat{\Phi}}^{-it}}$, with natural values on elementary tensor products.

(ii) we have :

$$V_{\psi}(\Delta_{\psi \underset{N^o}{a} \otimes_{\beta} \delta^{-it} \Delta_{\widehat{\Phi}}^{-it}) = (\Delta_{\psi \underset{N}{b} \otimes_{\alpha} \delta^{-it} \Delta_{\widehat{\Phi}}^{-it})V_{\psi}$$

(iii) for any x in A , t in \mathbb{R} , we have :

$$\mathbf{a}(\sigma_t^{\psi}(x)) = (\Delta_{\psi \underset{N}{b} \otimes_{\alpha} \delta^{-it} \Delta_{\widehat{\Phi}}^{-it}) \mathbf{a}(x) (\Delta_{\psi \underset{N}{b} \otimes_{\alpha} \delta^{it} \Delta_{\widehat{\Phi}}^{it})$$

Proof. Using 3.10(vii), $\Delta_{\widehat{\Phi}}$ is the closure of $PJ_{\Phi}\delta^{-1}J_{\Phi}$, and, therefore, $\overline{\delta\Delta_{\widehat{\Phi}}}$ is equal to $\overline{P\delta J_{\Phi}\delta^{-1}J_{\Phi}}$, and, using 8.8(i), we can define $\Delta_{\psi}^{it} {}_a\otimes_{\beta} \delta^{-it}\Delta_{\widehat{\Phi}}^{-it}$ and $\Delta_{\psi}^{it} {}_b\otimes_{\alpha} \delta^{-it}\Delta_{\widehat{\Phi}}^{-it}$ by writing :

$$\begin{aligned}\Delta_{\psi}^{it} {}_a\otimes_{\beta} \delta^{-it}\Delta_{\widehat{\Phi}}^{-it} &= (\Delta_{\psi}^{it} {}_a\otimes_{\beta} \delta^{-it}P^{-it})(1_{{}_a\otimes_{\beta} J_{\Phi}\delta^{-it}J_{\Phi}}) \\ \Delta_{\psi}^{it} {}_b\otimes_{\alpha} \delta^{-it}\Delta_{\widehat{\Phi}}^{-it} &= (\Delta_{\psi}^{it} {}_b\otimes_{\alpha} \delta^{-it}P^{-it})(1_{{}_b\otimes_{\alpha} J_{\Phi}\delta^{-it}J_{\Phi}})\end{aligned}$$

We then obtain (ii), thanks to 8.8(i) and 8.6(ii). Then (iii) is an easy corollary of (ii) and 8.6(i). \square

8.10. Corollary. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A ; let ψ a δ -invariant normal semi-finite faithful weight on A , and let us define the representation a of N on H_{ψ} such that, for $n \in N$:*

$$a(n) = J_{\psi}b(n^*)J_{\psi}$$

and let us suppose that ψ bears the density property, as defined in 8.1. Then :

- (i) \mathfrak{a} has a standard implementation;
- (ii) \mathfrak{a} satisfies property (A) if and only if \mathfrak{a} is saturated.

Proof. We obtain (i) by 8.8. We obtain then (ii) by using (i) and 7.7. \square

8.11. Corollary. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A and let ψ_1, ψ_2 be two δ -invariant normal faithful semi-finite weights on A , and let us suppose that both ψ_1 and ψ_2 bear the density property. Then, for all t in \mathbb{R} , $(D\psi_1 : D\psi_2)_t$ belongs to $A^{\mathfrak{a}}$.*

Proof. Let us consider the von Neumann algebra $B = A \otimes M_2(\mathbb{C})$; for n in N , let us define $\tilde{b}(n) = b(n) \otimes (e_{11} + e_{22})$, which is an injective $*$ -antihomomorphism from N into B , and, for $X = (x_{i,j})_{i,j}$ in B ; let σ denotes the flip from $H_{\Phi} \otimes \mathbb{C}^2$ onto $\mathbb{C}^2 \otimes H_{\Phi}$, and let us define :

$$\mathfrak{b}(X) = (1_{{}_b\otimes_{\alpha} \sigma})(\mathfrak{a}(x_{i,j}))_{i,j}(1_{{}_b\otimes_{\alpha} \sigma^*})$$

which belongs to $B_{\tilde{b}^* \otimes \alpha} M$.

It is easy to check that $(\tilde{b}, \mathfrak{b})$ is an action of \mathfrak{G} on B . Moreover, let ψ be the normal semi-finite faithful weight defined on B by, for $X = (x_{i,j})_{i,j}$ in B^+ , $\psi(X) = \psi_1(x_{11}) + \psi_2(x_{22})$; we then can verify that ψ is a δ -invariant weight with respect to \mathfrak{b} , and that ψ bears the density property.

Therefore, we have, by definition of $(D\psi_1 : D\psi_2)_t$:

$$\begin{aligned}(1_{{}_b\otimes_{\alpha} \sigma})[\mathfrak{a}(D\psi_1 : D\psi_2)_t \otimes e_{21}](1_{{}_b\otimes_{\alpha} \sigma^*}) &= \mathfrak{b}[(D\psi_1 : D\psi_2)_t \otimes e_{21}] \\ &= \mathfrak{b}\sigma_t^{\psi}(1 \otimes e_{21})\end{aligned}$$

which, using 8.8(ii), is equal to :

$$\begin{aligned}(\sigma_t^{\psi} {}_{\tilde{b}^* \otimes \alpha} \tau_{-t} \sigma_{-t}^{\Phi \circ R} \sigma_t^{\Phi}) \mathfrak{b}(1 \otimes e_{21}) &= (\sigma_t^{\psi} {}_{\tilde{b}^* \otimes \alpha} \tau_{-t} \sigma_{-t}^{\Phi \circ R} \sigma_t^{\Phi})[(1 \otimes e_{21}) {}_{\tilde{b} \otimes \alpha} 1] \\ &= [(D\psi_1 : D\psi_2)_t \otimes e_{21}] {}_{\tilde{b} \otimes \alpha} 1\end{aligned}$$

from which we get the result. \square

In that chapter, we define the crossed product of a von Neumann algebra by a measured quantum groupoid (9.1), and the dual action on this new von Neumann algebra (9.4). We prove that the dual action is integrable (9.8), and satisfies (9.7) the property (A) introduced in chapter 6.

9.1. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A ; we define the crossed-product of A by \mathfrak{G} (via the action \mathfrak{a}) as the von Neumann algebra generated by $\mathfrak{a}(A)$ and $1_{b \otimes_{\alpha} \widehat{M}'}_N$ on the Hilbert space $\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}}$. This von Neumann algebra will be denoted by $A \rtimes_{\mathfrak{a}} \mathfrak{G}$. Clearly, this von Neumann algebra is included in $A_{b \otimes_{\alpha} \mathcal{L}(H_{\Phi})}_N$.

9.2. Example. Let \mathcal{G} be a measured groupoid, and \mathfrak{a} an action of \mathcal{G} on a von Neumann algebra A , in the sense of [Y1], [Y2], [Y3]. We have seen in 6.3 that \mathfrak{a} can be as well considered as an action of the abelian measured quantum groupoid constructed from \mathcal{G} (3.13) on A . In [Y1] and [Y3] is given a construction of the crossed product of A by \mathcal{G} ; using ([Y1], 2.14), we get that the crossed product defined in 9.1 is the same.

9.3. Example. Let $M_0 \subset M_1$ be a depth 2 inclusion of von Neumann algebras, with a regular operator-valued weight T_1 from M_1 to M_0 , as defined in 2.2. Let \mathfrak{G}_1 be the measured quantum groupoid, whose underlying von Neumann algebra is $M'_0 \cap M_2$ and whose basis is $M'_0 \cap M_1$, constructed (3.15) from this inclusion, and \mathfrak{G}_2 the measured quantum groupoid constructed from the inclusion $M_1 \subset M_2$, which is isomorphic to $\widehat{\mathfrak{G}}_1^o$ (therefore, we get that $\widehat{\mathfrak{G}}_2^c = \widehat{\mathfrak{G}}_1$). Then (6.5), there exists ([EV], 7.3) a canonical action \mathfrak{a}_2 of \mathfrak{G}_2 on M_1 , which has the following properties :

(i) the basis of \mathfrak{G}_2 is $M'_1 \cap M_2$ (which, using j_1 , is anti-isomorphic to $M'_0 \cap M_1$), the underlying von Neumann algebra of \mathfrak{G}_2 is $M'_1 \cap M_3$ and, for x in M_1 , $\mathfrak{a}_2(x)$ belongs to $M_1_{j_1 * j_2} M'_1 \cap M_3$; in fact \mathfrak{a}_2 is given by the natural inclusion of M_1 into M_3 ;

(ii) we have $M_0 = M_1^{\mathfrak{a}_2}$;

(iii) there is an isomorphism $I_{\mathfrak{a}_2}$ from $M_1 \rtimes_{\mathfrak{a}_2} \mathfrak{G}_2$ onto M_2 ([EV], 7.6), such that, for any $x \in M_1$, we have $I_{\mathfrak{a}_2}(\mathfrak{a}_2(x)) = x$, and, for any $y \in M'_0 \cap M_2$, we have $I_{\mathfrak{a}_2}(1_{j_1 \otimes_{j_2} y}_{M'_1 \cap M_2}) = y$.

9.4. Theorem. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A , $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ the crossed product of A by \mathfrak{G} via the action \mathfrak{a} . Then :

(i) the operator $1_{\mathfrak{H}_{b \otimes_{\alpha} W^{\text{co}}}}_N$ is a corepresentation of $(\widehat{\mathfrak{G}})^c$ on the $N^o - N^o$ bimodule $1_{b \otimes_{\alpha} \widehat{\beta}}(\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}})_{1_{b \otimes_{\alpha} \widehat{\alpha}}}$.

(ii) this corepresentation implements an action $(1_{b \otimes_{\alpha} \widehat{\alpha}}, \tilde{\mathfrak{a}})$ of $(\widehat{\mathfrak{G}})^c$ on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, which verifies, for any $x \in A$, $y \in \widehat{M}'$:

$$\tilde{\mathfrak{a}}(\mathfrak{a}(x)) = \mathfrak{a}(x)_{\widehat{\alpha} \otimes_{\beta} 1_{N^o}}$$

$$\tilde{\mathfrak{a}}(1_{b \otimes_{\alpha} y}_N) = 1_{b \otimes_{\alpha} \widehat{\Gamma}^c(y)}_N$$

This action $(1_{b \otimes_{\alpha} \hat{\alpha}}, \tilde{\mathfrak{a}})$ (or $\tilde{\mathfrak{a}}$ for simplification) of $(\hat{\mathfrak{G}})^c$ will be called the dual action of \mathfrak{a} .

(iii) We have :

$$\mathfrak{a}(A) \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}} = A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap A_{b \otimes_{\alpha} M}$$

Proof. (i) is given by 5.6. Moreover, as $\mathfrak{a}(x)$ belongs to $A_{b \otimes_{\alpha} M}$, we have :

$$(1_{\mathfrak{H}} \otimes_{\alpha} W^{co})(\mathfrak{a}(x) \otimes_{\beta} 1)(1_{\mathfrak{H}} \otimes_{\alpha} W^{co})^* = \mathfrak{a}(x) \otimes_{\beta} 1$$

and, using 3.12(vi), we get :

$$(1_{\mathfrak{H}} \otimes_{\alpha} W^{co})(1 \otimes_{\alpha} y \otimes_{\beta} 1)(1_{\mathfrak{H}} \otimes_{\alpha} W^{co})^* = 1 \otimes_{\alpha} \hat{\Gamma}^c(y)$$

and, therefore, we have :

$$\begin{aligned} (1_{\mathfrak{H}} \otimes_{\alpha} W^{co})A \rtimes_{\mathfrak{a}} \mathfrak{G}(1_{\mathfrak{H}} \otimes_{\alpha} W^{co})^* \\ \subset (\mathfrak{a}(A) \otimes_{\beta} 1 \cup 1 \otimes_{\alpha} \widehat{M'} \otimes_{\beta} \widehat{M'})'' \subset A \rtimes_{\mathfrak{a}} \mathfrak{G} \otimes_{\beta} \widehat{M'} \end{aligned}$$

which proves (ii).

The inclusion $\mathfrak{a}(A) \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$ is given by (ii); let now X be in $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, then X belongs to $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$ if and only if we have :

$$(1 \otimes_{\alpha} W^{co})(X \otimes_{\beta} 1) = (X \otimes_{\beta} 1)(1 \otimes_{\alpha} W^{co})$$

which means that X belongs to $\mathcal{L}(\mathfrak{H})_{b \otimes_{\alpha} M}$. As $A \rtimes_{\mathfrak{a}} \mathfrak{G} \subset A_{b \otimes_{\alpha} \mathcal{L}(H_{\Phi})}$, this finishes the proof of (iii). \square

9.5. Example. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, and (id, β) its trivial action on N^o (6.2); then the crossed-product $N^o \rtimes_{\beta} \mathfrak{G}$ is equal to $\widehat{M'}$; moreover, the dual action $\tilde{\beta}$ is equal to $\hat{\Gamma}^c$, considered as an action of $(\hat{\mathfrak{G}})^c$ on $\widehat{M'}$ (6.10). Therefore, (β, Γ) , considered as an action of \mathfrak{G} on M , is the dual action of the trivial action (id, α) of $\hat{\mathfrak{G}}^o$ on N .

9.6. Example. Let \mathcal{G} be a measured groupoid, and \mathfrak{a} an action of \mathcal{G} on a von neumann algebra A , in the sense of [Y1], [Y2], [Y3]. We have seen in 9.2 that the crossed-product $A \rtimes_{\mathfrak{a}} \mathcal{G}$ in the sense of 9.1 is equal to the crossed-product in the sense of Yamanouchi; moreover, the dual action $\tilde{\mathfrak{a}}$ in the sense of 9.4 is an action of the measured quantum groupoid $\hat{\mathcal{G}}^c$ (whose underlying von Neumann algebra is the von Neumann algebra generated by the right convolution algebra of \mathcal{G}). Using ([Y3], 4.4), we see it is equal to the "dual coaction of \mathcal{G} " introduced in ([Y3], 4.9).

9.7. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A ; the dual action $\tilde{\mathfrak{a}}$ of $(\hat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ satisfies the property (A) defined in 7.1.

Proof. Let us consider the von Neumann algebra B generated on the Hilbert space $\mathfrak{H}_{b \otimes_\alpha}$ $H_\Phi \hat{\otimes}_{\beta} H_\Phi$ by $\tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G})$ and $1_{b \otimes_\alpha} 1_{\hat{\alpha} \otimes_\beta} \beta(N)'$. By 9.4(ii), it is generated by $\mathfrak{a}(A)_{\hat{\alpha} \otimes_\beta} 1_{N^o}$, $1_{b \otimes_\alpha} \hat{\Gamma}'(\widehat{M}')$ and $1_{b \otimes_\alpha} 1_{\hat{\alpha} \otimes_\beta} \beta(N)'$; but, using 7.2 applied to $(\widehat{\mathfrak{G}})^c$, we get that :

$$(\hat{\Gamma}'(\widehat{M}') \cup 1_{\hat{\alpha} \otimes_\beta} \beta(N)')'' = \widehat{M}'_{\hat{\alpha} \otimes_\beta} \mathcal{L}(H_\Phi)$$

and, therefore :

$$B = (\mathfrak{a}(A)_{\hat{\alpha} \otimes_\beta} 1 \cup 1_{b \otimes_\alpha} \widehat{M}'_{\hat{\alpha} \otimes_\beta} \mathcal{L}(H_\Phi))''$$

or :

$$\begin{aligned} B' &= \mathfrak{a}(A)'_{\hat{\alpha} \otimes_\beta} \mathcal{L}(H_\Phi) \cap \mathcal{L}(\mathfrak{H})_{b \otimes_\alpha} \widehat{M}_{\hat{\alpha} \otimes_\beta} 1 \\ &= (\mathcal{L}(\mathfrak{H})_{b \otimes_\alpha} \widehat{M} \cap \mathfrak{a}(A)')_{\hat{\alpha} \otimes_\beta} 1 \end{aligned}$$

which gives that $B = A \rtimes_{\mathfrak{a}} \mathfrak{G}_{\hat{\alpha} \otimes_\beta} \mathcal{L}(H_\Phi)$, and finishes the proof. \square

9.8. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A ; the dual action $\tilde{\mathfrak{a}}$ of $(\widehat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is integrable.*

Proof. Let y be positive in \widehat{M}' ; we have :

$$T_{\tilde{\mathfrak{a}}}(1_{b \otimes_\alpha} y) = (id_{\hat{\alpha} \otimes_\beta} \widehat{\Phi}^c)(1_{b \otimes_\alpha} \widehat{\Gamma}^c(y)) = 1_{b \otimes_\alpha} \widehat{T}^c(y)$$

As \widehat{T}^c is a semi-finite weight from \widehat{M}' to $\beta(N)$, we get that $T_{\tilde{\mathfrak{a}}}$ is semi-finite, which finishes the proof. \square

9.9. Definition. Let us take the notations of 9.8. Let ψ be a normal semi-finite faithful weight on $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$; then, we shall denote $\tilde{\psi}$ the lifted weight $\tilde{\psi} = \psi \circ T_{\tilde{\mathfrak{a}}}$, which is a normal semi-finite faithful weight on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$.

9.10. Lemma. *With the notations of 9.9, we have, for all $x \in (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$ and $a \in \mathfrak{N}_{\widehat{T}^c}$:*

$$\tilde{\psi}(x^*(1_{b \otimes_\alpha} a^* a)x) = \psi(x^*(1_{b \otimes_\alpha} \widehat{T}^c(a^* a))x)$$

Proof. We have, using 9.4(ii) :

$$\tilde{\mathfrak{a}}(1_{b \otimes_\alpha} a^* a) = 1_{b \otimes_\alpha} \widehat{\Gamma}^c(a^* a)$$

from which we get that :

$$T_{\tilde{\mathfrak{a}}}(1_{b \otimes_\alpha} a^* a) = 1_{b \otimes_\alpha} \widehat{T}^c(a^* a)$$

and, therefore :

$$T_{\tilde{\mathfrak{a}}}[x^*(1_{b \otimes_\alpha} a^* a)x] = x^*[1_{b \otimes_\alpha} \widehat{T}^c(a^* a)]x$$

from which we get the result. \square

9.11. **Proposition.** *Let's take the hypothesis and notations of 9.9; there exists an isometry V from $H_\psi \otimes_{\nu} H_\Phi$ into $H_{\tilde{\psi}}$, such that, we have, for all x in \mathfrak{N}_ψ and a in $\mathfrak{N}_{\widehat{T}^c} \cap \mathfrak{N}_{\widehat{\Phi}^c}$:*

$$V(\Lambda_\psi(x) \otimes_{\nu} \Lambda_{\widehat{\Phi}^c}(a)) = \Lambda_{\tilde{\psi}}((1 \otimes_N a)x)$$

Moreover, we have, for all $b \in \widehat{M'}$, $y \in (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$:

$$V(1 \otimes_{N^o} b) = \pi_{\tilde{\psi}}(1 \otimes_N b)V$$

$$V(J_\psi y J_\psi \otimes_{N^o} 1) = J_{\tilde{\psi}} \pi_{\tilde{\psi}}(y) J_{\tilde{\psi}} V$$

Proof. For any a in $\mathfrak{N}_{\widehat{T}^c} \cap \mathfrak{N}_{\widehat{\Phi}^c}$, we have $\Lambda_{\widehat{\Phi}^c}(a) = J_{\widehat{\Phi}} \Lambda_{\widehat{\Phi}}(J_{\widehat{\Phi}} a J_{\widehat{\Phi}})$, which belongs to $D({}_\alpha H_\Phi, \nu)$ (2.2), and $R^{\alpha, \nu}(\Lambda_{\widehat{\Phi}^c}(a)) = J_{\widehat{\Phi}} \Lambda_{\widehat{T}}(J_{\widehat{\Phi}} a J_{\widehat{\Phi}}) J_\nu$, and, therefore, we have:

$$\langle \Lambda_{\widehat{\Phi}^c}(a), \Lambda_{\widehat{\Phi}^c}(a) \rangle_{\alpha, \nu} = \alpha^{-1} [\widehat{T}(J_{\widehat{\Phi}} a^* a J_{\widehat{\Phi}})]^o = \beta^{-1} (\widehat{T}^c(a^* a))$$

and we get :

$$\|\Lambda_\psi(x) \otimes_{\nu} \Lambda_{\widehat{\Phi}^c}(a)\|^2 = (b \circ \beta^{-1}(\widehat{T}^c(a^* a)) \Lambda_\psi(x) | \Lambda_\psi(x))$$

which, using 9.10, is equal to $\|\Lambda_{\tilde{\psi}}((1 \otimes_N a)x)\|^2$. Which, by polarisation, proves the existence of the isometry V .

The formula $V(1 \otimes_{N^o} b) = \pi_{\tilde{\psi}}(1 \otimes_N b)V$ is then trivial.

As, for all $t \in \mathbb{R}$, and $y \in (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$, we have $\sigma_t^{\tilde{\psi}}(y) = \sigma_t^\psi(y)$, we get that if y is analytic with respect to ψ , y is also analytic with respect to $\tilde{\psi}$, and, for such an element :

$$\begin{aligned} V(J_\psi \sigma_{-i/2}^\psi(y^*) J_\psi \otimes_{N^o} 1) (\Lambda_\psi(x) \otimes_{\nu} \Lambda_{\widehat{\Phi}^c}(a)) &= V(\Lambda_\psi(xy) \otimes_{\nu} \Lambda_{\widehat{\Phi}^c}(a)) \\ &= \Lambda_{\tilde{\psi}}((1 \otimes_N a)xy) \end{aligned}$$

which is equal to :

$$J_{\tilde{\psi}} \pi_{\tilde{\psi}}(\sigma_{-i/2}^{\tilde{\psi}}(y^*)) J_{\tilde{\psi}} \Lambda_{\tilde{\psi}}((1 \otimes_N a)x) = J_{\tilde{\psi}} \pi_{\tilde{\psi}}(\sigma_{-i/2}^\psi(y^*)) J_{\tilde{\psi}} \Lambda_{\tilde{\psi}}((1 \otimes_N a)x)$$

which, by continuity, gives the result which finishes the proof. \square

9.12. **Lemma.** *Let's take the hypothesis and notations of 9.9 and 9.11; then, for all X in $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we have $\pi_{\tilde{\psi}}(X)V = VX$.*

Proof. Let $x \in A$; we have :

$$\begin{aligned} (\mathfrak{a} \otimes_{N^o} id) \mathfrak{a}(x) &= (id \otimes_N \Gamma) \mathfrak{a}(x) \\ &= (1 \otimes_N \widehat{W}^o) (\mathfrak{a}(x) \otimes_{N^o} 1) ((1 \otimes_N \widehat{W}^o)^*) \end{aligned}$$

and, therefore :

$$(\mathfrak{a}(x) \otimes_{N^o} 1) ((1 \otimes_N \widehat{W}^o)^*) = (1 \otimes_N \widehat{W}^o)^* (\mathfrak{a} \otimes_N id) \mathfrak{a}(x)$$

Let $\xi \in D({}_\alpha H_\Phi)$, $\eta \in D(H_\beta)$; we have :

$$\mathfrak{a}(x) (1 \otimes_N (id * \omega_{\xi, \eta}(\widehat{W}^o)^*)) = (\rho_\eta^{\beta, \alpha})^* (1 \otimes_N \widehat{W}^o)^* (\mathfrak{a} \otimes_N id) \mathfrak{a}(x) \rho_\xi^{b, \alpha}$$

Let $(e_i)_{i \in I}$ be an orthogonal (α, ν) -basis of H_Φ ; as $\sum_i \rho_{e_i}^{\beta, \alpha} (\rho_{e_i}^{\beta, \alpha})^* = 1$, it is equal to :

$$\begin{aligned} \sum_i (\rho_\eta^{\beta, \alpha})^* (1_{b \otimes_N \alpha} \widehat{W^o})^* \rho_{e_i}^{\beta, \alpha} (\rho_{e_i}^{\beta, \alpha})^* (\mathbf{a}_{\beta^* \alpha} id) \mathbf{a}(x) \rho_\xi^{b, \alpha} = \\ \sum_i (1_{b \otimes_N \alpha} (id * \omega_{e_i, \eta}) (\widehat{W^o}^*)) \mathbf{a}((id_{b^* \alpha} \omega_{\xi, e_i}) \mathbf{a}(x)) \end{aligned}$$

This sum being convergent σ^* -strongly.

Let y be in $\mathfrak{N}_T \cap \mathfrak{N}_\Phi \cap \mathfrak{N}_{RTT} \cap \mathfrak{N}_{\Phi \circ R}$; and let us put $\eta = J_{\widehat{\Phi}} J_\Phi \Lambda_\Phi(y)$; then $J_{\widehat{\Phi}} \eta$ belongs to $J_\Phi \Lambda_\Phi(\mathfrak{N}_\Phi) \cap D(\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$, thanks to 3.10(i); therefore, η belongs to $D(\alpha H_\Phi, \nu) \cap D((H_\Phi)_\beta, \nu^o)$; moreover, as $\pi'(J_{\widehat{\Phi}} \eta)$ is a bounded operator, we get (4.1) that $\omega_{J_{\widehat{\Phi}} \xi, J_{\widehat{\Phi}} \eta}$ belongs to I_Φ , and, therefore, by 4.8, that $(\omega_{J_{\widehat{\Phi}} \xi, J_{\widehat{\Phi}} \eta} * id)(W)$ belongs to $\mathfrak{N}_{\widehat{\Phi}}$; we then get easily that $(id * \omega_{\xi, \eta}) (\widehat{W^o}^*)$ belongs to $\mathfrak{N}_{\widehat{\Phi}^c}$, and that :

$$\begin{aligned} \Lambda_{\widehat{\Phi}^c}((id * \omega_{\xi, \eta}) (\widehat{W^o}^*)) &= J_{\widehat{\Phi}} \Lambda_{\widehat{\Phi}}((\omega_{J_{\widehat{\Phi}} \xi, J_{\widehat{\Phi}} \eta} * id)(W)) \\ &= J_{\widehat{\Phi}} a_\Phi(\omega_{J_{\widehat{\Phi}} \xi, J_{\widehat{\Phi}} \eta}) \\ &= J_{\widehat{\Phi}} \pi'(J_{\widehat{\Phi}} \eta)^* J_{\widehat{\Phi}} \xi \\ &= J_{\widehat{\Phi}} J_\Phi y^* J_\Phi J_{\widehat{\Phi}} \xi \end{aligned}$$

Let x' be in \mathfrak{N}_ψ , then, $\mathbf{a}(x)(1_{b \otimes_N \alpha} (id * \omega_{\xi, \eta}) (\widehat{W^o}^*)) x'$ belongs to $\mathfrak{N}_{\widehat{\psi}}$, by 9.11, and is the σ^* -strong limit of :

$$\sum_i (1_{b \otimes_N \alpha} (id * \omega_{e_i, \eta}) (\widehat{W^o}^*)) \mathbf{a}((id_{b^* \alpha} \omega_{\xi, e_i}) \mathbf{a}(x)) x'$$

Let J finite, $J \subset I$; then :

$$\Lambda_\psi \left[\sum_{i \in J} (1_{b \otimes_N \alpha} (id * \omega_{e_i, \eta}) (\widehat{W^o}^*)) \mathbf{a}((id_{b^* \alpha} \omega_{\xi, e_i}) \mathbf{a}(x)) x' \right]$$

is, using 9.11, equal to :

$$V \left(\sum_{i \in J} [(id_{b^* \alpha} \omega_{\xi, e_i}) \mathbf{a}(x) \Lambda_\psi(x')_{b \otimes_N \alpha} \Lambda_{\widehat{\Phi}^c}((id * \omega_{e_i, \eta}) (\widehat{W^o}^*))] \right)$$

which is :

$$V \left(\sum_{i \in J} [(id_{b^* \alpha} \omega_{\xi, e_i}) \mathbf{a}(x) \Lambda_\psi(x')_{b \otimes_N \alpha} J_{\widehat{\Phi}} J_\Phi y^* J_\Phi J_{\widehat{\Phi}} e_i] \right)$$

which can be written :

$$V(1_{b \otimes_N \alpha} J_{\widehat{\Phi}} J_\Phi y^* J_\Phi J_{\widehat{\Phi}}) \sum_{i \in J} \rho_{e_i}^{\beta, \alpha} (\rho_{e_i}^{\beta, \alpha})^* \mathbf{a}(x) (\Lambda_\psi(x')_{b \otimes_N \alpha} \xi)$$

which converges in norm, when J is growing, to :

$$\begin{aligned} V(1_{b \otimes_N \alpha} J_{\widehat{\Phi}} J_\Phi y^* J_\Phi J_{\widehat{\Phi}}) \mathbf{a}(x) (\Lambda_\psi(x')_{b \otimes_N \alpha} \xi) &= V \mathbf{a}(x) (\Lambda_\psi(x')_{b \otimes_N \alpha} J_{\widehat{\Phi}} J_\Phi y^* J_\Phi J_{\widehat{\Phi}} \xi) \\ &= V \mathbf{a}(x) (\Lambda_\psi(x')_{b \otimes_N \alpha} \Lambda_{\widehat{\Phi}^c}((id * \omega_{\xi, \eta}) (\widehat{W^o}^*)) \end{aligned}$$

As $\Lambda_{\tilde{\psi}}$ is σ^* -norm closed, we therefore get that :

$$\begin{aligned} V\mathbf{a}(x)(\Lambda_{\psi}(x') \underset{N}{b} \otimes_{\alpha} \Lambda_{\widehat{\Phi}^c}((id * \omega_{\xi, \eta})(\widehat{W^o}^*)) &= \Lambda_{\tilde{\psi}}[\mathbf{a}(x)(1 \underset{N}{b} \otimes_{\alpha} (id * \omega_{\xi, \eta})(\widehat{W^o}^*))x'] \\ &= \pi_{\tilde{\psi}}(\mathbf{a}(x))\Lambda_{\tilde{\psi}}[(1 \underset{N}{b} \otimes_{\alpha} (id * \omega_{\xi, \eta})(\widehat{W^o}^*))x'] \\ &= \pi_{\tilde{\psi}}(\mathbf{a}(x))V(\Lambda_{\psi}(x') \underset{N}{b} \otimes_{\alpha} \Lambda_{\widehat{\Phi}^c}((id * \omega_{\xi, \eta})(\widehat{W^o}^*))) \end{aligned}$$

and, by linearity and continuity, we get that $\pi_{\tilde{\psi}}(\mathbf{a}(x))V = V\mathbf{a}(x)$, for any $x \in A$. As we have obtained in 9.11 that $V(1 \underset{N^o}{b} \otimes_{\alpha} b) = \pi_{\tilde{\psi}}(1 \underset{N}{b} \otimes_{\alpha} b)V$ for any $b \in \widehat{M'}$, we obtain that $\pi_{\tilde{\psi}}(X)V = VX$ for any X in the involutive algebra generated by $\mathbf{a}(A)$ and $1 \underset{N}{b} \otimes_{\alpha} \widehat{M'}$; then, by strong limit, we obtain the result. \square

10. AN AUXILLIARY WEIGHT ON THE CROSSED-PRODUCT

Be given an action \mathbf{a} of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A , and a normal semi-finite faithful weight ψ on A , we construct, in this chapter, (10.1) an auxilliary weight $\tilde{\psi}_0$ on $A \rtimes_{\mathbf{a}} \mathfrak{G}$ which will allow us, thanks to chapter 8, to construct a standard implementation for the dual action (10.7). We obtain (10.7) that the dual action satisfies the saturation property introduced in chapter 7; we introduce a third technical property (property (B) in 10.10) which is satisfied by the dual action (10.12).

10.1. Proposition. *Let's take the hypothesis and notations of 9.9 and 9.11; then :*

(i) *there exists a unique normal semi-finite faithful weight $\tilde{\psi}_0$ on $A \rtimes_{\mathbf{a}} \mathfrak{G}$ such that :*

$$\mathfrak{N}_{\tilde{\psi}_0} = \{x \in \mathfrak{N}_{\tilde{\psi}}, \Lambda_{\tilde{\psi}}(x) \in ImV\}$$

and such that we may identify $H_{\tilde{\psi}_0}$ with ImV , $\Lambda_{\tilde{\psi}_0}$ with the restriction of $\Lambda_{\tilde{\psi}}$ to $\mathfrak{N}_{\tilde{\psi}_0}$, and, for any $y \in A \rtimes_{\mathbf{a}} \mathfrak{G}$, $\pi_{\tilde{\psi}_0}(y)$ to the restriction of $\pi_{\tilde{\psi}}(y)$ to ImV .

(ii) *let us write \tilde{V} the unitary from $H_{\psi} \underset{\nu}{b} \otimes_{\alpha} H_{\Phi}$ onto $H_{\tilde{\psi}_0}$ defined, for all $\Xi \in H_{\psi} \underset{\nu}{b} \otimes_{\alpha} H_{\Phi}$, by $\tilde{V}\Xi = V\Xi$; then we have, for all y in $A \rtimes_{\mathbf{a}} \mathfrak{G}$:*

$$\pi_{\tilde{\psi}_0}(y) = \tilde{V}y\tilde{V}^*$$

(iii) *the linear set generated by the elements of the form $(1 \underset{N}{b} \otimes_{\alpha} a)x$, with $a \in \mathfrak{N}_{\widehat{T}^c} \cap \mathfrak{N}_{\widehat{\Phi}^c}$ and $x \in \mathfrak{N}_{\tilde{\psi}}$, is dense in $A \rtimes_{\mathbf{a}} \mathfrak{G}$.*

Proof. Using 9.10, we get that $\{x \in \mathfrak{N}_{\tilde{\psi}}, \Lambda_{\tilde{\psi}}(x) \in ImV\}$ is an ideal of $A \rtimes_{\mathbf{a}} \mathfrak{G}$, which is dense by 9.11; then, we can apply ([V2], 7.4) to obtain (i); result (ii) is then straightforward. Result (iii) is also a corollary of 9.10 and 9.11. \square

10.2. Proposition. *Let's take the hypothesis and notations of 9.9, 9.11 and 10.1; let us define the unitary U_{ψ} from $H_{\psi} \underset{\nu^o}{a} \otimes_{\beta} H_{\Phi}$ onto $H_{\psi} \underset{\nu}{b} \otimes_{\alpha} H_{\Phi}$ by the formula :*

$$U_{\psi} = \tilde{V}^* J_{\tilde{\psi}_0} \tilde{V} (J_{\psi} \underset{\nu^o}{a} \otimes_{\beta} J_{\Phi})$$

Then :

(i) *for all y in $(A \rtimes_{\mathbf{a}} \mathfrak{G})^{\tilde{\mathbf{a}}}$, we have $y = U_{\psi}(\pi_{\psi}(y) \underset{N^o}{a} \otimes_{\beta} 1)U_{\psi}^*$;*

(ii) *for all $t \in \mathbb{R}$, we have $\sigma_t^{\tilde{\psi}_0}(y) = \sigma_t^{\psi}(y)$;*

(iii) *we have $U_{\psi}(J_{\psi} \underset{\nu}{b} \otimes_{\alpha} J_{\Phi}) = (J_{\psi} \underset{\nu^o}{a} \otimes_{\beta} J_{\Phi})U_{\psi}^*$.*

Proof. Let us first remark that it is possible to define on elementary tensors an antilinear surjective isometry $J_{\psi a \otimes_{\beta} J_{\hat{\Phi}}}$ from $H_{\psi a \otimes_{\beta} H_{\Phi}}$ onto $H_{\psi b \otimes_{\alpha} H_{\Phi}}$, whose inverse is $J_{\psi b \otimes_{\alpha} J_{\hat{\Phi}}}$, which is defined the same way.

Let us take now $y \in D(\sigma_{i/2}^{\psi})$; we have, for all $x \in \mathfrak{N}_{\psi}$, $a \in \mathfrak{N}_{\hat{\Phi}^c} \cap \mathfrak{N}_{\hat{T}^c}$:

$$\begin{aligned} \Lambda_{\tilde{\psi}_0}[(1 \otimes_N a)xy] &= \tilde{V}(\Lambda_{\psi}(xy) \otimes_{\alpha} \Lambda_{\hat{\Phi}^c}(a)) \\ &= \tilde{V}(J_{\psi} \pi_{\psi}(\sigma_{i/2}^{\psi}(y))^* J_{\psi} \otimes_{\alpha} 1)(\Lambda_{\psi}(x) \otimes_{\alpha} \Lambda_{\hat{\Phi}^c}(a)) \end{aligned}$$

which is equal to :

$$\tilde{V}(J_{\psi} \pi_{\psi}(\sigma_{i/2}^{\psi}(y))^* J_{\psi} \otimes_{\alpha} 1) \tilde{V}^* \Lambda_{\tilde{\psi}_0}[(1 \otimes_N a)x]$$

As the linear set generated by all the elements of the form $(1 \otimes_N a)x$ is, by construction of $\tilde{\psi}_0$, a core for $\Lambda_{\tilde{\psi}_0}$, we get that, for any $X \in \mathfrak{N}_{\tilde{\psi}_0}$, we have :

$$\Lambda_{\tilde{\psi}_0}(Xy) = \tilde{V}(J_{\psi} \pi_{\psi}(\sigma_{i/2}^{\psi}(y))^* J_{\psi} \otimes_{\alpha} 1) \tilde{V}^* \Lambda_{\tilde{\psi}_0}(X)$$

from which we deduce that y belongs to $D(\sigma_{i/2}^{\tilde{\psi}_0})$, and that :

$$J_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0}(\sigma_{i/2}^{\tilde{\psi}_0}(y))^* J_{\tilde{\psi}_0} = \tilde{V}(J_{\psi} \pi_{\psi}(\sigma_{i/2}^{\psi}(y))^* J_{\psi} \otimes_{\alpha} 1) \tilde{V}^*$$

which can be written also, thanks to 10.1(ii) :

$$\sigma_{i/2}^{\tilde{\psi}_0}(y) = U_{\psi}(\pi_{\psi}(\sigma_{i/2}^{\psi}(y)) \otimes_{\beta} 1) U_{\psi}^*$$

Taking the adjoints, we get that, if y' belongs to $D(\sigma_{-i/2}^{\psi})$, then y' belongs to $D(\sigma_{-i/2}^{\tilde{\psi}_0})$, and :

$$\sigma_{-i/2}^{\tilde{\psi}_0}(y') = U_{\psi}(\pi_{\psi}(\sigma_{-i/2}^{\psi}(y')) \otimes_{\beta} 1) U_{\psi}^*$$

If now z belongs to $D(\sigma_{-i}^{\psi})$, then z belongs to $D(\sigma_{-i/2}^{\tilde{\psi}_0})$, and :

$$\sigma_{-i/2}^{\tilde{\psi}_0}(z) = U_{\psi}(\pi_{\psi}(\sigma_{-i/2}^{\psi}(z)) \otimes_{\beta} 1) U_{\psi}^*$$

But, as $\sigma_{-i}^{\psi}(z)$ belongs to $D(\sigma_{i/2}^{\psi})$, $\sigma_{-i}^{\psi}(z)$ belongs to $D(\sigma_{i/2}^{\tilde{\psi}_0})$ and :

$$\sigma_{i/2}^{\tilde{\psi}_0}(\sigma_{-i}^{\psi}(z)) = U_{\psi}(\pi_{\psi}(\sigma_{i/2}^{\psi}(\sigma_{-i}^{\psi}(z))) \otimes_{\beta} 1) U_{\psi}^* = U_{\psi}(\pi_{\psi}(\sigma_{-i/2}^{\psi}(z)) \otimes_{\beta} 1) U_{\psi}^*$$

and we get that $\sigma_{-i/2}^{\tilde{\psi}_0}(z) = \sigma_{i/2}^{\tilde{\psi}_0}(\sigma_{-i}^{\psi}(z))$; therefore, z belongs to $D(\sigma_{-i}^{\tilde{\psi}_0})$, and $\sigma_{-i}^{\tilde{\psi}_0}(z) = \sigma_{-i}^{\psi}(z)$. Then, using ([H], 4.3 and 4.4), we obtain (ii). Let us return now to the formula obtained for $y \in D(\sigma_{i/2}^{\psi})$; we had obtained :

$$\sigma_{i/2}^{\tilde{\psi}_0}(y) = U_{\psi}(\pi_{\psi}(\sigma_{i/2}^{\psi}(y)) \otimes_{\beta} 1) U_{\psi}^*$$

Using (ii), we get :

$$\sigma_{i/2}^{\psi}(y) = U_{\psi}(\pi_{\psi}(\sigma_{i/2}^{\psi}(y)) \otimes_{\beta} 1) U_{\psi}^*$$

from which, by density, one obtains (i).

From the definition of U_ψ , one gets :

$$U_\psi^* = (J_\psi \underset{\nu}{b} \otimes_\alpha J_{\widehat{\Phi}}) \tilde{V}^* J_{\tilde{\psi}_0} \tilde{V}$$

and the formula (iii) is straightforward. \square

10.3. Proposition. *Let's take the hypothesis and notations of 9.9, 9.11 and 10.1; then, the weight $\tilde{\psi}_0$ is $\hat{\delta}^{-1}$ -invariant with respect to the dual action $\tilde{\mathbf{a}}$.*

Proof. By 8.2 applied to $\widehat{\mathfrak{G}}^c$, we know that $\widehat{\Phi}^c$ is $\hat{\delta}^{-1}$ -invariant with respect to the action $\widehat{\Gamma}^c$ of $\widehat{\mathfrak{G}}^c$ on \widehat{M}' . Moreover, by 4.5 applied to $\widehat{\Gamma}^c$, we get that :

$$\Lambda_{\widehat{\Phi}^c} [((id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi}) \widehat{\Gamma}^c(a))] = (id * \omega_{\hat{\delta}^{-1/2} \eta, \xi})(W^{oc}) \Lambda_{\widehat{\Phi}^c}(a)$$

for all $a \in \mathfrak{N}_{\widehat{\Phi}^c}$, $\xi \in D((H_\Phi)_\beta, \nu^o)$ and $\eta \in D((H_\Phi)_\beta, \nu^o)$, such that $\hat{\delta}^{-1/2} \eta$ belongs to $D(\hat{\alpha} H_\Phi, \nu)$.

Therefore, for any $x \in \mathfrak{N}_\psi$ and $a \in \mathfrak{N}_{\widehat{T}^c} \cap \mathfrak{N}_{\widehat{\Phi}^c}$, we have :

$$\Lambda_{\tilde{\psi}_0} [(id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi}) \tilde{\mathbf{a}}((1 \underset{N}{b} \otimes_\alpha a)x)] = \Lambda_{\tilde{\psi}_0} [(1 \underset{N}{b} \otimes_\alpha (id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi}) \widehat{\Gamma}^c(a)x)]$$

which is equal to :

$$\begin{aligned} & \tilde{V}[\Lambda_\psi(x) \underset{\nu}{b} \otimes_\alpha \Lambda_{\widehat{\Phi}^c}[(id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi}) \widehat{\Gamma}^c(a)]] \\ &= \tilde{V}[1 \underset{N}{b} \otimes_\alpha (id * \omega_{\hat{\delta}^{-1/2} \eta, \xi})(W^{oc})](\Lambda_\psi(x) \underset{\nu}{b} \otimes_\alpha \Lambda_{\widehat{\Phi}^c}(a)) \\ &= \tilde{V}[1 \underset{N}{b} \otimes_\alpha (id * \omega_{\hat{\delta}^{-1/2} \eta, \xi})(W^{oc})] \tilde{V}^* \Lambda_{\tilde{\psi}_0}((1 \underset{N}{b} \otimes_\alpha a) \mathbf{a}(x)) \end{aligned}$$

which, by definition of $\tilde{\psi}_0$, and the closedness of $\Lambda_{\tilde{\psi}_0}$, allows us to write, for all $X \in \mathfrak{N}_{\tilde{\psi}_0}$:

$$\Lambda_{\tilde{\psi}_0}((id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi}) \tilde{\mathbf{a}}(X)) = \tilde{V}[1 \underset{N}{b} \otimes_\alpha (id * \omega_{\hat{\delta}^{-1/2} \eta, \xi})(W^{oc})] \tilde{V}^* \Lambda_{\tilde{\psi}_0}(X)$$

Taking now $(\xi_i)_{i \in I}$ an orthogonal (β, ν^o) -basis of H_Φ , we have :

$$\tilde{\psi}_0((id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_\eta) \tilde{\mathbf{a}}(X^* X)) = \sum_i \|\Lambda_{\tilde{\psi}_0}((id \underset{N^o}{\hat{\alpha}} *_{\beta} \omega_{\eta, \xi_i}) \tilde{\mathbf{a}}(X))\|^2$$

which is equal to :

$$\begin{aligned} & \sum_i \|[1 \underset{N}{b} \otimes_\alpha (id * \omega_{\hat{\delta}^{-1/2} \eta, \xi_i})(W^{oc})] \tilde{V}^* \Lambda_{\tilde{\psi}_0}(X)\|^2 = \\ &= \|(1 \underset{N}{b} \otimes_\alpha W^{oc}) \tilde{V}^*(\Lambda_{\tilde{\psi}_0}(X) \underset{\beta_0}{\otimes} \underset{\nu}{\hat{\alpha}} \hat{\delta}^{-1/2} \eta)\|^2 = \|\Lambda_{\tilde{\psi}_0}(X) \underset{\beta_0}{\otimes} \underset{\nu}{\hat{\alpha}} \hat{\delta}^{-1/2} \eta\|^2 \end{aligned}$$

where we put, for $n \in N$, $\beta_0(n) = J_{\tilde{\psi}_0}(1 \underset{N}{b} \otimes_\alpha \hat{\alpha}(n^*)) J_{\tilde{\psi}_0}$. Which is the result. \square

10.4. Lemma. *Let (b, \mathbf{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A , and let ψ be a normal semi-finite faithful weight on A . Let $A \rtimes_{\mathbf{a}} \mathfrak{G}$ be the crossed product, and $\tilde{\psi}_0$ be the weight defined in 10.1. Then, the linear span generated by the elements $\pi_{\tilde{\psi}_0}(x^*) \tilde{V}(\Lambda_\psi(y) \underset{\nu}{b} \otimes_\alpha \Lambda_{\widehat{\Phi}^c}(a))$, where x, y belong to \mathfrak{N}_ψ and a belongs to $\mathfrak{N}_{\widehat{T}^c} \cap \mathfrak{N}_{\widehat{\Phi}^c} \cap \mathfrak{N}_{\widehat{T}^c}^* \cap \mathfrak{N}_{\widehat{\Phi}^c}^*$, is a core for $S_{\tilde{\psi}_0}$, and we have :*

$$S_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0}(x^*) \tilde{V}(\Lambda_\psi(y) \underset{\nu}{b} \otimes_\alpha \Lambda_{\widehat{\Phi}^c}(a)) = \pi_{\tilde{\psi}_0}(y^*) \tilde{V}(\Lambda_\psi(x) \underset{\nu}{b} \otimes_\alpha \Lambda_{\widehat{\Phi}^c}(a^*))$$

Proof. By definition of $\tilde{\psi}_0$ (10.3(ii)), the linear span generated by the elements $(1_{b \otimes_N \alpha} a)x$, where a belongs to $\mathfrak{N}_{\hat{\mathcal{T}}^c} \cap \mathfrak{N}_{\hat{\mathcal{F}}^c}$, and x belongs to \mathfrak{N}_ψ is a core for $\Lambda_{\tilde{\psi}_0}$. We deduce easily the result from this. \square

10.5. Proposition. *Let (b, \mathfrak{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A , and let ψ be a normal semi-finite faithful weight on A . Let $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ be the crossed product, and $\tilde{\psi}_0$ be the weight defined in 10.1. Then, the unitary U_ψ defined in 10.2 satisfies, for all $b \in M$:*

$$(1_{b \otimes_N \alpha} J_\Phi b J_\Phi) U_\psi = U_\psi (1_{a \otimes_\beta \beta} J_\Phi b J_\Phi)$$

Proof. Using the notations of 4.7 and 10.4, we have :

$$\begin{aligned} (1_{b \otimes_N \alpha} J_\Phi b J_\Phi) \tilde{V}^* S_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0}(x^*) \tilde{V}(\Lambda_\psi(y)_{b \otimes_\alpha} \Lambda_{\hat{\mathcal{F}}^c}(a^*)) = \\ (1_{b \otimes_N \alpha} J_\Phi b J_\Phi) \tilde{V}^* \pi_{\tilde{\psi}_0}(y^*) \tilde{V}(\Lambda_\psi(x)_{b \otimes_\alpha} \Lambda_{\hat{\mathcal{F}}^c}(a)) \end{aligned}$$

which is equal to :

$$\begin{aligned} (1_{b \otimes_N \alpha} J_\Phi b J_\Phi) y^* (\Lambda_\psi(x)_{b \otimes_\alpha} \Lambda_{\hat{\mathcal{F}}^c}(a)) &= y^* (1_{b \otimes_N \alpha} J_\Phi b J_\Phi) (\Lambda_\psi(x)_{b \otimes_\alpha} \Lambda_{\hat{\mathcal{F}}^c}(a)) \\ &= y^* (\Lambda_\psi(x)_{b \otimes_\alpha} J_\Phi b J_\Phi \Lambda_{\hat{\mathcal{F}}^c}(a)) \end{aligned}$$

Using 3.10(v), we get that, for all $t \in \mathbb{R}$, $\Delta_{\hat{\mathcal{F}}}^{it} J_\Phi b J_\Phi \Delta_{\hat{\mathcal{F}}}^{-it}$ belongs to M' , and we define this way an automorphism group μ_t of M' ; if we suppose that $J_\Phi R(b^*) J_\Phi$ belongs to $D(\mu_{-i/2})$, we get that :

$$J_\Phi b J_\Phi \Lambda_{\hat{\mathcal{F}}^c}(a) = J_\Phi \Delta_{\hat{\mathcal{F}}}^{-1/2} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi) \Lambda_{\hat{\mathcal{F}}^c}(a^*)$$

and, therefore, using 10.4, we get :

$$\begin{aligned} (1_{b \otimes_N \alpha} J_\Phi b J_\Phi) \tilde{V}^* S_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0}(x^*) \tilde{V}((\Lambda_\psi(y)_{b \otimes_\alpha} \Lambda_{\hat{\mathcal{F}}^c}(a^*))) \\ = y^* (\Lambda_\psi(x)_{b \otimes_\alpha} J_\Phi \Delta_{\hat{\mathcal{F}}}^{-1/2} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi) \Lambda_{\hat{\mathcal{F}}^c}(a^*)) \\ = \tilde{V}^* \pi_{\tilde{\psi}_0}(y^*) \tilde{V}(\Lambda_\psi(x)_{b \otimes_\alpha} J_\Phi \Delta_{\hat{\mathcal{F}}}^{-1/2} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi) \Lambda_{\hat{\mathcal{F}}^c}(a^*)) \\ = \tilde{V}^* S_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0}(x^*) \tilde{V}(\Lambda_\psi(y)_{b \otimes_\alpha} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi) \Lambda_{\hat{\mathcal{F}}^c}(a^*)) \\ = \tilde{V}^* S_{\tilde{\psi}_0} \tilde{V} x^* (\Lambda_\psi(y)_{b \otimes_\alpha} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi) \Lambda_{\hat{\mathcal{F}}^c}(a^*)) \end{aligned}$$

and, by the closedness of $S_{\tilde{\psi}_0}$, we get :

$$(1_{b \otimes_N \alpha} J_\Phi b J_\Phi) \tilde{V}^* S_{\tilde{\psi}_0} \tilde{V} \subset \tilde{V}^* S_{\tilde{\psi}_0} \tilde{V} (1_{b \otimes_N \alpha} \mu_{-i/2}(J_\Phi R(b^*) J_\Phi))$$

Taking the adjoints, we have $J_\Phi R(b) J_\Phi \in D(\mu_{i/2})$ and :

$$(1_{b \otimes_N \alpha} \mu_{i/2}(J_\Phi R(b) J_\Phi)) \tilde{V}^* F_{\tilde{\psi}_0} \tilde{V} \subset \tilde{V}^* F_{\tilde{\psi}_0} \tilde{V} (1_{b \otimes_N \alpha} J_\Phi b^* J_\Phi)$$

Therefore, if we suppose that $J_\Phi b J_\Phi$ belongs to μ_{-i} , we get :

$$(1_{b \otimes_N \alpha} J_\Phi b J_\Phi) \tilde{V}^* \Delta_{\tilde{\psi}_0} \tilde{V} \subset \tilde{V}^* \Delta_{\tilde{\psi}_0} \tilde{V} ((1_{b \otimes_N \alpha} \mu_{-i}(J_\Phi b J_\Phi))$$

So, for any $X \in \mathcal{L}(H_\psi \underset{\nu}{b} \otimes_\alpha H_\phi)$, let $\epsilon_t(X) = \tilde{V}^* \Delta_{\tilde{\psi}_0}^{it} \tilde{V} X \tilde{V}^* \Delta_{\tilde{\psi}_0}^{-it} \tilde{V}$; we then get that $(1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi)$ belongs to $D(\epsilon_{-i})$, and, therefore, for any b in M , we have :

$$\epsilon_t(1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi) = 1 \underset{N}{b} \otimes_\alpha \mu_t(J_\Phi b J_\Phi)$$

and :

$$(1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi) \tilde{V}^* J_{\tilde{\psi}_0} \tilde{V} = \tilde{V}^* J_{\tilde{\psi}_0} \tilde{V} (1 \underset{N}{b} \otimes_\alpha J_\Phi R(b^*) J_\Phi)$$

and, therefore :

$$(1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi) U_\psi = U_\psi (1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi)$$

□

10.6. Corollary. *Let (b, \mathfrak{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A , and let ψ be a normal semi-finite faithful weight on A . Let $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ be the crossed product, and $\tilde{\psi}_0$ be the weight defined in 10.1. Then, for all $n \in N$, we have :*

$$J_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0} (1 \underset{N}{b} \otimes_\alpha \hat{\alpha}(n))^* J_{\tilde{\psi}_0} = \tilde{V} (1 \underset{N}{b} \otimes_\alpha \hat{\beta}(n)) \tilde{V}^*$$

Proof. We have, using 10.1(ii),

$$J_{\tilde{\psi}_0} \pi_{\tilde{\psi}_0} (1 \underset{N}{b} \otimes_\alpha \hat{\alpha}(n))^* J_{\tilde{\psi}_0} = J_{\tilde{\psi}_0} \tilde{V} (1 \underset{N}{b} \otimes_\alpha \hat{\alpha}(n))^* \tilde{V}^* J_{\tilde{\psi}_0}$$

which, using 10.2, is equal to :

$$\tilde{V} U_\psi (J_\psi \underset{N}{b} \otimes_\alpha J_{\hat{\Phi}}) (1 \underset{N}{b} \otimes_\alpha \hat{\alpha}(n))^* (J_\psi \underset{N^o}{a} \otimes_\beta J_{\hat{\Phi}}) U_\psi^* \tilde{V}^*$$

and, using 10.5, is equal to :

$$\tilde{V} U_\psi (1 \underset{N^o}{a} \otimes_\beta \hat{\beta}(n)) U_\psi^* \tilde{V}^* = \tilde{V} (1 \underset{N}{b} \otimes_\alpha \hat{\beta}(n)) \tilde{V}^*$$

□

10.7. Theorem. *Let (b, \mathfrak{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A ; let $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ be the crossed product, $\tilde{\mathfrak{a}}$ the dual action of $\hat{\mathfrak{G}}^c$ on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, and let ψ be a normal semi-finite faithful weight on $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$. Let $\tilde{\psi}_0$ be the weight defined in 10.1. Then,*

- (i) *the weight $\tilde{\psi}_0$ is $\hat{\delta}^{-1}$ -invariant with respect to the dual action $\tilde{\mathfrak{a}}$, and bears the density property defined in 8.1.*
- (ii) *there exists a standard implementation of the dual action on the Hilbert space $H_{\tilde{\psi}_0}$.*
- (iii) *the dual action is saturated.*

Proof. As $V[D((H_\psi)_b, \nu^o) \underset{\nu}{b} \otimes_\alpha D(\hat{\alpha} H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^o)]$ is dense in $H_{\tilde{\psi}_0}$, we get that $\tilde{\psi}_0$ bears the density property defined in 8.1; so (i) is then given by 10.3. Then (ii) is given by 8.10(i), and (iii) is given by 9.7 and 8.10(ii). □

10.8. Theorem. *Let (b, \mathbf{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A ; let $A \rtimes_{\mathbf{a}} \mathfrak{G}$ be the crossed product, $\tilde{\mathbf{a}}$ the dual action of $\hat{\mathfrak{G}}^c$ on $A \rtimes_{\mathbf{a}} \mathfrak{G}$, and let ψ be a normal semi-finite faithful weight on $(A \rtimes_{\mathbf{a}} \mathfrak{G})^{\tilde{\mathbf{a}}}$. Let $\tilde{\psi}$ be the lifted weight defined in 9.9, and $\tilde{\psi}_0$ be the weight defined in 10.1. Then :*

- (i) *we have $\tilde{\psi} = \tilde{\psi}_0$;*
- (ii) *the linear set generated by the elements of the form $(1 \otimes_N a)x$, with $a \in \mathfrak{N}_{\hat{\mathfrak{G}}^c} \cap \mathfrak{N}_{\hat{T}^c}$ and $x \in \mathfrak{N}_{\psi}$, is a core for $\Lambda_{\tilde{\psi}}$.*
- (iii) *the weight $\tilde{\psi}$ is $\hat{\delta}^{-1}$ -invariant with respect to the dual action $\tilde{\mathbf{a}}$, and bears the density property defined in 8.1.*

Proof. Thanks to 10.7(i), we can apply 8.8(ii) to the action $\tilde{\mathbf{a}}$ and to the weight $\tilde{\psi}_0$, and we have, for any $x \in A \rtimes_{\mathbf{a}} \mathfrak{G}$:

$$\tilde{\mathbf{a}}(\sigma_t^{\tilde{\psi}_0}(x)) = (\sigma_t^{\tilde{\psi}_0} \hat{\alpha}^*_{\beta} \hat{\tau}_{-t}^c \sigma_{-t}^{\hat{\Phi}^c \circ R^c} \sigma_t^{\hat{\Phi}^c}) \tilde{\mathbf{a}}(x)$$

and, using 3.8(vi) and (vii) applied to $\hat{\Phi}^c$, we get that, for any positive x in $A \rtimes_{\mathbf{a}} \mathfrak{G}$, we have $T_{\tilde{\mathbf{a}}}(\sigma_t^{\tilde{\psi}_0}(x)) = \sigma_t^{\tilde{\psi}_0}(T_{\tilde{\mathbf{a}}}(x))$, which implies, using 10.2(ii), that :

$$T_{\tilde{\mathbf{a}}}(\sigma_t^{\tilde{\psi}_0}(x)) = \sigma_t^{\psi}(T_{\tilde{\mathbf{a}}}(x))$$

and, therefore, $\tilde{\psi}(\sigma_t^{\tilde{\psi}_0}(x)) = \tilde{\psi}(x)$.

By construction (10.1(i)), we have $\tilde{\psi}(x) = \tilde{\psi}_0(x)$ for all $x \in \mathfrak{M}_{\tilde{\psi}_0}$; from which we can deduce now (i). Then (ii) is just given by (i) and the definition of $\tilde{\psi}_0$, and (iii) by (i) and 10.7(i). \square

10.9. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a\mathfrak{H}_b$ a $N - N$ bimodule, A a von Neumann algebra such that $b(N) \subset A \subset a(N)'$, V a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$ which implements an action (b, \mathbf{a}) of \mathfrak{G} on A ; let $A \rtimes_{\mathbf{a}} \mathfrak{G}$ be the crossed-product, and $\tilde{\mathbf{a}}$ the dual action of $(\hat{\mathfrak{G}})^c$ on the crossed product; then :*

- (i) $A \rtimes_{\mathbf{a}} \mathfrak{G} \subset V(\mathcal{L}(\mathfrak{H}) \hat{\alpha}^*_{\beta} \widehat{M}') V^*$
- (ii) $(A \rtimes_{\mathbf{a}} \mathfrak{G})^{\tilde{\mathbf{a}}} \subset \text{Sata}$.

Proof. We have obtained in 7.4(ii) that $1 \otimes_N \widehat{M}' \subset V(\mathcal{L}(\mathfrak{H}) \hat{\alpha}^*_{\beta} \widehat{M}') V^*$; as $\mathbf{a}(A) = V(A \hat{\alpha}_{\beta} 1) V^*$, we obtain (i) by definition of $A \rtimes_{\mathbf{a}} \mathfrak{G}$. Then (ii) is a corollary of (i) and 9.4(iii). \square

10.10. Definition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, A a von Neumann algebra, (b, \mathbf{a}) an action of \mathfrak{G} on A ; following ([ES1], II5), we shall say that \mathbf{a} satisfies property (B) if we have $\mathbf{a}(A) = (A \rtimes_{\mathbf{a}} \mathfrak{G})^{\tilde{\mathbf{a}}}$.

10.11. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a\mathfrak{H}_b$ a $N - N$ bimodule, A a von Neumann algebra such that $b(N) \subset A \subset a(N)'$, V a corepresentation of \mathfrak{G} on ${}_a\mathfrak{H}_b$ which implements an action (b, \mathbf{a}) of \mathfrak{G} on A ; let $A \rtimes_{\mathbf{a}} \mathfrak{G}$ be the crossed-product, and $\tilde{\mathbf{a}}$ the dual action of $(\hat{\mathfrak{G}})^c$ on the crossed product; let us suppose that the action \mathbf{a} is saturated, in the sense of 7.3. Then, the action \mathbf{a} satisfies property (B) (in the sense of 10.10).*

Proof. Using 9.4(iii) and 10.9(ii), we have :

$$\mathbf{a}(A) \subset (A \rtimes_{\mathbf{a}} \mathfrak{G})^{\tilde{\mathbf{a}}} \subset \text{Sata}$$

Therefore, the result is trivial. \square

10.12. Corollary. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A ; the dual action $\tilde{\mathfrak{a}}$ of $(\mathfrak{G})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ satisfies property (B) of 10.10.*

Proof. This is a straightforward corollary of 10.7(ii) and 10.11. \square

11. BIDUALITY

In that chapter, we prove a first version of biduality theorems (11.3, 11.4), which implies that any action satisfies properties (A) (11.5(ii)) and (B) (11.5(i)). We then get the biduality theorem (11.6) and commutation theorem (11.7).

11.1. Lemma. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A , $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ the crossed product of A by \mathfrak{G} via this action. Then, we have :*

$$[(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A) \cup 1_{b^*_{\alpha}} \widehat{M}'_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})]'' = A \rtimes_{\mathfrak{a}} \mathfrak{G}_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})$$

Proof. We have $\mathfrak{a}(A) \subset A_{b^*_{\alpha}} \mathcal{L}(H_{\Phi})$, and, therefore:

$$(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A) \subset \mathfrak{a}(A)_{b^*_{\alpha}} \mathcal{L}(H_{\Phi})$$

from which we get :

$$[(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A) \cup 1_{b^*_{\alpha}} \widehat{M}'_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})]'' \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})$$

Conversely, let $X \in \mathcal{L}(\mathfrak{H})_{b^*_{\alpha}} \widehat{M}$, such that $X_{\beta^*_{\alpha}} 1$ commutes with $(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A)$. As, for $x \in A$, we have, using 3.12(v) :

$$(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(x) = (id_{b^*_{\alpha}} \Gamma)\mathfrak{a}(x) = (1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu})^* (\mathfrak{a}(x)_{\hat{\alpha} \otimes \beta} 1) (1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu})$$

and, therefore :

$$\mathfrak{a}(x)_{\hat{\alpha} \otimes \beta} 1 = (1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu}) (\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(x) (1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu})^*$$

As X belongs to $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha}} \widehat{M}$, we get that :

$$(1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu}) (X_{\beta^*_{\alpha}} 1) = (X_{\hat{\alpha} \otimes \beta} 1) (1_{b^*_{\alpha}} \sigma_{\nu} W^o \sigma_{\nu})$$

and, therefore, we get that $X_{\hat{\alpha} \otimes \beta} 1$ commutes with $\mathfrak{a}(x)_{\hat{\alpha} \otimes \beta} 1$, which means that X belongs to $\mathfrak{a}(A)'$; as X belongs to $\mathcal{L}(\mathfrak{H})_{b^*_{\alpha}} \widehat{M}$, we finally get that X belongs to $(A \rtimes_{\mathfrak{a}} \mathfrak{G})'$. So, we have proved that :

$$[(\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A) \cup 1_{b^*_{\alpha}} \widehat{M}'_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})]' \subset [A \rtimes_{\mathfrak{a}} \mathfrak{G}_{\beta^*_{\alpha}} \mathcal{L}(H_{\Phi})]'$$

from which we get the result. \square

11.2. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A , $\tilde{\mathfrak{a}}$ the dual action of $(\widehat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, $\tilde{\tilde{\mathfrak{a}}}$ the bidual action of $(\mathfrak{G})^{oc}$ on the double crossed-product $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$; we shall denote :

$$\underline{A} = (\mathfrak{a}(A) \cup 1_{b \otimes_N \alpha} \alpha(N)')''$$

and, for any $X \in \underline{A}$:

$$\underline{\mathfrak{a}}(X) = (1_{b \otimes_N \alpha} \sigma_{\nu^o} W \sigma_{\nu^o}) (id_{b \otimes_N \alpha} \varsigma_N) (\mathfrak{a}_{b \otimes_N \alpha} id)(X) (1_{b \otimes_N \alpha} \sigma_{\nu^o} W \sigma_{\nu^o})^*$$

Moreover, for any $Y \in \mathcal{L}(\mathfrak{H}_{b \otimes_{\nu} \alpha} H_{\Phi} \otimes_{\beta} H_{\Phi})$, we define :

$$\Theta(Y) = (1_{b \otimes_{\nu} \alpha} \sigma_{\nu} W^o \sigma_{\nu}) Y (1_{b \otimes_N \alpha} \sigma_{\nu} W^o \sigma_{\nu})^* \in \mathcal{L}(\mathfrak{H}_{b \otimes_{\nu} \alpha} H_{\Phi} \otimes_{\hat{\alpha} \otimes_{\beta} \nu^o} H_{\Phi})$$

Then, for any $X \in \underline{A}$, we have :

$$\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(\underline{A}) = (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$$

$$(\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)_{b \otimes_N \alpha} \mathcal{J}_{\mathfrak{G}}) \underline{\mathfrak{a}}(X) = \tilde{\tilde{\mathfrak{a}}} \circ \Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(X)$$

and, for any $Z \in A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we get that $\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(Z) = \tilde{\mathfrak{a}}(Z)$.

Proof. Let $x \in A$; then $\mathfrak{a}(x) \in \underline{A}$, $(\mathfrak{a}_{b \otimes_N \alpha} id) \mathfrak{a}(x) = (id_{b \otimes_N \alpha} \Gamma) \mathfrak{a}(x)$, and, using 3.12, we have :

$$\Theta(\mathfrak{a}_{b \otimes_N \alpha} id) \mathfrak{a}(x) = \mathfrak{a}(x)_{\hat{\alpha} \otimes_{\beta} 1_{N^o}}$$

Let $y \in \widehat{M'}$; then $1_{b \otimes_N \alpha} y \in \underline{A}$, $(\mathfrak{a}_{b \otimes_N \alpha} id)(1_{b \otimes_N \alpha} y) = 1_{b \otimes_N \alpha} 1_{\beta \otimes_N \alpha} y$, and, using 3.10(iii) and 3.12(iv), we get :

$$\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(1_{b \otimes_N \alpha} y) = 1_{b \otimes_N \alpha} \widehat{\Gamma}^c(y)$$

So, for any $Z \in A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we get that $\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(Z) = \tilde{\mathfrak{a}}(Z)$. Let now $z \in M'$; then $1_{b \otimes_N \alpha} z \in \underline{A}$, $(\mathfrak{a}_{b \otimes_N \alpha} id)(1_{b \otimes_N \alpha} z) = 1_{b \otimes_N \alpha} 1_{\beta \otimes_N \alpha} z$, and :

$$\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(1_{b \otimes_N \alpha} z) = 1_{b \otimes_N \alpha} 1_{\hat{\alpha} \otimes_{\beta} z_{N^o}}$$

from which we get that $\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)(\underline{A}) = (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$. Moreover, using 3.6, we deduce from $(\mathfrak{a}_{b \otimes_N \alpha} id) \mathfrak{a}(x) = (id_{b \otimes_N \alpha} \Gamma) \mathfrak{a}(x)$, that, for all $x \in A$, we have $\underline{\mathfrak{a}}(\mathfrak{a}(x)) = \mathfrak{a}(x)_{\hat{\beta} \otimes_{\alpha} 1_N}$.

From which we deduce that :

$$\begin{aligned} (\Theta(\mathfrak{a}_{b \otimes_N \alpha} id)_{b \otimes_N \alpha} \mathcal{J}_{\mathfrak{G}}) \underline{\mathfrak{a}}(\mathfrak{a}(x)) &= \mathfrak{a}(x)_{\hat{\alpha} \otimes_{\beta} 1_{N^o}} \otimes_{\hat{\alpha}} 1_N \\ &= \tilde{\mathfrak{a}}(\mathfrak{a}(x))_{\hat{\beta} \otimes_{\hat{\alpha}} 1_N} \\ &= \tilde{\tilde{\mathfrak{a}}}(\tilde{\mathfrak{a}}(\mathfrak{a}(x))) \\ &= \tilde{\tilde{\mathfrak{a}}} \circ \Theta(\mathfrak{a}_{b \otimes_N \alpha} id) \mathfrak{a}(x) \end{aligned}$$

For $y \in \widehat{M}'$, we have $\underline{\mathbf{a}}(1_{b \otimes_{\alpha} y}) = 1_{b \otimes_{\alpha} y} \beta \otimes_{\alpha} 1$ and, therefore :

$$\begin{aligned}
(\Theta(\mathbf{a}_{b \otimes_{\alpha} id})_{b \otimes_{\alpha} \mathcal{J}_{\mathfrak{G}}}) \underline{\mathbf{a}}(1_{b \otimes_{\alpha} y}) &= 1_{b \otimes_{\alpha} \widehat{\Gamma}^c(y)} \beta \otimes_{\alpha} 1 \\
&= \tilde{\mathbf{a}}(1_{b \otimes_{\alpha} y}) \beta \otimes_{\alpha} 1 \\
&= \tilde{\mathbf{a}}(\tilde{\mathbf{a}}(1_{b \otimes_{\alpha} y})) \\
&= \tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b \otimes_{\alpha} id})(1_{b \otimes_{\alpha} y})
\end{aligned}$$

from which we deduce that, for all $Z \in A \rtimes_{\mathbf{a}} \mathfrak{G}$, we have :

$$(\Theta(\mathbf{a}_{b \otimes_{\alpha} id})_{b \otimes_{\alpha} \mathcal{J}_{\mathfrak{G}}}) \underline{\mathbf{a}}(Z) = \tilde{\mathbf{a}}(\tilde{\mathbf{a}}(Z)) = \tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b \otimes_{\alpha} id})(Z)$$

For $z \in M'$, we have $(\mathbf{a}_{b \otimes_{\alpha} id})(1_{b \otimes_{\alpha} z}) = 1_{b \otimes_{\alpha} z} \beta \otimes_{\alpha} z$ and :

$$\underline{\mathbf{a}}(1_{b \otimes_{\alpha} z}) = (1_{b \otimes_{\alpha} \sigma_{\nu^o}})(1_{b \otimes_{\alpha} (W(1_{\beta \otimes_{\alpha} z} W^*))})(1_{b \otimes_{\alpha} \sigma_{\nu^o}})$$

As $W = (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) W^* (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi})$, we get that :

$$\underline{\mathbf{a}}(1_{b \otimes_{\alpha} z}) = (id_{b \otimes_{\alpha} \varsigma_{N^o}})(1_{b \otimes_{\alpha} (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \Gamma(J_{\Phi} z J_{\Phi}) (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi})})$$

and :

$$\begin{aligned}
(\Theta(\mathbf{a}_{b \otimes_{\alpha} id})_{b \otimes_{\alpha} \mathcal{J}_{\mathfrak{G}}}) \underline{\mathbf{a}}(1_{b \otimes_{\alpha} z}) &= 1_{b \otimes_{\alpha} 1_{\widehat{\alpha} \otimes_{\beta} \Gamma^{oc}(z)}} \\
&= \tilde{\mathbf{a}}(1_{b \otimes_{\alpha} 1_{\widehat{\alpha} \otimes_{\beta} z}}) \\
&= \tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b \otimes_{\alpha} id})(1_{b \otimes_{\alpha} z})
\end{aligned}$$

from which we get the result. \square

11.3. Proposition. Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathbf{a}) an action of \mathfrak{G} on A , $\tilde{\mathbf{a}}$ the dual action of $(\widehat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathbf{a}} \mathfrak{G}$, $\tilde{\tilde{\mathbf{a}}}$ the bidual action of $(\mathfrak{G})^{oc}$ on the double crossed-product $(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o$; then, with the notations of 11.2 :

- (i) $(1_{b \otimes_{\alpha} \widehat{\beta}}, \underline{\mathbf{a}})$ is an action of \mathfrak{G} on \underline{A} ;
- (ii) this action satisfies property (B);
- (iii) $\underline{A}^{\underline{\mathbf{a}}} = A \rtimes_{\mathbf{a}} \mathfrak{G}$

Proof. Thanks to 11.2, we get that, for any $n \in N$:

$$\begin{aligned}
\underline{\mathbf{a}}(1_{b \otimes_{\alpha} \widehat{\beta}(n)}) &= (id_{b \otimes_{\alpha} \varsigma_{N^o}})(1_{b \otimes_{\alpha} (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \Gamma(\alpha(n)) (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi})}) \\
&= 1_{b \otimes_{\alpha} 1_{\beta \otimes_{\alpha} \beta(n)}}
\end{aligned}$$

On the other hand, we get, using 11.2, for any $X \in \underline{A}$, that :

$$(\Theta(\mathbf{a}_{b \otimes_{\alpha} id})_{b \otimes_{\alpha} \mathcal{J}_{\mathfrak{G}}}) \underline{\mathbf{a}}(X) \beta \otimes_{\alpha} id \underline{\mathbf{a}}(X)$$

is equal to :

$$\begin{aligned}
(\tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}}) \underline{\mathbf{a}}(X) &= (\tilde{\mathbf{a}}_{\beta \otimes_{\hat{\alpha}} id} \tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b_N^* \alpha} id)(X) \\
&= (id_{b_N^* \alpha} \Gamma^{oc}) \tilde{\mathbf{a}} \circ \Theta(\mathbf{a}_{b_N^* \alpha} id)(X) \\
&= (\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \Gamma^{oc} \circ \mathcal{J}_{\mathfrak{G}}) \underline{\mathbf{a}}(X)
\end{aligned}$$

which is equal to $(\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}}) (id_{\beta^* \alpha} \Gamma) \underline{\mathbf{a}}(X)$, from which we get (i). On the other hand, we have :

$$\begin{aligned}
(\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}})(\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G}) &= (\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}})(\underline{\mathbf{a}}(\underline{A}) \cup 1_{b_N^* \alpha} 1_{\beta \otimes_{\hat{\alpha}} \alpha} \widehat{M}')'' \\
&= (\tilde{\mathbf{a}}[(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o] \cup 1_{b_N^* \alpha} 1_{\beta \otimes_{\hat{\alpha}} \alpha} \widehat{M}')'' \\
&= [(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o] \rtimes_{\tilde{\mathbf{a}}} \mathfrak{G}^{oc}
\end{aligned}$$

Using then 9.4(iii), we get that :

$$(\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}})((\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G})^{\tilde{\mathbf{a}}}) = (\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}})((\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G} \cap \underline{A}_{b_N^* \alpha} M))$$

which, using 11.2 and 9.4(iii) again, is equal to :

$$[(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o] \rtimes_{\tilde{\mathbf{a}}} \mathfrak{G}^{oc} \cap [(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o]_{\hat{\alpha}^* \beta} M' = ([(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o] \rtimes_{\tilde{\mathbf{a}}} \mathfrak{G}^{oc})^{\tilde{\mathbf{a}}}_{N^o}$$

which, using 10.12, is equal to $\tilde{\mathbf{a}}[(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o]$, and, therefore, by 11.2 again, is equal to $(\Theta(\mathbf{a}_{b_N^* \alpha} id)_{b_N^* \alpha} \mathcal{J}_{\mathfrak{G}}) \underline{\mathbf{a}}(\underline{A})$. From which we get that $(\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G})^{\tilde{\mathbf{a}}} = \underline{\mathbf{a}}(\underline{A})$ which is (ii).

Using 11.3, we get that X in \underline{A} belongs to $\underline{A}^{\mathbf{a}}$ if and only if $\Theta(\mathbf{a}_{b_N^* \alpha} id)(X)$ belongs to $[(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o]^{\tilde{\mathbf{a}}}$, which, using 10.12, is equal to $\tilde{\mathbf{a}}(A \rtimes_{\mathbf{a}} \mathfrak{G})$, or, using 11.3 again, is equal to $\Theta(\mathbf{a}_{b_N^* \alpha} id)(A \rtimes_{\mathbf{a}} \mathfrak{G})$; from which we get (iii). \square

11.4. Proposition. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathbf{a}) an action of \mathfrak{G} on A , $\tilde{\mathbf{a}}$ the dual action of $(\mathfrak{G})^c$ on the crossed product $A \rtimes_{\mathbf{a}} \mathfrak{G}$, $\tilde{\tilde{\mathbf{a}}}$ the bidual action of $(\mathfrak{G})^{oc}$ on the double crossed-product $(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^o$; let's use the notations of 11.2 and 11.3. Then, we have :*

$$\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G} = (id_{b_N^* \alpha} \varsigma_{N^o})[(1_{b_N^* \alpha} W)(A \rtimes_{\mathbf{a}} \mathfrak{G}_{\beta^* \alpha} \mathcal{L}(H_{\Phi}))(1_{b_N^* \alpha} W)^*]$$

Proof. By definition of \underline{A} , we have :

$$\underline{\mathbf{a}}(\underline{A}) = (id_{b_N^* \alpha} \varsigma_{N^o})[(1_{b_N^* \alpha} W)((\mathbf{a}_{b_N^* \alpha} id) \mathbf{a}(A) \cup 1_{b_N^* \alpha} 1_{\beta \otimes_{\hat{\alpha}} \alpha} \alpha(N)')''(1_{b_N^* \alpha} W)^*]$$

and $\underline{A} \rtimes_{\underline{\mathbf{a}}} \mathfrak{G} = (\underline{\mathbf{a}}(\underline{A}) \cup 1_{b_N^* \alpha} 1_{\beta \otimes_{\hat{\alpha}} \alpha} \widehat{M}')''$. But, on the other hand, using 3.11(iii), we get that $W^*(\widehat{M}'_{\alpha \otimes_{\hat{\beta}} 1})W$ is equal to :

$$\begin{aligned}
(J_{\hat{\Phi} \alpha \otimes_{\hat{\beta}} J_{\Phi}} W)(J_{\hat{\Phi} \alpha \otimes_{\hat{\beta}} J_{\Phi}})(\widehat{M}'_{\alpha \otimes_{\hat{\beta}} 1})(J_{\hat{\Phi} \beta \otimes_{\alpha} J_{\Phi}}) W^*(J_{\hat{\Phi} \beta \otimes_{\alpha} J_{\Phi}}) \\
= (J_{\hat{\Phi} \alpha \otimes_{\hat{\beta}} J_{\Phi}})_{N^o} \widehat{\Gamma}^o(\widehat{M})(J_{\hat{\Phi} \beta \otimes_{\alpha} J_{\Phi}})_N
\end{aligned}$$

Therefore, we get that :

$$\widehat{M}'_{\alpha \otimes_{\beta} 1} = W(J_{\widehat{\Phi}} \alpha \otimes_{\beta} J_{\Phi}) \widehat{\Gamma}^o(\widehat{M})(J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) W^*$$

and :

$$1_{\beta \otimes_{\alpha} \widehat{M}'} = \varsigma_{N^o}[W(J_{\widehat{\Phi}} \alpha \otimes_{\beta} J_{\Phi}) \widehat{\Gamma}^o(\widehat{M})(J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) W^*]$$

Moreover, we have :

$$\begin{aligned} (J_{\widehat{\Phi}} \alpha \otimes_{\beta} J_{\Phi}) \widehat{\Gamma}^o(\widehat{M})(J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \cup (1_{\beta \otimes_{\alpha} \alpha(N)'})'' &= \\ &= (J_{\widehat{\Phi}} \alpha \otimes_{\beta} J_{\Phi}) [\widehat{\Gamma}^o(\widehat{M}) \cup 1_{\alpha \otimes_{\beta} \hat{\beta}(N)'}]'' ((J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \end{aligned}$$

which, by 7.2 applied to $\widehat{\mathfrak{G}}^o$, is equal to :

$$(J_{\widehat{\Phi}} \alpha \otimes_{\beta} J_{\Phi}) (\widehat{M}_{\alpha \otimes_{\beta} \mathcal{L}(H_{\Phi})}) (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) = \widehat{M}'_{\beta \otimes_{\alpha} \mathcal{L}(H_{\Phi})}$$

Therefore, we get that $\underline{A} \rtimes_{\underline{\mathfrak{a}}} \mathfrak{G}$ is equal to :

$$(id_{b^*_{\alpha} \varsigma_{N^o}})[(1_{b \otimes_{\alpha} W})((\mathfrak{a}_{b^*_{\alpha}} id)\mathfrak{a}(A) \cup 1_{b \otimes_{\alpha} \widehat{M}'_{\beta \otimes_{\alpha} \mathcal{L}(H_{\Phi})}})(1_{b \otimes_{\alpha} W})^*]$$

which, using 11.1, gives the result. \square

11.5. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A , $\tilde{\mathfrak{a}}$ the dual action of $(\widehat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$. Then :*

(i) \mathfrak{a} satisfies property (B), i.e. we have :

$$(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}} = \mathfrak{a}(A)$$

(ii) \mathfrak{a} satisfies property (A), i.e. we have :

$$(\mathfrak{a}(A) \cup 1_{b \otimes_{\alpha} \alpha(N)'})'' = A_{b^*_{\alpha} \mathcal{L}(H_{\Phi})}$$

Proof. Let's use the constructions and notations of 11.2, 11.3 and 11.4. Using 9.4(iii), we have $(\underline{A} \rtimes_{\underline{\mathfrak{a}}} \mathfrak{G})^{\tilde{\mathfrak{a}}} = \underline{A} \rtimes_{\underline{\mathfrak{a}}} \mathfrak{G} \cap \mathcal{L}(\mathfrak{H}_{b \otimes_{\alpha} H_{\Phi}})_{\beta^*_{\alpha} M}$, and, therefore, using 11.4 and 9.4(iii) again, we have :

$$(\underline{A} \rtimes_{\underline{\mathfrak{a}}} \mathfrak{G})^{\tilde{\mathfrak{a}}} = (id_{b^*_{\alpha} \varsigma_{N^o}})[(1_{b \otimes_{\alpha} W})((A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}_{\beta^*_{\alpha} \mathcal{L}(H_{\Phi})})(1_{b \otimes_{\alpha} W})^*]$$

But, by 11.3, we have :

$$(\underline{A} \rtimes_{\underline{\mathfrak{a}}} \mathfrak{G})^{\tilde{\mathfrak{a}}} = \underline{\mathfrak{a}}(\underline{A}) = (id_{b^*_{\alpha} \varsigma_{N^o}})[(1_{b \otimes_{\alpha} W})(\mathfrak{a}_{b^*_{\alpha}} id)(\underline{A})_{\beta^*_{\alpha} \mathcal{L}(H_{\Phi})})(1_{b \otimes_{\alpha} W})^*]$$

from which we deduce that $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}_{\beta^*_{\alpha} \mathcal{L}(H_{\Phi})} = (\mathfrak{a}_{b^*_{\alpha}} id)(\underline{A})$, from which we get that

$(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}_{\beta^*_{\alpha} \mathcal{L}(H_{\Phi})} \subset \mathfrak{a}(A)_{\beta^*_{\alpha} \mathcal{L}(H_{\Phi})}$ and we deduce that $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}} \subset \mathfrak{a}(A)$, which,

thanks to 9.4(iii), gives (i).

But we have now :

$$\begin{aligned}
(\mathfrak{a}_{b^*_{N\alpha}} id)(A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)) &= \mathfrak{a}(A)_{\beta^*_{N\alpha}} \mathcal{L}(H_\Phi) \\
&= (A \rtimes_{\mathfrak{a}} \mathfrak{G})_{\beta^*_{N\alpha}}^{\tilde{\mathfrak{a}}} \mathcal{L}(H_\Phi) \\
&= (\mathfrak{a}_{b^*_{N\alpha}} id)(\underline{A})
\end{aligned}$$

which gives (ii). \square

11.6. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A , $\tilde{\mathfrak{a}}$ the dual action of $(\widehat{\mathfrak{G}})^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, $\tilde{\tilde{\mathfrak{a}}}$ the bidual action of $(\mathfrak{G})^{oc}$ on the double crossed-product $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$; let us define, for any $X \in A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)$:*

$$\underline{\mathfrak{a}}(X) = (1_{b \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})(id_{b^*_{N\alpha}} \varsigma_N)(\mathfrak{a}_{b^*_{N\alpha}} id)(X)(1_{b \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})^*$$

Moreover, for any $Y \in \mathcal{L}(\mathfrak{H}_{b \otimes_{\alpha} H_\Phi \otimes_{\alpha} H_\Phi})$, we define :

$$\Theta(Y) = (1_{b \otimes_{\alpha} \sigma_{\nu} W^o \sigma_{\nu}}) Y (1_{b \otimes_{\alpha} \sigma_{\nu} W^o \sigma_{\nu}})^* \in \mathcal{L}(\mathfrak{H}_{b \otimes_{\alpha} H_\Phi \otimes_{\alpha} H_\Phi})$$

Then, for any $X \in A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)$, we have :

$$\Theta(\mathfrak{a}_{b^*_{N\alpha}} id)(A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)) = (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$$

$$(\Theta(\mathfrak{a}_{b^*_{N\alpha}} id)_{b^*_{N\alpha}} \mathcal{I}_{\mathfrak{G}}) \underline{\mathfrak{a}}(X) = \tilde{\tilde{\mathfrak{a}}} \circ \Theta(\mathfrak{a}_{b^*_{N\alpha}} id)(X)$$

and, for any $Z \in A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we get that $\Theta(\mathfrak{a}_{b^*_{N\alpha}} id)(Z) = \tilde{\mathfrak{a}}(Z)$.

Then $(1_{b \otimes_{\alpha} \hat{\beta}}, \underline{\mathfrak{a}})$ is an action of \mathfrak{G} on $A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)$, and :

$$(A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi))^{\underline{\mathfrak{a}}} = A \rtimes_{\mathfrak{a}} \mathfrak{G}$$

Moreover, for any $X \in (A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi))^+$, we have :

$$T_{\tilde{\tilde{\mathfrak{a}}}} \Theta(\mathfrak{a}_{b^*_{N\alpha}} id) = \Theta(\mathfrak{a}_{b^*_{N\alpha}} id) T_{\underline{\mathfrak{a}}}(X) = \tilde{\mathfrak{a}}(T_{\underline{\mathfrak{a}}}(X))$$

Proof. This result is given by 11.2, 11.3 and 11.5(ii). \square

11.7. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid, ${}_a \mathfrak{H}_b$ a $N - N$ bimodule, A a von Neumann algebra such that $b(N) \subset A \subset a(N)'$, V a corepresentation of \mathfrak{G} on ${}_a \mathfrak{H}_b$ which implements an action (b, \mathfrak{a}) of \mathfrak{G} on A and an action (a, \mathfrak{a}') of \mathfrak{G}^o on A' . Then :*

$$(i) A \rtimes_{\mathfrak{a}} \mathfrak{G} = V(\mathcal{L}(\mathfrak{H})_{a^*_{\beta}} \widehat{M'}) V^* \cap A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)$$

$$(ii) A' \rtimes_{\mathfrak{a}'} \mathfrak{G}^o = V^*(A \rtimes_{\mathfrak{a}} \mathfrak{G})' V.$$

Proof. Using 11.6, we get that $X \in A_{b^*_{N\alpha}} \mathcal{L}(H_\Phi)$ belongs to $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ if and only if

$X \in A_{b^*_{N\alpha}} \hat{\beta}(N)'$, and $\underline{\mathfrak{a}}(X) = X_{\hat{\beta} \otimes_{\alpha} 1}$, or, equivalently :

$$(\mathfrak{a}_{b^*_{N\alpha}} id) \mathfrak{a}(X) = (1_{b \otimes_{\alpha} W^*})(1_{b \otimes_{\alpha} \sigma_{\nu}})(X_{\hat{\beta} \otimes_{\alpha} 1})(1_{b \otimes_{\alpha} \sigma_{\nu}})(1_{b \otimes_{\alpha} W})$$

which means (7.4) that X belongs to $A_{b^*_{\alpha} \hat{\beta}(N)'} \cap \text{Sat}V$, and, therefore, by 7.4(i), we get :

$$A \rtimes_{\mathfrak{a}} \mathfrak{G} = V(\mathcal{L}(\mathfrak{H})_{a^*_{\beta} \widehat{M}'} V^* \cap A_{b^*_{\alpha} \hat{\beta}(N)'})$$

We have seen in 7.4 that $V(\mathcal{L}(\mathfrak{H})_{a^*_{\beta} \widehat{M}'} V^* \cap A_{b^*_{\alpha} \hat{\beta}(N)'})$ belongs to $a(N)'_{b^*_{\alpha} \hat{\beta}(N)'}$, which gives (i).

From (i), we get that :

$$(A \rtimes_{\mathfrak{a}} \mathfrak{G})' = (V(1_{b \otimes_{\alpha} \widehat{M}}) V^* \cup A'_{b \otimes_{\alpha} 1})''$$

and :

$$V^*(A \rtimes_{\mathfrak{a}} \mathfrak{G})' V = [V(A'_{b \otimes_{\alpha} 1}) V^* \cup 1_{b \otimes_{\alpha} \widehat{M}}]''$$

which gives (ii). \square

11.8. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra acting on a Hilbert space \mathfrak{H} , (b, \mathfrak{a}) an action of \mathfrak{G} on A ; then, \mathfrak{a} is saturated, i.e. we have :*

$$\mathfrak{a}(A) = \{X \in A_{b^*_{\alpha} M}, (\mathfrak{a}_{b^*_{\alpha} id}(X) = (id_{b^*_{\alpha} \Gamma})(X))\}$$

Proof. Let us suppose that there exists a corepresentation V on \mathfrak{H} which implements \mathfrak{a} ; using 7.6 and 11.5(i), we get that \mathfrak{a} is saturated (7.3), which means that we have the result. Applying this fact to 6.7, we get that $(\mathfrak{a}_{b^*_{\alpha} id})\mathfrak{a}(A)$ is equal to :

$$\{X \in \mathfrak{a}(A)_{b^*_{\alpha} M}, (\mathfrak{a}_{b^*_{\alpha} id} \beta^*_{\alpha} id)(X) = (id_{b^*_{\alpha} id} \beta^*_{\alpha} \Gamma)(X)\}$$

which is the image under $(\mathfrak{a}_{b^*_{\alpha} id})$ of :

$$\{Y \in A_{b^*_{\alpha} M}; (\mathfrak{a}_{b^*_{\alpha} id} \beta^*_{\alpha} id)(\mathfrak{a}_{b^*_{\alpha} id}(Y) = (id_{b^*_{\alpha} id} \beta^*_{\alpha} \Gamma)(\mathfrak{a}_{b^*_{\alpha} id}(Y))\}$$

As, $(id_{b^*_{\alpha} id} \beta^*_{\alpha} \Gamma)(\mathfrak{a}_{b^*_{\alpha} id}(Y) = (\mathfrak{a}_{b^*_{\alpha} id} \beta^*_{\alpha} id)(id_{b^*_{\alpha} \Gamma})(Y)$, we get the result. \square

11.9. Theorem. *Let $\mathfrak{G} = (N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid; let A be a von Neumann algebra, (b, \mathfrak{a}) an action of \mathfrak{G} on A and $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ its crossed product. Then, the linear subspace generated by all elements of the form $(1_{b \otimes_{\alpha} a})\mathfrak{a}(x)$, with $a \in \widehat{M}'$ and $x \in A$, is dense in $A \rtimes_{\mathfrak{a}} \mathfrak{G}$.*

Proof. Using 11.5(i), this is a straightforward corollary of 10.1(iii). \square

11.10. Remark. These theorems (11.5, 11.6, 11.8) are the generalization, up to the measured quantum groupoid framework, of biduality theorems ([V2] 2.6, 2.7) obtained for actions of locally compact quantum groups, and we followed the same strategy as [V2] (which is, in fact, the strategy of [ES1]). These theorems generalizes as well the duality theorems for groupoid actions ([Y3], 6.5) and integrable groupoid coactions ([Y3], 7.8).

12. CHARACTERIZATION OF CROSSED-PRODUCTS

On this chapter, inspired by ([ES1], V), a measured quantum groupoid \mathfrak{G} be given, we characterize crossed-products among the von Neumann algebras on which there exists an action of $\widehat{\mathfrak{G}}^c$ (12.3). A corollary (12.5) will be used in chapter 13.

12.1. **Notations.** In this chapter, we shall consider a measured quantum groupoid \mathfrak{G} , its dual $\widehat{\mathfrak{G}}$, and a von Neumann algebra B on a Hilbert space \mathfrak{H} . We suppose that there exist :

- (i) a normal faithful morphism μ from \widehat{M}' into B ;
- (ii) a normal faithful morphism \mathfrak{b} from B into $B_{\mu \circ \hat{\alpha} * \beta} \widehat{M}'$, such that $(\mu \circ \hat{\alpha}, \mathfrak{b})$ is an action of $\widehat{\mathfrak{G}}^c$ on B ;
- (iii) we have, moreover : $\mathfrak{b} \circ \mu = (\mu_{\hat{\alpha} * \beta} id) \widehat{\Gamma}^c$.

This last formula implies that, for any $n \in N$, $\mathfrak{b} \circ \mu(\beta(n))$ belongs to $(\mu \circ \hat{\alpha}(N))'$ and :

$$\mathfrak{b} \circ \mu(\beta(n)) = (\mu_{\hat{\alpha} * \beta} id) \widehat{\Gamma}^c(\beta(n)) = (\mu_{\hat{\alpha} * \beta} id)(\beta(n)_{\hat{\alpha} \otimes \beta} 1) = \mu(\beta(n))_{\mu \circ \hat{\alpha} \otimes \beta} 1$$

and, therefore, by definition, $\mu \circ \beta(N) \subset B^a$.

12.2. **Lemma.** *Let's take the hypothesis and notations of 12.1. Then*

(i) *there exists a corepresentation Y of \mathfrak{G} on the bimodule ${}_{\mu \circ \hat{\alpha}} \mathfrak{H}_{\mu \circ \beta}$. such that, for any $\xi \in D({}_\alpha H_\Phi, \nu)$, $\eta \in D((H_\Phi)_\beta, \nu^o)$, we have :*

$$(id * \omega_{\xi, \eta})(Y) = \mu[(id * \omega_{\xi, \eta})((\sigma_\nu W^o \sigma_\nu)^*)]$$

(ii) *for any $b' \in B'$, we have $Y(b'_{\mu \circ \hat{\alpha} \otimes \beta} 1) = (b'_{\mu \circ \beta \otimes \alpha} 1)Y$;*

(iii) *for any $b \in B$, $Y\mathfrak{b}(x)Y^*$ belongs to $B^b_{\mu \circ \beta * \alpha} \mathcal{L}(H_\Phi)$; moreover, if $x \in B^b$, then $Y(x_{\mu \circ \beta \otimes \alpha} 1)Y^*$ belongs to $B^b_{\mu \circ \beta * \alpha} M$;*

(iv) *the corepresentation Y of \mathfrak{G} implements on B^b an action of \mathfrak{G} , we shall denote \mathfrak{d} . More precisely, $(\mu \circ \beta, \mathfrak{d})$ is an action of \mathfrak{G} on B^b .*

Proof. By 3.10(v), H_Φ can be considered as the standard Hilbert space of \widehat{M}' , we can identify with \widehat{M}^o . Therefore, the Hilbert space \mathfrak{H} is isomorphic (2.3) to the relative tensor product $\mathfrak{H}_{\mu \otimes \hat{\pi}} (H_\Phi)$, where $\hat{\pi}$ denotes here the canonical representation of \widehat{M} (resp. \widehat{M}')

on H_Φ , and this isomorphism is an isomorphism of right \widehat{M} -modules, which means that, modulo the identification of these Hilbert spaces, we have, for all $x \in \widehat{M}'$, $\mu(x) = 1_{\mu \otimes \hat{\pi}} x$.

Let us put $Y = 1_{\mathfrak{H}} \mu \otimes \hat{\pi} (\sigma_\nu W^o \sigma_\nu)^*$. So, Y is a unitary from $\mathfrak{H}_{\mu \circ \hat{\alpha} \otimes \beta} H_\Phi$ onto $\mathfrak{H}_{\mu \circ \beta \otimes \alpha} H_\Phi$.

As $(\sigma_\nu W^o \sigma_\nu)^*$ is a corepresentation of \mathfrak{G} on the $N - N$ bimodule ${}_{\hat{\alpha}}(H_\Phi)_\beta$ (5.6), we get easily that Y is a corepresentation of \mathfrak{G} on the $N - N$ bimodule ${}_{\mu \circ \hat{\alpha}} \mathfrak{H}_{\mu \circ \beta}$.

Moreover, we have then :

$$\begin{aligned} (id * \omega_{\xi, \eta})(Y) &= (id_{\widehat{M}} * \mu \otimes \hat{\pi} id * \omega_{\xi, \eta})[1_{\widehat{M}} \mu \otimes \hat{\pi} (\sigma_\nu W^o \sigma_\nu)^*] \\ &= 1_{\widehat{M}} \mu \otimes \hat{\pi} (id * \omega_{\xi, \eta})((\sigma_\nu W^o \sigma_\nu)^*) \\ &= \mu[(id * \omega_{\xi, \eta})((\sigma_\nu W^o \sigma_\nu)^*)] \end{aligned}$$

which gives (i). Then (ii) is a straightforward corollary of (i). From (ii), we get easily that $Y\mathfrak{b}(b)Y^*$ belongs to $B_{\mu \circ \beta * \alpha} \mathcal{L}(H_\Phi)$, and, therefore, that $(id_{\mu \circ \beta * \alpha} \omega_\xi)(Y\mathfrak{b}(b)Y^*)$

belongs to B . Let $(\eta_i)_{i \in I}$ be an orthogonal (β, ν^o) -basis of H_Φ . We get that the element

$(id_{\mu \circ \beta * \alpha} \omega_\xi)(Y \mathfrak{b}(b) Y^*)$ is equal to the sum :

$$\begin{aligned} \sum_{i,j} (id_{\mu \circ \beta * \alpha} \omega_\xi)(Y \theta^{\beta, \nu^o}(\eta_i, \eta_i) \mathfrak{b}(b) \theta^{\beta, \nu^o}(\eta_j, \eta_j) Y^*) &= \\ &= \sum_{i,j} (id * \omega_{\xi, \eta_i})(Y) (id_{\mu \circ \hat{\alpha} * \beta} \omega_{\eta_i, \eta_j}) \mathfrak{b}(b) [(id * \omega_{\xi, \eta_j})(Y)]^* \end{aligned}$$

And, as we know that $(\sigma_\nu W^o \sigma_\nu)^* = \widehat{W}^c$, we get that :

$$\mathfrak{b}[(id_{\mu \circ \beta * \alpha} \omega_\xi)(Y \mathfrak{b}(b) Y^*)]$$

is equal to the sum over (i, j) of all following elements :

$$\mathfrak{b} \circ \mu[(id * \omega_{\xi, \eta_i})(\widehat{W}^c)(id_{\mu \circ \hat{\alpha} * \beta} id_{\hat{\alpha} * \beta} \omega_{\eta_i, \eta_j})(id_{\mu \circ \hat{\alpha} * \beta} \widehat{\Gamma}^c) \mathfrak{b}(b)] \mathfrak{b} \circ \mu(id * \omega_{\xi, \eta_j})(\widehat{W}^c)^*$$

Using the pentagonal relation for \widehat{W}^c , we get that :

$$\mathfrak{b} \circ \mu[(id * \omega_{\xi, \eta_i})(\widehat{W}^c)] = (\mu_{\hat{\alpha} * \beta} id) \widehat{\Gamma}^c[(id * \omega_{\xi, \eta_i})(\widehat{W}^c)]$$

is equal to $(\mu_{\hat{\alpha} * \beta} id)(id * id * \omega_{\xi, \eta_i})[\sigma_{\beta, \hat{\alpha}}^{2,3}(\widehat{W}^c_{\alpha \otimes \beta} 1)(1_{\hat{\alpha} \otimes \beta} \sigma_\nu)(1_{\hat{\alpha} \otimes \beta} \widehat{W}^c)]$. And, as $(id_{\mu \circ \hat{\alpha} * \beta} \widehat{\Gamma}^c) \mathfrak{b}(b)$ is equal to $(1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \widehat{W}^c)^* (1_{\mu \circ \hat{\alpha} \otimes \beta} \sigma_{\nu^o}) (\mathfrak{b}(b)_{\hat{\alpha} \otimes \beta} 1) (1_{\mu \circ \hat{\alpha} \otimes \beta} \sigma_{\nu^o})^* (1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \widehat{W}^c)$, we obtain finally that this sum is equal to :

$$(\mu_{\hat{\alpha} * \beta} id)(id * id * \omega_\xi)[\sigma_{\beta, \hat{\alpha}}^{2,3}(\widehat{W}^c \mathfrak{b}(b) (\widehat{W}^c)^*_{\alpha \otimes \beta} 1)(\sigma_{\beta, \hat{\alpha}}^{2,3})^*]$$

which can be written :

$$(id_{\mu \circ \beta * \alpha} \omega_\xi)(\sigma_{\beta, \hat{\alpha}}^{2,3}(Y \mathfrak{b}(b) Y^*_{\hat{\alpha} \otimes \beta} 1)(\sigma_{\beta, \hat{\alpha}}^{2,3})^*) = (id_{\mu \circ \beta * \alpha} \omega_\xi)(Y \mathfrak{b}(b) Y^*)_{\hat{\alpha} \otimes \beta} 1$$

and, therefore, we get that $Y \mathfrak{b}(b) Y^*$ belongs to $B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_\Phi)$.

Let $x \in B^{\mathfrak{b}} \subset \mu \circ \hat{\alpha}(N)'$. We have obtained that $Y(x_{\mu \circ \hat{\alpha} \otimes \beta} 1) Y^*$ belongs to $B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_\Phi)$; as it belongs clearly to $\mathcal{L}(\mathfrak{H})_{\mu \circ \beta * \alpha} M$, we get that $Y(x_{\mu \circ \hat{\alpha} \otimes \beta} 1) Y^*$ belongs to $B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} M$, which finishes the proof of (iii).

If $n \in N$, we have :

$$\begin{aligned} \mathfrak{d}(\mu \circ \beta(n)) &= (1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \widehat{W}^c)(1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \beta(n)_{\hat{\alpha} \otimes \beta} 1)(1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \widehat{W}^c)^* \\ &= 1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} \Gamma(\beta(n)) \\ &= 1_{\mathfrak{H}} \mu_{\widehat{M}} \otimes_{\widehat{\pi}} 1_{\beta \otimes \alpha} \beta(n) \\ &= 1_{\mu \circ \beta \otimes \alpha} \beta(n) \end{aligned}$$

which gives (iv). □

12.3. **Theorem.** Let \mathfrak{G} be a measured quantum groupoid, $\widehat{\mathfrak{G}}$ its dual measured quantum groupoid, B a von Neumann algebra on a Hilbert space \mathfrak{H} . Then, are equivalent :

(i) there exists a von Neumann algebra A , an action (b, \mathfrak{a}) of \mathfrak{G} on A , and an isomorphism \mathcal{J} between the von Neumann algebras B and $A \rtimes_{\mathfrak{a}} \mathfrak{G}$.

(ii) there exists a normal faithful morphism μ from \widehat{M}' into B , a normal faithful morphism \mathfrak{b} from B into $B_{\mu \circ \hat{\alpha} * \beta} \widehat{M}'$ such that $(\mu \circ \hat{\alpha}, \mathfrak{b})$ is an action of $\widehat{\mathfrak{G}}^c$ on B which satisfies

$$\mathfrak{b} \circ \mu = (\mu_{\hat{\alpha} * \beta} id) \widehat{\Gamma}^c.$$

Moreover, we have then :

$$\mathfrak{b} \circ \mathcal{J} = (\mathcal{J}_{\hat{\alpha} * \beta} id) \mathfrak{b}$$

$$\mathcal{J} \circ T_{\mathfrak{b}} = T_{\tilde{\mathfrak{a}}} \circ \mathcal{J}$$

$$\mathcal{J}(B^{\mathfrak{b}}) = \mathfrak{a}(A)$$

$$B = (B^{\mathfrak{b}} \cup \mu(\widehat{M}'))''$$

and, for any $x \in \widehat{M}'$, we have :

$$\mathcal{J} \circ \mu(x) = 1_{\mathfrak{b} \otimes_{\alpha} x}$$

Proof. Let us suppose (i); then, if we define \mathfrak{b} by $\mathfrak{b} = (\mathcal{J}_{\hat{\alpha} * \beta} id)^{-1} \circ \tilde{\mathfrak{a}} \circ \mathcal{J}$, and, for $x \in \widehat{M}'$, if we put $\mu(x) = \mathcal{J}^{-1}(1_{\mathfrak{b} \otimes_{\alpha} x})$, then, we have (ii).

Let us suppose (ii); we can use all notations and results of 12.2. Let us consider the action $(\mu \circ \beta, \mathfrak{d})$ of \mathfrak{G} on $B^{\mathfrak{b}}$; we have :

$$\begin{aligned} B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G} &= (\mathfrak{d}(B^{\mathfrak{b}}) \cup 1_{\mu \circ \beta \otimes_{\alpha}} \widehat{M}')'' \\ &= (Y(B^{\mathfrak{b}}_{\mu \circ \hat{\alpha} \otimes \beta} 1) Y^* \cup 1_{\mu \circ \beta \otimes_{\alpha}} \widehat{M}')'' \end{aligned}$$

But, for any $x \in \widehat{M}'$, we have :

$$\begin{aligned} Y^*(1_{\mu \circ \beta \otimes_{\alpha}} x) Y &= (1_{\mathfrak{H}} \mu \otimes_{\widehat{M}} \widehat{W}^{c*}) (1_{\mathfrak{H}} \mu \otimes_{\widehat{M}} 1_{\mu \circ \beta \otimes_{\alpha}} x) (1_{\mathfrak{H}} \mu \otimes_{\widehat{M}} \widehat{W}^c) \\ &= 1_{\mathfrak{H}} \mu \otimes_{\widehat{M}} \widehat{\Gamma}^c(x) \\ &= (\mu_{\hat{\alpha} * \beta} id) \widehat{\Gamma}^c(x) \\ &= \mathfrak{b} \circ \mu(x) \end{aligned}$$

and, therefore, we have :

$$\begin{aligned} B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G} &= (Y(B^{\mathfrak{b}}_{\mu \circ \hat{\alpha} \otimes \beta} 1) Y^* \cup 1_{\mu \circ \beta \otimes_{\alpha}} \widehat{M}')'' \\ &= (Y \mathfrak{b}(B^{\mathfrak{b}}) Y^* \cup Y \mathfrak{b}(\mu(\widehat{M}')) Y^*)'' \\ &= Y \mathfrak{b}((B^{\mathfrak{b}} \cup \mu(\widehat{M}'))'' Y^* \\ &\subset Y \mathfrak{b}(B) Y^* \end{aligned}$$

For $y \in B$, let us define $\mathcal{J}(y) = Y\mathfrak{b}(y)Y^*$; so, we get $B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G} \subset \mathcal{J}(B)$.
Using 12.2(iii), we get that $\mathcal{J}(B) \subset B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})$, and, therefore :

$$\mathcal{J}(B) \subset B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi}) \cap Y(\mathcal{L}(\mathfrak{H})_{\mu \circ \hat{\alpha} * \beta} \widehat{M'})Y^*$$

which, thanks to 11.7(i), is equal to $B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G}$.

So, we have proved that $\mathcal{J}(B) = B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G}$, and we get (i), with $A = B^{\mathfrak{b}}$ and $(b, \mathfrak{a}) = (\mu \circ \beta, \mathfrak{d})$.
We have already obtained that $\tilde{\mathfrak{a}} \circ \mathcal{J} = (\mathcal{J}_{\hat{\alpha} * \beta} id) \mathfrak{b}$, and, for any $x \in \widehat{M'}$, we have
 $\mathcal{J} \circ \mu(x) = 1_{b \otimes_{\alpha} x}$ from which we get, from the definition of the crossed-product, that
 $B = (B^{\mathfrak{b}} \cup \mu(\widehat{M'}))''$. \square

12.4. Corollary. *Let's take the hypothesis and notations of 12.1 and 12.2; then, there exists an isomorphism \mathcal{J} from $B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})$ onto $B \rtimes_{\mathfrak{b}} \mathfrak{G}^c$ such that, for all $x \in B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})$, we have :*

$$\tilde{\mathfrak{b}}\mathcal{J}(x) = (\mathcal{J}_{\mu \circ \beta * \alpha} \mathcal{J}_{\mathfrak{G}})\mathfrak{e}(x)$$

where

$$\mathfrak{e}(x) = (1_{\mu \circ \beta \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})(id_{\mu \circ \beta * \alpha} \varsigma_N)(\mathfrak{d}_{\mu \circ \beta * \alpha} id)(x)(1_{\mu \circ \beta \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})^*$$

and $\mathcal{J}_{\mathfrak{G}}$ had been defined in 3.12(vi).

Proof. Thanks to 12.3, there exist an action $(\mu \circ \beta, \mathfrak{d})$ of \mathfrak{G} on $B^{\mathfrak{b}}$, and an isomorphism \mathcal{J} from B onto $B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G}$ such that $\tilde{\mathfrak{d}} \circ \mathcal{J} = (\mathcal{J}_{\hat{\alpha} * \beta} id) \mathfrak{b}$. Therefore, we have, using 11.6, there exists an isomorphism Θ such that :

$$\begin{aligned} (\mathcal{J}_{\hat{\alpha} * \beta} id)(B \rtimes_{\mathfrak{b}} \mathfrak{G}^c) &= (B^{\mathfrak{b}} \rtimes_{\mathfrak{d}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{d}}} \widehat{\mathfrak{G}}^c \\ &= \Theta(\mathfrak{d}_{\mu \circ \beta * \alpha} id)(B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})) \end{aligned}$$

Let's use again 11.6; for all $x \in B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})$, let's define :

$$\mathfrak{e}(x) = (1_{\mu \circ \beta \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})(id_{\mu \circ \beta * \alpha} \varsigma_N)(\mathfrak{d}_{\mu \circ \beta * \alpha} id)(x)(1_{\mu \circ \beta \otimes_{\alpha} \sigma_{\nu^o}} W \sigma_{\nu^o})^*$$

We have then :

$$\tilde{\mathfrak{d}}\Theta(\mathfrak{d}_{\mu \circ \beta * \alpha} id)(x) = (\Theta(\mathfrak{d}_{\mu \circ \beta * \alpha} id)_{\mu \circ \beta * \alpha} \mathcal{J}_{\mathfrak{G}})\mathfrak{e}(x)$$

Let us define $\mathcal{J} = (\mathcal{J}_{\hat{\alpha} * \beta} id)^{-1} \circ \Theta(\mathfrak{d}_{\mu \circ \beta * \alpha} id)$ which is an isomorphism from $B^{\mathfrak{b}}_{\mu \circ \beta * \alpha} \mathcal{L}(H_{\Phi})$ onto $B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c$; we have then :

$$(\mathcal{J}_{\mu \circ \beta * \alpha} \mathcal{J}_{\mathfrak{G}})\mathfrak{e}(x) = \tilde{\mathfrak{d}}\Theta(\mathfrak{d}_{\mu \circ \beta * \alpha} id)(x) = \tilde{\mathfrak{b}}\mathcal{J}(x)$$

which finishes the proof. \square

12.5. **Corollary.** Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A ; let A_0 a von Neumann subalgebra of $A^{\mathfrak{a}}$, and let us write $D = A \cap A'_0$, and $\mathfrak{d} = \mathfrak{a}|_D$; then :

- (i) (b, \mathfrak{d}) is an action of \mathfrak{G} on D ;
- (ii) $D \rtimes_{\mathfrak{d}} \mathfrak{G} = A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap A'_0 \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$.

Proof. Let us define $B = A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap A'_0 \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$. For $x \in \widehat{M}'$, let us put $\mu(x) = 1 \otimes_{N^o} x$, which belongs to B , and let us write $\mathfrak{b} = \tilde{\mathfrak{a}}|_B$; for all $b \in B$, and $a_0 \in A_0$, we have :

$$\begin{aligned} \mathfrak{b}(b)(a_0 \otimes_{N^o} 1) &= \tilde{\mathfrak{a}}(b) \tilde{\mathfrak{a}}(a_0 \otimes_{N^o} 1) \\ &= \tilde{\mathfrak{a}}(a_0 \otimes_{N^o} 1) \tilde{\mathfrak{a}}(b) \\ &= (a_0 \otimes_{N^o} 1) \mathfrak{b}(b) \end{aligned}$$

from which we get that $\mathfrak{b}(b)$ belongs to $A'_0 \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$, and, therefore, to $B \rtimes_{\hat{\alpha}^* \beta} M'$. So, $(\mu \circ \hat{\alpha}, \mathfrak{b})$ is an action of $\widehat{\mathfrak{G}}^c$ on B , and we can apply 12.3, from which we get that B is the crossed product of $B^{\mathfrak{b}}$ by an action of \mathfrak{G} .

But we have $B^{\mathfrak{b}} = (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}} \cap A'_0 \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$, which is equal to $\mathfrak{a}(A) \cap A'_0 \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$ (11.5(i)), and, therefore, to $\mathfrak{a}(D)$. We then easily finish the proof. \square

13. DUAL WEIGHT; BIDUAL WEIGHT; DEPTH 2 INCLUSION ASSOCIATED TO AN ACTION

Be given an action (b, \mathfrak{a}) of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A and a normal semi-finite faithful weight ψ on A , we define in 13.1 a dual weight $\tilde{\psi}$ on the crossed-product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, using property (B) proved in 11.5(i) and the weight constructed in chapter 10. We obtain a characterization of these weights (13.3), and the GNS construction associated to this weight (13.4). Moreover, we study (13.5) then the unitary U_{ψ} constructed in 10.2. Using then the isomorphism obtained in chapter 11 between the double crossed product and $A \rtimes_{b^* \alpha} \mathcal{L}(H_{\Phi})$, we obtain a characterization of the bidual weight (13.7), which will allow us to construct, using 13.5, Jones' tower associated to the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ (13.8 and 13.9(i)). We prove then that this inclusion is depth 2 (13.9(iv)) and that the operator-valued weight $T_{\tilde{\mathfrak{a}}}$ of this inclusion is regular in the sense of 2.2(13.10).

13.1. **Definition.** Let (b, \mathfrak{a}) be an action of a measured quantum groupoid \mathfrak{G} on a von Neumann algebra A , and let ψ be a normal semi-finite faithful weight on A ; using 9.8, we get that the dual action $\tilde{\mathfrak{a}}$ of $\widehat{\mathfrak{G}}^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is integrable, and, therefore, by definition, the operator-valued weight $T_{\tilde{\mathfrak{a}}}$ from $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ on $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$ is semi-finite. On the other hand, using 11.5(i), we know that $\mathfrak{a}(A) = (A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\tilde{\mathfrak{a}}}$. So, the formula $\tilde{\psi} = \psi \circ \mathfrak{a}^{-1} \circ T_{\tilde{\mathfrak{a}}}$ defines a normal semi-finite faithful weight on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we shall call the dual weight of ψ .

13.2. **Example.** Let \mathcal{G} be a measured groupoid, and \mathfrak{a} an action of \mathcal{G} on a von Neumann algebra, as defined in [Y1], [Y2], [Y3]. We have seen in 9.2 that the crossed-product defined by Yamanouchi is the same one as ours, and in 9.6 that the dual coaction of \mathcal{G} introduced by Yamanouchi is our dual action $\tilde{\mathfrak{a}}$ (of $\widehat{\mathcal{G}}^c$) on the crossed-product $A \rtimes_{\mathfrak{a}} \mathcal{G}$. Starting from a normal semi-finite faithful weight on A , in [Y1], [Y2], Yamanouchi defines

a dual weight on $A \rtimes_{\mathfrak{a}} \mathcal{G}$ ([Y2], 4.10), using an operator-valued weight defined in ([Y2], 3.11), which is equal ([Y2], bottom of the page 665) to our operator-valued weight $T_{\tilde{\mathfrak{a}}}$; therefore, Yamanouchi's dual weight is the same as ours.

13.3. Theorem. *Let \mathcal{G} be a quantum measured groupoid, and (b, \mathfrak{a}) an action of \mathcal{G} on a von Neumann algebra A ; let φ be a normal semi-finite faithful weight on the crossed product $A \rtimes_{\mathfrak{a}} \mathcal{G}$; then, are equivalent :*

- (i) *there exists a normal semi-finite faithful weight ψ on A such that $\varphi = \tilde{\psi}$;*
- (ii) *the weight φ is $\hat{\delta}^{-1}$ -invariant, with respect to the dual action $\tilde{\mathfrak{a}}$ of $\hat{\mathcal{G}}^c$ on the crossed product $A \rtimes_{\mathfrak{a}} \mathcal{G}$, and bears the density property defined in 8.1.*

Proof. Let us suppose (i): then, we have (ii) by a simple application of 10.8(ii).

Let us suppose (ii), and let now θ be a normal semi-finite faithful weight on A , and $\tilde{\theta}$ its dual weight; by 8.11, we obtain that $(D\varphi : D\tilde{\theta})_t$ belongs to $(A \rtimes_{\mathfrak{a}} \mathcal{G})^{\tilde{\mathfrak{a}}}$, which is equal to $\mathfrak{a}(A)$ by 11.5(i). As $\sigma_t^{\tilde{\theta}} \circ \alpha = \alpha \circ \sigma_t^{\theta}$ by definition of the dual weight, there exists a normal semi-finite faithful weight φ on A such that $\mathfrak{a}[(D\psi : D\theta)_t] = (D\varphi : D\tilde{\theta})_t$.

On the other hand, by definition of the dual weights, we have :

$$(D\tilde{\psi} : D\tilde{\theta})_t = \mathfrak{a}[(D\psi : D\theta)_t]$$

From which we deduce that $(D\varphi : D\tilde{\theta})_t = (D\tilde{\psi} : D\tilde{\theta})_t$, which gives the result. \square

13.4. Theorem. *Let \mathcal{G} be a quantum measured groupoid, and (b, \mathfrak{a}) an action of \mathcal{G} on a von Neumann algebra A ; let ψ be a normal faithful semi-finite weight on A , and let $\tilde{\psi}$ be the dual weight constructed in 13.1 on the crossed product $A \rtimes_{\mathfrak{a}} \mathcal{G}$; then, the linear set generated by all the elements $(1_{b \otimes_{\alpha} a})\mathfrak{a}(x)$, for all $x \in \mathfrak{N}_{\psi}$, $a \in \mathfrak{N}_{\hat{\mathcal{G}}^c} \cap \mathfrak{N}_{\hat{T}^c}$, is a core for $\Lambda_{\tilde{\psi}}$, and it is possible to identify the GNS representation of $A \rtimes_{\mathfrak{a}} \mathcal{G}$ associated to the weight $\tilde{\psi}$ with the natural representation on $H_{\psi} \otimes_{\alpha} H_{\Phi}$ by writing :*

$$\Lambda_{\psi}(x) \otimes_{\alpha} \Lambda_{\hat{\mathcal{G}}^c}(a) = \Lambda_{\tilde{\psi}}[(1_{b \otimes_{\alpha} a})\mathfrak{a}(x)]$$

Moreover, using that identification, the linear set generated by the elements of the form $\mathfrak{a}(y^*)(\Lambda_{\psi}(x) \otimes_{\alpha} \Lambda_{\hat{\mathcal{G}}^c}(a))$, for x, y in \mathfrak{N}_{ψ} , and a in $\mathfrak{N}_{\hat{\mathcal{G}}^c} \cap \mathfrak{N}_{\hat{T}^c} \cap \mathfrak{N}_{\hat{\mathcal{G}}^c}^* \cap \mathfrak{N}_{\hat{T}^c}^*$ is a core for $S_{\tilde{\psi}}$, and we have :

$$S_{\tilde{\psi}}\mathfrak{a}(y^*)(\Lambda_{\psi}(x) \otimes_{\alpha} \Lambda_{\hat{\mathcal{G}}^c}(a)) = \mathfrak{a}(x^*)(\Lambda_{\psi}(y) \otimes_{\alpha} \Lambda_{\hat{\mathcal{G}}^c}(a^*))$$

Proof. Thanks to 10.8(i) and (ii), the unitary \tilde{V} constructed in 10.1(ii) is equal to the isometry V constructed in 9.11. So, using this unitary, we can identify $H_{\tilde{\psi}}$ with $H_{\psi} \otimes_{\alpha} H_{\Phi}$, which leads to the first result, using 11.5(i), 9.11 and 10.1(ii). The second result comes then from 10.4. \square

13.5. Proposition. *Let \mathcal{G} be a quantum measured groupoid, and (b, \mathfrak{a}) an action of \mathcal{G} on a von Neumann algebra A ; let ψ be a normal faithful semi-finite weight on A , and let $\tilde{\psi}$ be the dual weight constructed in 13.1 on the crossed product $A \rtimes_{\mathfrak{a}} \mathcal{G}$; let us identify $H_{\tilde{\psi}}$ with $H_{\psi} \otimes_{\alpha} H_{\Phi}$ as in 13.4. Then, the unitary $U_{\psi}^{\mathfrak{a}}$ from $H_{\psi} \otimes_{\beta} H_{\Phi}$ onto $H_{\psi} \otimes_{\alpha} H_{\Phi}$ defined by :*

$$U_{\psi}^{\mathfrak{a}} = J_{\tilde{\psi}}(J_{\psi} \otimes_{\beta} J_{\Phi})$$

satisfies :

$$U_\psi^{\mathfrak{a}}(J_\psi \underset{N}{b} \otimes_\alpha J_\Phi) = (U_\psi^{\mathfrak{a}})^*(J_\psi \underset{N^o}{a} \otimes_\beta J_\Phi)$$

and we have :

(i) for all $y \in A$:

$$\mathfrak{a}(y) = U_\psi^{\mathfrak{a}}(y \underset{N^o}{a} \otimes_\beta 1)(U_\psi^{\mathfrak{a}})^*$$

(ii) for all $b \in M$:

$$(1 \underset{N}{b} \otimes_\alpha J_\Phi b J_\Phi) U_\psi^{\mathfrak{a}} = U_\psi^{\mathfrak{a}}(1 \underset{N^o}{a} \otimes_\beta J_\Phi b J_\Phi)$$

(iii) for all $n \in N$:

$$U_\psi^{\mathfrak{a}}(b(n) \underset{N^o}{a} \otimes_\beta 1) = (1 \underset{N}{b} \otimes_\alpha \beta(n)) U_\psi^{\mathfrak{a}}$$

$$U_\psi^{\mathfrak{a}}(1 \underset{N^o}{a} \otimes_\beta \alpha(n)) = (a(n) \underset{N}{b} \otimes_\alpha 1) U_\psi^{\mathfrak{a}}$$

Proof. Using again 10.8(i) and 11.5(i), the first result is given by 10.2(iii); (i) is given by 10.2(i), (ii) by 10.5. Applying (i) to $y = b(n)$, we get the first result of (iii); we get then the second result of (iii) by using $U_\psi^{\mathfrak{a}}(J_\psi \underset{N}{b} \otimes_\alpha J_\Phi) = (U_\psi^{\mathfrak{a}})^*(J_\psi \underset{N^o}{a} \otimes_\beta J_\Phi)$. \square

13.6. Lemma. Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A ; let $(1 \underset{N}{b} \otimes_\alpha \hat{\beta}, \underline{\mathfrak{a}})$ be the action of \mathfrak{G} on $A \underset{N}{b} *_\alpha \mathcal{L}(H)$ constructed in 11.6. Then, for all a in $\mathfrak{N}_{\widehat{\Phi}^c}$ and X in A^+ , we have :

$$T_{\underline{\mathfrak{a}}}(\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha} X (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^*) = (1 \underset{N}{b} \otimes_\alpha a) \mathfrak{a}(X) (1 \underset{N}{b} \otimes_\alpha a^*)$$

Proof. Let us first remark that $\Lambda_{\widehat{\Phi}^c}(a) = J_{\widehat{\Phi}} \Lambda_{\widehat{\Phi}}(J_{\widehat{\Phi}} a J_{\widehat{\Phi}})$ which belongs to $D({}_\alpha H_\Phi, \nu)$, by 2.2; on the other hand, let us suppose that A is acting on an Hilbert space \mathfrak{H} ; we verify straightforwardly that $\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha} X (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^*$ commutes with all $Y \underset{N}{b} \otimes_\alpha 1$, with $Y \in A'$, and, therefore, belongs to $(A \underset{N}{b} *_\alpha \mathcal{L}(H))^+$.

Let η in $D(\mathfrak{H}_b, \nu^o)$; for all ξ in $D((H_\Phi)_{\hat{\beta}}, \nu^o)$, we have, using 11.6 :

$$\begin{aligned} &< T_{\underline{\mathfrak{a}}}(\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha} X (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^*), \omega_{\eta \underset{N}{b} \otimes_\alpha \xi} > = \\ &\Phi[(\omega_{\eta \underset{N}{b} \otimes_\alpha id})(1 \underset{N}{b} \otimes_\alpha (\rho_\xi^{\alpha, \hat{\beta}})^* W \rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{\beta, \alpha}) \mathfrak{a}(X) (1 \underset{N}{b} \otimes_\alpha (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^* W^* \rho_\xi^{\alpha, \hat{\beta}})] \end{aligned}$$

We know that $(\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^* W^* \rho_\xi^{\alpha, \hat{\beta}} = (id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)$ belongs to \mathfrak{N}_Φ , thanks to 9.11 applied to $\widehat{\mathfrak{G}}$, and that :

$$\Lambda_\Phi((id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)) = a_{\widehat{\Phi}}(\omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)}) = a^* \xi$$

and, therefore, we get that $< T_{\underline{\mathfrak{a}}}(\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha} X (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^*), \omega_{\eta \underset{N}{b} \otimes_\alpha \xi} >$ is equal to :

$$\begin{aligned} &\Phi((id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)^* (\omega_{\eta \underset{N}{b} \otimes_\alpha id})(X) (id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)) = \\ &= ((\omega_{\eta \underset{N}{b} \otimes_\alpha id})(X) \Lambda_\Phi((id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)) | \Lambda_\Phi(id * \omega_{\xi, \Lambda_{\widehat{\Phi}^c}(a)})(W^*)) \\ &= ((\omega_{\eta \underset{N}{b} \otimes_\alpha id})(X) a^* \xi | a^* \xi) \end{aligned}$$

and, finally, we get that :

$$< T_{\underline{\mathfrak{a}}}(\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha} X (\rho_{\Lambda_{\widehat{\Phi}^c}(a)}^{b,\alpha})^*), \omega_{\eta \underset{N}{b} \otimes_\alpha \xi} > = ((1 \underset{N}{b} \otimes_\alpha a) \mathfrak{a}(X) (1 \underset{N}{b} \otimes_\alpha a^*) (\eta \underset{N}{b} \otimes_\alpha \xi) | \eta \underset{N}{b} \otimes_\alpha \xi)$$

from which we get the result. \square

13.7. Theorem. *Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A ; let ψ be a normal semi-finite faithful weight on A ; let $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ be the crossed-product, as defined in 9.1, and $\tilde{\psi}$ the dual weight, as defined in 13.1; let us consider the dual action, as defined in 9.4, the bicrossed-product and the bidual weight $\tilde{\tilde{\psi}}$. We have defined in 11.6 an isomorphism Θ such that :*

$$\Theta(\mathfrak{a}_{b^*_{\alpha}} id)(A_{b^*_{\alpha}} \mathcal{L}(H_{\Phi})) = (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^o$$

We have then :

$$\frac{d\tilde{\psi} \circ \Theta(\mathfrak{a}_{b^*_{\alpha}} id)}{d\psi^o} = \Delta_{\tilde{\psi}}$$

Proof. Let us remark that $\tilde{\psi} \circ \Theta(\mathfrak{a}_{b^*_{\alpha}} id)$ is a normal semi-finite weight on $A_{b^*_{\alpha}} \mathcal{L}(H_{\Phi})$, whose commutant is $A'_{b \otimes_{\alpha} 1}$, which is isomorphic to A' , and, therefore to A^o . Therefore, the spatial derivative has a meaning. Let us write it h for simplification. By definition of this spatial derivative, for all Ξ in $D(H_{\psi} b \otimes_{\alpha} H_{\Phi}, \psi^o)$, Ξ belongs to $D(h^{1/2})$ if and only if $\theta^{\psi^o}(\Xi, \Xi)$ belongs to $\mathfrak{M}_{\tilde{\psi} \circ \Theta(\mathfrak{a}_{b^*_{\alpha}} id)}^+$, and we have then :

$$\|h^{1/2}\Xi\|^2 = \tilde{\psi} \circ \Theta(\mathfrak{a}_{b^*_{\alpha}} id)(\theta^{\psi^o}(\Xi, \Xi))$$

which, using 11.6, gives that :

$$\|h^{1/2}\Xi\|^2 = \tilde{\psi} \circ T_b(\theta^{\psi^o}(\Xi, \Xi))$$

Let now x, y in \mathfrak{N}_{ψ} , and a in $\mathfrak{N}_{\hat{T}^c} \cap \mathfrak{N}_{\hat{T}^c}^* \cap \mathfrak{N}_{\hat{\Phi}} \cap \mathfrak{N}_{\hat{\Phi}}^*$; we have, for all $z \in \mathfrak{N}_{\psi}$:

$$\begin{aligned} (J_{\psi} z^* J_{\psi} b \otimes_{\alpha} 1)(\mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)})) &= \mathfrak{a}(x^*)(J_{\psi} z^* J_{\psi} \Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)}) \\ &= \mathfrak{a}(x^*)(y J_{\psi} \Lambda_{\psi}(z) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)}) \\ &= \mathfrak{a}(x^*) \rho_{\Lambda_{\hat{\Phi}^c(a)}}^{b, \alpha} y J_{\psi} \Lambda_{\psi}(z) \end{aligned}$$

from which we get that $\mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)})$ belongs to $D(H_{\psi} b \otimes_{\alpha} H_{\Phi}, \psi^o)$, and that :

$$R^{\psi^o}(\mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)})) = \mathfrak{a}(x^*) \rho_{\Lambda_{\hat{\Phi}^c(a)}}^{b, \alpha} y$$

Therefore, for x, y in \mathfrak{N}_{ψ} , a in $\mathfrak{N}_{\hat{T}^c} \cap \mathfrak{N}_{\hat{T}^c}^* \cap \mathfrak{N}_{\hat{\Phi}} \cap \mathfrak{N}_{\hat{\Phi}}^*$, we get that :

$$\theta^{\psi^o}(\mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)}), \mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)})) = \mathfrak{a}(x^*) \rho_{\Lambda_{\hat{\Phi}^c(a)}}^{b, \alpha} y y^* (\rho_{\Lambda_{\hat{\Phi}^c(a)}}^{b, \alpha})^* \mathfrak{a}(x)$$

which, thanks to 13.6, belongs to $\mathfrak{M}_{T_{\underline{a}}}^+$, and we have :

$$\begin{aligned} T_{\underline{a}}(\theta^{\psi^o}(\mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)}), \mathfrak{a}(x^*)(\Lambda_{\psi}(y) b \otimes_{\alpha} \Lambda_{\hat{\Phi}^c(a)}))) &= \\ \mathfrak{a}(x^*)(1 b \otimes_{\alpha} a) \mathfrak{a}(y y^*)(1 b \otimes_{\alpha} a^*) \mathfrak{a}(x) \end{aligned}$$

Using now 13.4, we get that this last operator belongs to \mathfrak{M}_ψ^+ , and, therefore, that $\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})$ belongs to $D(h^{1/2})$. And, by 13.4, we get that $\mathfrak{a}(y^*)(1 {}_b \otimes_\alpha a^*)\mathfrak{a}(x)$ belongs to $\mathfrak{N}_{\tilde{\psi}} \cap \mathfrak{N}_\psi^*$, and that :

$$\tilde{\psi} \circ T_{\underline{a}}(\theta^{\psi^o}(\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)}), \mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})))$$

is equal to :

$$\begin{aligned} \|\Lambda_{\tilde{\psi}}(\mathfrak{a}(y^*)(1 {}_b \otimes_\alpha a^*)\mathfrak{a}(x))\|^2 &= \|S_{\tilde{\psi}}\Lambda_{\tilde{\psi}}[\mathfrak{a}(x^*)(1 {}_b \otimes_\alpha a)\mathfrak{a}(y)]\|^2 \\ &= \|\Delta_{\tilde{\psi}}^{1/2}\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})\|^2 \end{aligned}$$

Therefore, we get that :

$$\|h^{1/2}\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})\| = \|\Delta_{\tilde{\psi}}^{1/2}\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})\|$$

and, using again 13.4, as the vectors $\mathfrak{a}(x^*)(\Lambda_\psi(y) {}_b \otimes_\alpha \Lambda_{\widehat{\Phi}^c(a)})$ are a core for $\Delta_{\tilde{\psi}}^{1/2}$, we have $\|h^{1/2}\xi\| = \|\Delta_{\tilde{\psi}}^{1/2}\xi\|$ for all ξ in $D(h^{1/2})$, from which we get that $\Delta_{\tilde{\psi}} \subset h$, and, as they are self-adjoint operators, we get the result. \square

13.8. Theorem. *Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A , $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ its crossed product; let $(1 {}_b \otimes_\alpha \hat{\beta}, \underline{a})$ be the action of \mathfrak{G} on $A {}_b \otimes_\alpha \mathcal{L}(H_\Phi)$ introduced in 11.6; then, the inclusion :*

$$\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G} \subset A {}_b \otimes_\alpha \mathcal{L}(H_\Phi)$$

is standard, and the operator-valued weight $T_{\underline{a}}$ from $A {}_b \otimes_\alpha \mathcal{L}(H_\Phi)$ to $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is the basic construction made from the operator-valued weight $T_{\hat{\mathfrak{a}}}$ from $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ to $\mathfrak{a}(A)$.

Proof. Let ψ be a normal semi-finite faithful weight on A , and $\tilde{\psi}$ its dual weight on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$; let us represent the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ on the Hilbert space $H_{\tilde{\psi}}$, which had been identified (13.4) with $H_\psi {}_b \otimes_\alpha H_\Phi$, equipped with the natural representation of the inclusion. We have then, using 13.5 :

$$\begin{aligned} J_{\tilde{\psi}}\mathfrak{a}(A)J_{\tilde{\psi}} &= J_{\tilde{\psi}}U_\psi^{\mathfrak{a}}(A {}_a \otimes_\beta 1)(U_\psi^{\mathfrak{a}})^*J_{\tilde{\psi}} \\ &= (J_\psi {}_a \otimes_\beta J_{\widehat{\Phi}})(A {}_a \otimes_\beta 1)(J_\psi {}_b \otimes_\alpha J_{\widehat{\Phi}}) \\ &= A' {}_b \otimes_\alpha 1 \end{aligned}$$

and, therefore, we get :

$$J_{\tilde{\psi}}\mathfrak{a}(A)'J_{\tilde{\psi}} = (A' {}_b \otimes_\alpha 1)' = A {}_b \otimes_\alpha \mathcal{L}(H_\Phi)$$

which gives that $A {}_b \otimes_\alpha \mathcal{L}(H_\Phi)$ is the standard construction made from the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$.

Let T be the standard construction made from the operator-valued weight $T_{\hat{\mathfrak{a}}}$. By definition (2.2), we have :

$$\frac{d\tilde{\psi} \circ T}{d\psi^o} = \Delta_{\tilde{\psi}}$$

which, thanks to 13.7, gives that $\tilde{\psi} \circ T = \tilde{\psi} \circ \Theta(\mathfrak{a}_{b^* \alpha} id)$, and, using 11.6, we have :

$$\tilde{\psi} \circ T = \tilde{\psi} \circ T_{\underline{\mathfrak{a}}}$$

from which we get that $T = T_{\underline{\mathfrak{a}}}$, which finishes the proof. \square

13.9. Theorem. *Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A , $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ its crossed product; then :*

(i) *the Jones' tower associated to the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is :*

$$\mathfrak{a}(A)_{\hat{\alpha} \otimes \beta} 1 \subset \tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G})_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o}$$

(ii) *moreover, the operator-valued weight $T_{(\tilde{\mathfrak{a}})}$ from $(A \rtimes_{\mathfrak{a}} \mathfrak{G})_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})$ to $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$*

is the operator-valued weight constructed by successive basic constructions in this tower.

(iii) *the derived tower is :*

$$\begin{aligned} \tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap \mathfrak{a}(A)') &\subset (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c \cap \mathfrak{a}(A)'_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o} \\ &\subset [A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap \mathfrak{a}(A)']_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o} \end{aligned}$$

(iv) *let us write $B = A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap \mathfrak{a}(A)'$, and $\mathfrak{b} = \tilde{\mathfrak{a}}|_B$. Then $(1_{b^* \otimes \alpha} \hat{\alpha}, \mathfrak{b})$ is an action of $\widehat{\mathfrak{G}}^c$ on B , and we have :*

$$B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c = (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c \cap \mathfrak{a}(A)'_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o}$$

(v) *the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is depth 2.*

Proof. We have got in 13.8 that the inclusion :

$$\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G} \subset A_{b^* \alpha} \mathcal{L}(H_{\Phi})_N$$

is standard. Let's apply the isomorphism $\Theta(\mathfrak{a}_{b^* \alpha} id)$ from $A_{b^* \alpha} \mathcal{L}(H_{\Phi})_N$ onto $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$ defined in 11.6; we then get the inclusion :

$$\mathfrak{a}(A)_{\hat{\alpha} \otimes \beta} 1 \subset \tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$$

of von Neumann algebras on $H_{\psi} \otimes_{\nu} H_{\Phi} \otimes_{\nu^o} H_{\Phi}$; let's apply now 13.8 to the inclusion :

$$\tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \subset (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\tilde{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$$

We get that the basic construction made from this inclusion gives the von Neumann algebra $(A \rtimes_{\mathfrak{a}} \mathfrak{G})_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o}$, which gives (i).

By 11.6 again, we see that the isomorphism $\Theta(\mathfrak{a}_{b^* \alpha} id)$ sends the operator-valued weight $T_{\underline{\mathfrak{a}}}$ on $T_{\underline{\mathfrak{a}}}^*$. So, applying again 13.8 to this operator-valued weight, we get (ii).

Now, (iii) is just a corollary from (i), and (iv) is just an application of 12.5. So, we get that the derived tower obtained in (iii) can be written as :

$$\mathfrak{b}(B) \subset B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c \subset B_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi})_{N^o}$$

which is standard, by 13.8 applied to the action \mathfrak{b} ; which finishes the proof. \square

13.10. **Theorem.** Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A , $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ its crossed product, $\underline{\mathfrak{a}}$ the action of \mathfrak{G} on $A_{b^* \alpha} \mathcal{L}(H_{\Phi})$ introduced in 11.6. Then :

(i) for any $x \in M'^+$, $1_{b \otimes_{\alpha} N} x$ belongs to $A_{b^* \alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'$, and we have :

$$T_{\underline{\mathfrak{a}}}(1_{b \otimes_{\alpha} N} x) = 1_{b \otimes_{\alpha} N} T^{co}(x)$$

(ii) the restriction of $T_{\underline{\mathfrak{a}}}$ to $A_{b^* \alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'$ is semi-finite;

(iii) the operator-valued weight $T_{\underline{\mathfrak{a}}}$ from $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ to $\mathfrak{a}(A)$ is regular in the sense of 2.2.

Proof. Let $x \in M'$; we have, using 3.11 :

$$\begin{aligned} W(1_{\beta \otimes_{\alpha} N} x) W^* &= (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) W^* (1_{\alpha \otimes_{\beta} N^o} J_{\Phi} x J_{\Phi}) W (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \\ &= (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \Gamma(J_{\Phi} x J_{\Phi}) (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \end{aligned}$$

and, therefore, using 11.6 :

$$\underline{\mathfrak{a}}(1_{b \otimes_{\alpha} N} x) = (id_{b^* \alpha} \varsigma_N) (1_{b \otimes_{\alpha} N} [(J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi}) \Gamma(J_{\Phi} x J_{\Phi}) (J_{\widehat{\Phi}} \beta \otimes_{\alpha} J_{\Phi})])$$

from which we get, for x positive :

$$\begin{aligned} T_{\underline{\mathfrak{a}}}(1_{b \otimes_{\alpha} N} x) &= 1_{b \otimes_{\alpha} N} J_{\Phi} (\Phi \circ R_{\beta^* \alpha} id) \Gamma(J_{\Phi} x J_{\Phi}) J_{\Phi} \\ &= 1_{b \otimes_{\alpha} N} J_{\Phi} R T R (J_{\Phi} x J_{\Phi}) J_{\Phi} \\ &= 1_{b \otimes_{\alpha} N} T^{co}(x) \end{aligned}$$

which is (i). Then, (ii) is just a corollary of (i).

Let us apply now (ii) to the action $\tilde{\mathfrak{a}}$, we get that the restriction of $T_{(\tilde{\mathfrak{a}})}$ to the von Neumann algebra $(A \rtimes_{\mathfrak{a}} \mathfrak{G})_{\hat{\alpha}^* \beta} \mathcal{L}(H_{\Phi}) \cap \tilde{\mathfrak{a}}(A \rtimes_{\mathfrak{a}} \mathfrak{G})'$ is semi-finite, which, with (ii), gives (iii). \square

13.11. **Remark.** The fact that the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is depth 2 (13.9(v)) had been obtained in ([V2], 5.10) for actions of locally compact quantum groups in a somehow different way.

14. THE MEASURED QUANTUM GROUPOID ASSOCIATED TO AN ACTION

In this section, we apply the results of ([EV], [E4]), recalled in 3.15, 6.5 and 9.3, to the depth 2 inclusion (13.9) $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$, where (b, \mathfrak{a}) is an action of \mathfrak{G} on a von Neumann algebra A . From such data, we therefore obtain a new quantum measured groupoid $\mathfrak{G}(\mathfrak{a})$ (14.2), and we show that the dual action $\tilde{\mathfrak{a}}$ of $\widehat{\mathfrak{G}}^c$ can be considered as an action of $\mathfrak{G}(\tilde{\mathfrak{a}})$ (14.5). As the underlying von Neumann algebra of $\mathfrak{G}(\mathfrak{a})$ is a crossed-product of the relative commutant $A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap \mathfrak{a}(A)'$ by the restriction of the action $\tilde{\mathfrak{a}}$ of $\widehat{\mathfrak{G}}^c$, we show that the initial measured quantum groupoid can be naturally sent into $\mathfrak{G}(\mathfrak{a})$ (14.7), and, moreover, that, in the particular case when we are starting with an outer action of a locally compact quantum group, we recover the initial locally compact quantum group (14.9).

14.1. Theorem. Let \mathfrak{G} be a measured quantum groupoid, and (b, \mathfrak{a}) an action of \mathfrak{G} on a von Neumann algebra A . Let ψ be a normal faithful semi-finite weight on A , and $\tilde{\psi}$ be the dual weight on the crossed product $A \rtimes_{\mathfrak{a}} \mathfrak{G}$. Let us write B for $A \rtimes_{\mathfrak{a}} \mathfrak{G} \cap \mathfrak{a}(A)'$, \mathfrak{b} for $\tilde{\mathfrak{a}}|_B$, and \tilde{M} for $A_{b^* \alpha_N} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'$; then :

(i) let us write id for the inclusion of B into \tilde{M} , and j for the anti- $*$ -homomorphism $j(x) = J_{\tilde{\psi}} x^* J_{\tilde{\psi}}$, which sends B into $A_{b^* \alpha_N} \mathcal{L}(H_{\Phi}) \cap (A \rtimes_{\mathfrak{a}} \mathfrak{G})'$ (and, therefore, into \tilde{M}).

Then, there exists $\tilde{\Gamma}$, such that :

$$(B, \tilde{M}, id, j, \tilde{\Gamma})$$

is a Hopf-bimodule.

(ii) let us write also $j(x) = J_{\tilde{\psi}} x^* J_{\tilde{\psi}}$ the $*$ -anti-automorphism of \tilde{M} ; then j is a co-inverse for the coproduct constructed in (i).

(iii) let us write \tilde{T} for the restriction of $T_{\mathfrak{a}}$ to \tilde{M} ; then, \tilde{T} is a normal, semi-finite, faithful operator-valued weight from \tilde{M} onto B , which is left-invariant with respect to the coproduct $\tilde{\Gamma}$.

(iv) the isomorphism $\Theta(\mathfrak{a}_{b^* \alpha_N} id)$ sends \tilde{M} onto $B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c$ and B onto $\mathfrak{b}(B)$. Let us denote j_1 the $*$ -anti-automorphism of $B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c$ obtained by transporting j via this isomorphism, and $\tilde{\Gamma}_1$ the coproduct obtained the same way. Then :

$$(B, B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c, \mathfrak{b}, j_1 \circ \mathfrak{b}, \tilde{\Gamma}_1)$$

is a Hopf-bimodule, isomorphic to the one obtained in (i), j_1 is a co-inverse, and $T_{\mathfrak{b}}$ is a left-invariant operator-valued weight.

Proof. We had got in 13.9 that the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathfrak{G}$ is depth 2, and, in 13.10, that the operator-valued weight $T_{\mathfrak{a}}$ is regular. So, we can apply 3.15(i) and (ii). As, by 13.8, the basic construction made from this inclusion gives the algebra $A_{b^* \alpha_N} \mathcal{L}(H_{\Phi})$, and the beginning of the derived tower is $B \subset \tilde{M}$, we get (i), (ii) and (iii). Using 13.9(iv), we get the beginning of (iv); moreover, using 11.2, we get that this isomorphism sends the restriction of $T_{\mathfrak{a}}$ on the restriction of $T_{\tilde{\mathfrak{a}}}$, which is $T_{\mathfrak{b}}$. \square

14.2. Theorem. Let us take the notations of 14.1. Let us suppose that there exists a normal semi-finite faithful weight χ on B , invariant under the modular automorphism group $\sigma_t^{T_{\tilde{\mathfrak{a}}}}$; then :

$$(B, \tilde{M}, id, j, \tilde{\Gamma}, \tilde{T}, j \circ \tilde{T} \circ j, \chi) = (B, B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c, \mathfrak{b}, j_1 \circ \mathfrak{b}, \tilde{\Gamma}_1, T_{\mathfrak{b}}, j_1 \circ T_{\mathfrak{b}} \circ j_1, \chi)$$

is a measured quantum groupoid, we shall denote by $\mathfrak{G}(\mathfrak{a})$. Moreover, the dual action $\tilde{\mathfrak{a}}$ satisfies the same hypothesis, and $\widehat{\mathfrak{G}(\tilde{\mathfrak{a}})} = \widehat{\mathfrak{G}(\mathfrak{a})}^o$.

Proof. This is just 3.15 and 13.9. \square

14.3. Theorem. Let us take the notations of 14.1 and 14.2; there exists an action $(id, \bar{\mathfrak{a}})$ of $\mathfrak{G}(\tilde{\mathfrak{a}})$ on $A \rtimes_{\mathfrak{a}} \mathfrak{G}$ (where id means the inclusion of B into $A \rtimes_{\mathfrak{a}} \mathfrak{G}$, which is a anti- $*$ -homomorphism of the basis B^o of $\mathfrak{G}(\tilde{\mathfrak{a}})$), such that $(A \rtimes_{\mathfrak{a}} \mathfrak{G})^{\bar{\mathfrak{a}}} = \mathfrak{a}(A)$, and such that the crossed-product $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \mathfrak{G}(\tilde{\mathfrak{a}})$ is isomorphic to $A_{b^* \alpha_N} \mathcal{L}(H_{\Phi})$. More precisely, there exists an isomorphism $I_{\bar{\mathfrak{a}}}$ from $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \mathfrak{G}(\tilde{\mathfrak{a}})$ onto $A_{b^* \alpha_N} \mathcal{L}(H_{\Phi})$, such that :

(i) for any $x \in A \rtimes_{\mathfrak{a}} \mathfrak{G}$, we have $I_{\bar{\mathfrak{a}}}(\bar{\mathfrak{a}}(x)) = x$;

(ii) for any $y \in A_{b^*_{\alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'} \cap \mathfrak{a}(A)'$, we have $I_{\bar{\mathfrak{a}}}(1_{id \otimes_{\hat{j} \circ \mathfrak{b}} y}) = y$, where $\hat{j}(z) = J_{\tilde{\chi}} z^* J_{\tilde{\chi}}$, for all $z \in B \rtimes_{\mathfrak{b}} \widehat{\mathfrak{G}}^c$.

Proof. This is just 9.3 applied to 13.9 and 14.2. \square

14.4. **Theorem.** Let us take the notations of 14.1, 14.2 and 14.3; then, the application defined, for any $x \in \mathfrak{N}_{\tilde{\psi}}$, $y \in \mathfrak{N}_{\chi}$, $b \in \mathfrak{N}_{\Phi^{oc}} \cap \mathfrak{N}_{T^{oc}}$ by :

$$\mathcal{U}[\Lambda_{\tilde{\psi}}(x)_{id \otimes_{\chi^o} \hat{j} \circ \mathfrak{b}} (\Lambda_{\chi}(y)_{\hat{\alpha} \otimes_{\nu^o} \beta} \Lambda_{\Phi^{oc}}(b))] = \Lambda_{\tilde{\psi}}(yx)_{\hat{\alpha} \otimes_{\nu^o} \beta} \Lambda_{\Phi^{oc}}(b)$$

is a unitary from $H_{\tilde{\psi}}_{id \otimes_{\chi^o} \hat{j} \circ \mathfrak{b}} H_{\tilde{\chi}}$ onto $H_{\tilde{\psi}}_{\hat{\alpha} \otimes_{\nu^o} \beta} H_{\Phi}$, such that, for all $X \in (A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$, we have :

$$\mathcal{U}X\mathcal{U}^* = \Theta \circ (\mathfrak{a}_{b^*_{\alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'} id) I_{\bar{\mathfrak{a}}}(X)$$

and, in particular, for all $Y \in A \rtimes_{\mathfrak{a}} \mathfrak{G}$, $c \in M'$, we have :

$$\mathcal{U}\bar{\mathfrak{a}}(Y)\mathcal{U}^* = \tilde{\mathfrak{a}}(Y)$$

$$\mathcal{U}(1_{id \otimes_{B^o} \hat{j} \circ \mathfrak{b}} (1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} c))\mathcal{U}^* = 1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} c$$

Proof. Thanks to 14.3 and 11.6, we get that $\Theta \circ (\mathfrak{a}_{b^*_{\alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'} id) I_{\bar{\mathfrak{a}}}$ is an isomorphism between $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \mathfrak{G}(\tilde{\mathfrak{a}})$ and $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \widehat{\mathfrak{G}}^c$, which verifies, for all Y and c :

$$\Theta \circ (\mathfrak{a}_{b^*_{\alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'} id) I_{\bar{\mathfrak{a}}}(\bar{\mathfrak{a}}(Y)) = \tilde{\mathfrak{a}}(Y)$$

$$\Theta \circ (\mathfrak{a}_{b^*_{\alpha} \mathcal{L}(H_{\Phi}) \cap \mathfrak{a}(A)'} id) I_{\bar{\mathfrak{a}}}(1_{id \otimes_{B^o} \hat{j} \circ \mathfrak{b}} (1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} c)) = 1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} c$$

On the other hand, let us put, for all $n \in N$, $\tilde{b}(n) = J_{\tilde{\chi}} \hat{\alpha}(n^*) J_{\tilde{\chi}}$. We have, by definition of $U_{\chi}^{\mathfrak{b}}$, for all $y \in B$:

$$\begin{aligned} \hat{j}(\mathfrak{b}(y)) &= J_{\tilde{\chi}} \mathfrak{b}(y^*) J_{\tilde{\chi}} \\ &= J_{\tilde{\chi}} U_{\chi}^{\mathfrak{b}}(y^*_{\tilde{b} \otimes_{\hat{\alpha}} 1})(U_{\chi}^{\mathfrak{b}})^* J_{\tilde{\chi}} \\ &= J_{\chi} y^* J_{\chi} \hat{\alpha} \otimes_{\nu^o} \beta 1 \\ &= s(y)_{\hat{\alpha} \otimes_{\nu^o} \beta} 1 \end{aligned}$$

where $s(y) = J_{\chi} y^* J_{\chi}$. Therefore, the identification of $\Lambda_{\tilde{\psi}}(x)_{id \otimes_{\chi^o} \hat{j} \circ \mathfrak{b}} \Lambda_{\chi}(y)$ with $\Lambda_{\tilde{\psi}}(yx)_{\hat{\alpha} \otimes_{\nu^o} \beta} \Lambda_{\Phi^{oc}}(b)$ (2.3) gives the definition of \mathcal{U} .

Let us write $\widetilde{(\psi)}$ for the dual weight of $\tilde{\psi}$ on $(A \rtimes_{\mathfrak{a}} \mathfrak{G}) \rtimes_{\bar{\mathfrak{a}}} \mathfrak{G}(\tilde{\mathfrak{a}})$. Using 13.4 applied to the weight $\tilde{\chi}$, then to the weight $\widetilde{(\psi)}$, we get that :

$$\begin{aligned} \Lambda_{\tilde{\psi}}(x)_{id \otimes_{\chi^o} \hat{j} \circ \mathfrak{b}} (\Lambda_{\chi}(y)_{\hat{\alpha} \otimes_{\nu^o} \beta} \Lambda_{\Phi^{oc}}(b)) &= \Lambda_{\tilde{\psi}}(x)_{id \otimes_{\chi^o} \hat{j} \circ \mathfrak{b}} \Lambda_{\tilde{\chi}}((1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} b) \mathfrak{b}(y)) \\ &= \Lambda_{\widetilde{(\psi)}}[(1_{id \otimes_{B^o} \hat{j} \circ \mathfrak{b}} (1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} b) \mathfrak{b}(y)) \bar{\mathfrak{a}}(x)] \end{aligned}$$

As we have $\bar{\mathfrak{a}}(y) = 1_{id \otimes_{B^o} \hat{j} \circ \mathfrak{b}} \mathfrak{b}(y)$, we finally get, using again 13.4 applied to the bidual weight $\tilde{\psi}$:

$$\mathcal{U} \Lambda_{\widetilde{(\psi)}}[(1_{id \otimes_{B^o} \hat{j} \circ \mathfrak{b}} (1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} b) \bar{\mathfrak{a}}(yx))] = \Lambda_{\tilde{\psi}}((1_{b \otimes_{\alpha} N} 1_{\hat{\alpha} \otimes_{\beta} N^o} b) \tilde{\mathfrak{a}}(yx))$$

which we can write, using the isomorphism $J = \Theta \circ (\mathbf{a}_{b^* \alpha} id) I_{\bar{\mathbf{a}}} :$

$$\mathcal{U} \Lambda_{(\tilde{\psi})}[(1_{id \otimes_{B^o} \hat{b}} (1_{b \otimes_{\alpha} 1_{\hat{\alpha} \otimes_{\beta} b}})) \bar{\mathbf{a}}(yx)] = \Lambda_{\tilde{\psi}}(J[(1_{id \otimes_{B^o} \hat{b}} (1_{b \otimes_{\alpha} 1_{\hat{\alpha} \otimes_{\beta} b}})) \bar{\mathbf{a}}(yx)])$$

Using 10.8(ii) applied to $(\tilde{\psi})$ and to $\tilde{\psi}$, we see that these elements are a core for, respectively, $\Lambda_{(\tilde{\psi})}$ and $\Lambda_{\tilde{\psi}}$, and we finally get that $\mathcal{U} \Lambda_{(\tilde{\psi})}(Z) = \Lambda_{\tilde{\psi}}(J(Z))$, for all $Z \in \mathfrak{N}_{(\tilde{\psi})}$. From which one gets that :

$$\mathcal{U} X \mathcal{U}^* = \Theta \circ (\mathbf{a}_{b^* \alpha} id) I_{\bar{\mathbf{a}}}(X)$$

which finishes the proof. \square

14.5. Corollary. *Let us take the notations of 14.1, 14.2 and 14.3. If we make the identification of $H_{\tilde{\psi}} id \otimes_{B^o} H_{\tilde{\chi}}$ with $H_{\tilde{\chi}}$ by writing, for any $x \in \mathfrak{N}_{\tilde{\psi}}$, $y \in \mathfrak{N}_{\tilde{\chi}}$, $b \in \mathfrak{N}_{\Phi^{oc}} \cap \mathfrak{N}_{T^{oc}}$:*

$$\Lambda_{\tilde{\psi}}(x) id \otimes_{B^o} (\Lambda_{\tilde{\chi}}(y) \hat{\alpha} \otimes_{\beta} \Lambda_{\Phi^{oc}}(b)) = \Lambda_{\tilde{\psi}}(yx) \hat{\alpha} \otimes_{\beta} \Lambda_{\Phi^{oc}}(b)$$

we then identify $\tilde{\mathbf{a}}$ with $\bar{\mathbf{a}}$, and, therefore, $(id, \tilde{\mathbf{a}})$ (where id is the inclusion of B^o into $A \rtimes_{\mathbf{a}} \mathfrak{G}$) is an action of $\mathfrak{G}(\tilde{\mathbf{a}})$ on $A \rtimes_{\mathbf{a}} \mathfrak{G}$.

Proof. Clear by 14.4. \square

14.6. Corollary. *Let us use the notations of 14.1, 14.2, 14.3. Then, the dual action $(j_1 \circ b, \tilde{\mathbf{a}})$ of $\widehat{\mathfrak{G}(\tilde{\mathbf{a}})}^c = \mathfrak{G}(\mathbf{a})$ on $(A \rtimes_{\mathbf{a}} \mathfrak{G}) \rtimes_{\tilde{\mathbf{a}}} \widehat{\mathfrak{G}}^c$ can be, as well, be identified with the action $(1_{b \otimes_{\alpha} 1_{\hat{\alpha} \otimes_{\beta} \hat{\beta}}}, \tilde{\mathbf{a}})$ of \mathfrak{G}^{oc} , and the action $(1_{b \otimes_{\alpha} \hat{\beta}}, \underline{\mathbf{a}})$ of \mathfrak{G} on $A_{b^* \alpha} \mathcal{L}(H_{\Phi})$ can be considered as well as an action $(j, \underline{\mathbf{a}})$ of $\mathfrak{G}(\mathbf{a})^{oc}$.*

Proof. Left to the reader. \square

14.7. Proposition. *Let us use the notations introduced in 14.1 and 14.4; let us define, for all $x \in M'$, $\mu(x) = 1_{b \otimes_{\alpha} x}$. We define this way an injective $*$ -homomorphism from*

M' into \tilde{M} . Moreover, we have :

(i) for all $x \in M'$, $j_1(\mu(x)) = \mu(R^c(x))$; in particular, for all $n \in N$, $\mu(\hat{\alpha}(n))$ belongs to \tilde{N} and $j_1(\mu(\hat{\alpha}(n))) = 1_{b \otimes_{\alpha} \hat{\beta}(n)}$.

(ii) for all positive $x \in M'^+$, we have $T_{\underline{\mathbf{a}}}(\mu(x)) = \mu(T^{oc}(x))$.

(iii) for all $x \in M'$, we have :

$$\tilde{\Gamma}(\mu(x)) \simeq (\mu * \mu) \Gamma^{oc}(x)$$

Proof. We have, using successively 13.5, 3.11(v) and 13.5(ii) :

$$\begin{aligned} j_1(1_{b \otimes_{\alpha} x}) &= J_{\tilde{\psi}}(1_{b \otimes_{\alpha} x^*}) J_{\tilde{\psi}} \\ &= U_{\tilde{\psi}}^{\mathbf{a}}(J_{\tilde{\psi}} 1_{b \otimes_{\alpha} x^*} J_{\tilde{\Phi}})(1_{b \otimes_{\alpha} x^*})(J_{\tilde{\psi}} 1_{a \otimes_{\beta} J_{\tilde{\Phi}}})(U_{\tilde{\psi}}^{\mathbf{a}})^* \\ &= U_{\tilde{\psi}}^{\mathbf{a}}(1_{a \otimes_{\beta} R^c(x)})(U_{\tilde{\psi}}^{\mathbf{a}})^* \\ &= 1_{b \otimes_{\alpha} R^c(x)} \end{aligned}$$

which gives (i). Result (ii) is just a rewriting of [13.10\(i\)](#).

Let us apply now [14.4](#) and [14.6](#) to $1_{b \otimes_N \alpha} y$, with $y \in \widehat{M}'$; we get :

$$\begin{aligned} 1_{b \otimes_N \alpha} 1_{\hat{\alpha} \otimes_{N^o} \beta} \Gamma^{oc}(y) &= \tilde{\alpha}(1_{b \otimes_N \alpha} 1_{\hat{\alpha} \otimes_{N^o} \beta} y) \\ &= \tilde{\alpha}(1_{b \otimes_N \alpha} 1_{\hat{\alpha} \otimes_{N^o} \beta} y) \\ &= 1_{b \otimes_N \alpha} 1_{id \otimes_j} \tilde{\Gamma}(1_{b \otimes_N \alpha} y) \end{aligned}$$

which, up to the identifications, gives the result. \square

14.8. Example. Let \mathbf{G} be a locally compact quantum group, and let \mathfrak{a} be an action of \mathbf{G} on a von Neumann algebra A ; by applying [6.4](#) and [14.6](#) to \mathfrak{a} , we can construct a measured quantum groupoid $\mathfrak{G}(\mathfrak{a})$, whose underlying von Neumann algebra is the relative commutant $A \otimes \mathcal{L}(H_\Phi) \cap \mathfrak{a}(A)'$, the basis being $B = A \rtimes_{\mathfrak{a}} \mathbf{G} \cap \mathfrak{a}(A)'$; it can be described also by looking at the restriction \mathfrak{b} of the dual action $\tilde{\alpha}$ of $\widehat{\mathbf{G}}^c$ to B and looking at the inclusion of $\mathfrak{b}(B)$ into the crossed product $B \rtimes_{\mathfrak{b}} \widehat{\mathbf{G}}^c$, which is isomorphic to the inclusion of B into $A \otimes \mathcal{L}(H_\Phi) \cap \mathfrak{a}(A)'$. This example will be studied later on in another article, and we need to compare this example with Vainerman's construction made in [\[Va\]](#).

14.9. Remark. Let us recall that an action \mathfrak{a} of a locally compact quantum group \mathbf{G} on a von Neumann algebra A is called outer when $A \rtimes_{\mathfrak{a}} \mathbf{G} \cap \mathfrak{a}(A)' = \mathbb{C}$ (which means that the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathbf{G}$ is irreducible). It was proven in ([\[EN\]](#), [\[E2\]](#)) that any depth 2 irreducible inclusion of factors $M_0 \subset M_1$ generates a locally compact quantum group \mathbf{G} , and an outer action \mathfrak{a} of $\widehat{\mathbf{G}}^c$ on M_1 , such that $M_1^{\mathfrak{a}} = M_0$, and that the crossed product $M_1 \rtimes_{\mathfrak{a}} \widehat{\mathbf{G}}^c$ is isomorphic to the basic construction made from the inclusion $M_0 \subset M_1$; conversely, Vaes proved ([\[V2\]](#)) that, for any action \mathfrak{a} of a locally compact quantum group \mathbf{G} on a von Neumann algebra A , the inclusion $\mathfrak{a}(A) \subset A \rtimes_{\mathfrak{a}} \mathbf{G}$ is depth 2.

Here, we have constructed from this depth 2 inclusion, a measured quantum groupoid $\mathfrak{G}(\mathfrak{a})$; when the action \mathfrak{a} is outer, this measured quantum groupoid is a locally compact quantum group ([\[EN\]](#), [\[E2\]](#)), and we get [\(14.8\)](#) that this locally compact quantum group is \mathbf{G}^{oc} . This result was certainly known by specialists, but we were not able to find a proof anywhere.

As Vaes ([\[V3\]](#)) proved that any locally compact quantum group has an outer action (on some type III_1 factor), this proves that any locally compact quantum group comes from an irreducible depth 2 inclusion of factors, by the construction made in ([\[EN\]](#), [\[E2\]](#)).

Appendix

APPENDIX A. COINVERSE AND SCALING GROUP OF A MEASURED QUANTUM GROUPOID

In this chapter, we are dealing with a Hopf-bimodule $(N, \alpha, \beta, M, \Gamma)$, equipped with a left-invariant operator-valued weight T , and a right-invariant operator-valued weight T' . If ν denotes a normal semi-finite faithful weight on the basis, let Φ (resp. Ψ) be the lifted normal faithful semi-finite weight on M by T (resp. T'). Then, with the additional hypothesis that the two modular automorphism groups associated to the two weight Φ and Ψ commute (we then say that ν is relatively invariant with respect to T and T' (3.7), we can construct a co-inverse, a scaling group and an antipod, using slight generalizations of the constructions made in ([L2],9) for "adapted measured quantum groupoids".

A.1. Lemma. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exist a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M . Let us denote δ the modulus of Ψ with respect to Φ and λ the scaling operator (2.6). We shall use the notations of 2.2.1. Then :*

(i) *let $x \in \mathcal{T}_{\Psi, T'}$ and $n \in \mathbb{N}$ and $y = e_n x$, with the notations of 2.6; then y belongs to $\mathfrak{N}_{\Psi} \cap \mathfrak{N}_{T'}$, is analytical with respect to Ψ , and the operator $\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}$ is bounded, and its closure $\overline{\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}}$ belongs to \mathfrak{N}_{Φ} ; moreover, with the identifications made in 2.6, we have :*

$$\Lambda_{\Phi}(\overline{\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}}) = J_{\Psi}\Lambda_{\Psi}(y)$$

(ii) *let E be the linear space generated by all such elements of the form $\overline{\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}}$, for all $x \in \mathcal{T}_{\Psi, T'}$ and $n \in \mathbb{N}$; then E is a weakly dense subspace of \mathfrak{N}_{Φ} , and, for all $z \in E$, $\Lambda_{\Phi}(z) \in D((H_{\Phi})_{\beta}, \nu^o)$;*
(iii) *the linear set of all products $\langle \Lambda_{\Phi}(z), \Lambda_{\Phi}(z') \rangle_{\beta, \nu^o}$ (for z, z' in E) is a dense subspace of N .*

Proof. As e_n is analytical with respect to Ψ , y belongs to $\mathfrak{N}_{\Psi} \cap \mathfrak{N}_{T'}$, is analytical with respect to Ψ , and $\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}$ is bounded ([V1], 1.2); as δ^{-1} is the modulus of Φ with respect to Ψ , we get that $\overline{\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}}$ belongs to \mathfrak{N}_{Φ} ; we identify $\Lambda_{\Phi}(\overline{\sigma_{-i/2}^{\Psi}(y^*)\delta^{1/2}})$ with $\Lambda_{\Psi}(\sigma_{-i/2}^{\Psi}(y^*)) = J_{\Psi}\Lambda_{\Psi}(y)$, which is (i).

The subspace E contains all elements of the form $\sigma_{-i/2}^{\Psi}(x^*)\delta^{1/2}\overline{\sigma_{-i/2}^{\Psi}(e_n)}$ ($x \in \mathcal{T}_{\Psi, T'}$), and, by density of $\mathcal{T}_{\Psi, T'}$ in M , we get that the closure of E contains all elements of the form $a e_n \delta^{-1/2} \delta^{1/2} \sigma_{-i/2}^{\Psi}(e_n) = a e_n \sigma_{-i/2}^{\Psi}(e_n)$, for all $a \in M$; now, as $e_n \sigma_{-i/2}^{\Psi}(e_n)$ is converging to 1, we finally get that E is dense in M ; as $\Lambda_{\Phi}(E) \subset J_{\Psi}\Lambda_{\Psi}(\mathfrak{N}_{\Psi} \cap \mathfrak{N}_{T'})$, we get, by 2.2, that, for all z in E , $\Lambda_{\Phi}(z)$ belongs to $D((H_{\Phi})_{\beta}, \nu^o)$; more precisely, we have :

$$R^{\beta, \nu^o}(\Lambda_{\Phi}(\overline{\sigma_{-i/2}^{\Psi}(x^*)\delta^{1/2}\sigma_{-i/2}^{\Psi}(e_n)})) = R^{\beta, \nu^o}(J_{\Psi}\Lambda_{\Psi}(e_n x)) = \Lambda_{T'}(e_n x)$$

Therefore, the set of elements of the form $\langle \Lambda_{\Phi}(z), \Lambda_{\Phi}(z') \rangle_{\beta, \nu^o}$ contains all elements of the form $\beta^{-1} \circ T'(x^* e_n e_n x)$, for all x in $\mathcal{T}_{\Psi, T'}$ and $n \in \mathbb{N}$; as we have :

$$T'(x^* e_n e_n x) = \Lambda_{T'}(e_n x)^* \Lambda_{T'}(e_n x) = \Lambda_{T'}(x)^* e_n^* e_n \Lambda_{T'}(x)$$

we get that its closure contains all elements of the form $\beta^{-1} \circ T'(x^* x)$, and, therefore, it contains $\beta^{-1} \circ T'(\mathfrak{M}_{T'}^+)$, which finishes the proof. \square

A.2. Definition. As in ([L2], 9.2), we can define, for all $\lambda \in \mathbb{C}$, a closed operator $\Delta_{\Phi}^{\lambda} \otimes_{N^o} \Delta_{\Phi}^{\lambda}$, with natural values on elementary tensor products; it is possible also to define a unitary antilinear operator $J_{\Phi} \otimes_{N^o} J_{\Phi}$ from $H_{\Phi} \otimes_{N^o} H_{\Phi}$ onto $H_{\Phi} \otimes_N H_{\Phi}$ (whose inverse will be $J_{\Phi} \otimes_N J_{\Phi}$); by composition, we define then a closed antilinear operator $S_{\Phi} \otimes_{N^o} S_{\Phi}$, with natural values on elementary tensor products, whose adjoint will be $F_{\Phi} \otimes_N F_{\Phi}$.

A.3. Proposition. For all a, c in $(\mathfrak{N}_{\Phi} \cap \mathfrak{N}_T)^*(\mathfrak{N}_{\Psi} \cap \mathfrak{N}_{T'})$, b, d in $\mathcal{T}_{\Psi, T'}$ and g, h in E , the following vector :

$$U_{H_{\Phi}}^* \Gamma(g^*) [\Lambda_{\Phi}(h) \otimes_{\nu} (\lambda_{\Lambda_{\Psi}(\sigma_{-i}^{\Psi}(b^*))}^{\beta, \alpha})^* U_{H_{\Psi}}(\Lambda_{\Psi}(a) \otimes_{\nu^o} \Lambda_{\Phi}((cd)^*))]$$

belongs to $D(S_{\Phi} \otimes_{\nu^*} S_{\Phi})$, and the value of $\sigma_{\nu}(S_{\Phi} \otimes_{\nu^*} S_{\Phi})$ on this vector is equal to :

$$U_{H_{\Phi}}^* \Gamma(h^*) [\Lambda_{\Phi}(g) \otimes_{\nu} (\lambda_{\Lambda_{\Psi}(\sigma_{-i}^{\Psi}(d^*))}^{\beta, \alpha})^* U_{H_{\Psi}}(\Lambda_{\Psi}(c) \otimes_{\nu^o} \Lambda_{\Phi}((ab)^*))]$$

Proof. The proof is identical to ([L2], 9.9), thanks to A.1(ii). \square

A.4. Proposition. There exists a closed densely defined anti-linear operator G on H_{Φ} such that the linear span of :

$$(\lambda_{\Lambda_{\Psi}(\sigma_{-i}^{\Psi}(b^*))}^{\beta, \alpha})^* U_{H_{\Psi}}(\Lambda_{\Psi}(a) \otimes_{\nu^o} \Lambda_{\Phi}((cd)^*))$$

with a, c in $(\mathfrak{N}_{\Phi} \cap \mathfrak{N}_T)^*(\mathfrak{N}_{\Psi} \cap \mathfrak{N}_{T'})$, b, d in $\mathcal{T}_{\Psi, T'}$, is a core of G , and we have :

$$G[(\lambda_{\Lambda_{\Psi}(\sigma_{-i}^{\Psi}(b^*))}^{\beta, \alpha})^* U_{H_{\Psi}}(\Lambda_{\Psi}(a) \otimes_{\nu^o} \Lambda_{\Phi}((cd)^*))] = (\lambda_{\Lambda_{\Psi}(\sigma_{-i}^{\Psi}(d^*))}^{\beta, \alpha})^* U_{H_{\Psi}}(\Lambda_{\Psi}(c) \otimes_{\nu^o} \Lambda_{\Phi}((ab)^*))$$

Proof. The proof is identical to ([L2], 9.10), thanks to A.1(iii). \square

A.5. Theorem. Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exist a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M . Let G be the closed densely defined antilinear operator defined in A.4, and let $G = ID^{1/2}$ its polar decomposition. Then, the operator D is positive self-adjoint and non singular; there exists a one-parameter automorphism group τ_t on M defined, for $x \in M$, by :

$$\tau_t(x) = D^{-it} x D^{it}$$

We have, for all $n \in N$ and $t \in \mathbb{R}$:

$$\tau_t(\alpha(n)) = \alpha(\sigma_t^{\nu}(n))$$

$$\tau_t(\beta(n)) = \beta(\sigma_t^{\nu}(n))$$

which allows us to define $\tau_t \beta^* \alpha \tau_t$, $\tau_t \beta^* \alpha \sigma_t^{\Phi}$ and $\sigma_t^{\Psi} \beta^* \alpha \tau_{-t}$ on $M \otimes_{N^o} M$; moreover, we have :

$$\Gamma \circ \tau_t = (\tau_t \beta^* \alpha \tau_t) \Gamma$$

$$\Gamma \circ \sigma_t^{\Phi} = (\tau_t \beta^* \alpha \sigma_t^{\Phi}) \Gamma$$

$$\Gamma \circ \sigma_t^{\Psi} = (\sigma_t^{\Psi} \beta^* \alpha \tau_{-t}) \Gamma$$

Proof. The proof is identical to [L2], 9.12 to 9.28. \square

A.6. Theorem. Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M . Let G be the closed densely defined antilinear operator defined in A.4, and let $G = ID^{1/2}$ its polar decomposition. Then, the operator I is antilinear, isometric, surjective, and we have $I = I^* = I^2$; there exists a $*$ -antiautomorphism R on M defined, for $x \in M$, by :

$$R(x) = Ix^*I$$

such that, for all $t \in \mathbb{R}$, we get $R \circ \tau_t = \tau_t \circ R$ and $R^2 = id$.

For any a, b in $\mathfrak{N}_\Psi \cap \mathfrak{N}_{T'}$ we have :

$$R((\omega_{J_\Psi \Lambda_\Psi(a)} \underset{N}{\beta^* \alpha} id) \Gamma(b^* b)) = (\omega_{J_\Psi \Lambda_\Psi(b)} \underset{N}{\beta^* \alpha} id) \Gamma(a^* a)$$

and for any c, d in $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$, we have :

$$R((id \underset{N}{\beta^* \alpha} \omega_{J_\Phi \Lambda_\Phi(c)}) \Gamma(d^* d)) = (id \underset{N}{\beta^* \alpha} \omega_{J_\Phi \Lambda_\Phi(d)}) \Gamma(c^* c)$$

For all $n \in N$, we have $R(\alpha(n)) = \beta(n)$, which allows us to define $R_{\underset{N}{\beta^* \alpha}} R$ from $M_{\underset{N}{\beta^* \alpha}} M$ onto $M_{\underset{N^o}{\alpha^* \beta}} M$ (whose inverse will be $R_{\underset{N^o}{\alpha^* \beta}} R$), and we have :

$$\Gamma \circ R = \varsigma_{N^o} (R_{\underset{N}{\beta^* \alpha}} R) \Gamma$$

Proof. The proof is identical to [L2], 9.38 to 9.42. \square

A.7. Theorem. Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T_L$; then :

(i) M is the weak closure of the linear span of all elements of the form $(\omega_{\underset{N}{\beta^* \alpha}} id) \Gamma(x)$,

for all $x \in M$ and $\omega \in M_*$ such that there exists $k > 0$ with $\omega \circ \beta \leq k\nu$.

(ii) M is the weak closure of the linear span of all elements of the form $(id \underset{N}{\beta^* \alpha} \omega) \Gamma(x)$,

for all $x \in M$ and $\omega \in M_*$ such that there exists $k > 0$ with $\omega \circ \alpha \leq k\nu$.

(iii) M is the weak closure of the linear span of all elements of the form $(id * \omega_{v,w})(W)$, where v belongs to $D(\alpha H_\Phi, \nu)$ and w belongs to $D((H_\Phi)_\beta, \nu^o)$.

Proof. The proof is identical to [L2], 9.25. \square

A.8. Definition. Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M ; let τ_t the one-parameter automorphism group constructed in A.5 and let R be the involutive $*$ -antiautomorphism constructed in A.6. We shall call τ_t the scaling group of $(N, \alpha, \beta, M, \Gamma, T, T', \nu)$ and R the coinverse of $(N, \alpha, \beta, M, \Gamma, T, T', \nu)$. Thanks to A.6 and A.7, we see that, T and ν being given, R does not depend on the choice of the right-invariant operator-valued weight T' .

Similarly, from A.5, one gets that, for all x in M , $\omega \in M_*$ such that there exists $k > 0$ with $\omega \circ \alpha \leq k\nu$, $\omega' \in M_*$ such that there exists $k > 0$ with $\omega' \circ \beta \leq k\nu$, one has :

$$\tau_t((id_{\beta * \alpha} \omega) \Gamma(x)) = (id_{\beta * \alpha} \omega \circ \sigma_{-t}^\Phi) \Gamma \sigma_t^\Phi(x)$$

$$\tau_t((\omega'_{\beta * \alpha} id) \Gamma(x)) = (\omega' \circ \sigma_t^\Psi \beta * \alpha id) \Gamma \sigma_{-t}^\Psi(x)$$

So, T and ν being given, τ_t does not depend on the choice of the right-invariant operator-valued weight T' .

A.9. Theorem. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$; then, for any ξ, η in $D(\alpha H_\Phi, \nu) \cap D((H_\Phi)_{\hat{\beta}}, \nu^\circ)$, $(id * \omega_{\xi, \eta})(W)$ belongs to $D(\tau_{i/2})$, and, if we define $S = R\tau_{i/2}$, we have :*

$$S((id * \omega_{\xi, \eta})(W)) = (id * \omega_{\eta, \xi})(W)^*$$

More generally, for any x in $D(S) = D(\tau_{i/2})$, we get that $S(x)^*$ belongs to $D(S)$ and $S(S(x)^*)^* = x$; S will be called the antipod of the measured quantum groupoid, and, therefore, the co-inverse and the scaling group, given by polar decomposition of the antipod, rely only upon the pseudo-multiplicative W .

Proof. It is proved similarly to [L2] 9.35 and 9.36. \square

A.10. Proposition. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule, equipped with a left-invariant operator-valued weight T , and a right-invariant valued weight T' ; let ν be a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7; let τ_t be the scaling group of $(N, \alpha, \beta, M, \Gamma, T, T', \nu)$ and R the coinverse of $(N, \alpha, \beta, M, \Gamma, T_L, T_R, \nu)$; then :*

- (i) *the operator-valued weight $RT'R$ is left-invariant, the operator valued-weight RTR is right-invariant, and ν is relatively invariant with respect to $RT'R$ and RTR .*
- (ii) *τ_t is the scaling group of $(N, \alpha, \beta, M, \Gamma, RT'R, RTR, \nu)$*

Proof. Let $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M by T and T' ; the lifted weight by $RT'R$ (resp. RTR) is then $\Psi \circ R$ (resp. $\Phi \circ R$). As $\sigma_t^{\Psi \circ R} = R \circ \sigma_{-t}^\Psi \circ R$ and $\sigma_s^{\Phi \circ R} = R \circ \sigma_{-s}^\Phi \circ R$, we get that $\sigma^{\Psi \circ R}$ and $\sigma^{\Phi \circ R}$ commute, which is (i).

From A.5 and A.6, we get that :

$$\begin{aligned} \Gamma \circ \sigma_t^{\Psi \circ R} &= \Gamma \circ R \circ \sigma_{-t}^\Psi \circ R = \varsigma_{N^\circ}(R_{\beta * \alpha} R) \Gamma \circ \sigma_{-t}^\Psi \circ R \\ &= \varsigma_{N^\circ}(R \circ \sigma_{-t}^\Psi \circ R_{\alpha * \beta} R \circ \tau_t \circ R) \varsigma_N \Gamma = (\tau_t \beta * \alpha \sigma_t^{\Psi \circ R}) \Gamma \end{aligned}$$

from which we get that, for all $x \in M$ and $\omega \in M_*$ such that there exists $k > 0$ such that $\omega \circ \alpha < k\nu$, we have :

$$\tau_t((id_{\beta * \alpha} \omega) \Gamma(x)) = (id_{\beta * \alpha} \omega \circ \sigma_{-t}^{\Psi \circ R}) \Gamma(\sigma_t^{\Psi \circ R}(x))$$

from which we get, by A.7, that τ_t is the scaling group associated to $RT_R R$, $RT_L R$ and ν . \square

In this section, with the same hypothesis as in appendix A, we construct two one-parameter automorphism groups on the basis N (B.2), and we prove (B.7) that these automorphisms leave invariant the quasi-invariant weight ν . We prove also in B.7 that the weight ν is also quasi-invariant with respect to T and RTR . We finish by proving (B.8) that our axioms and Lesieur's axioms are equivalent.

B.1. Lemma. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule, equipped with a left-invariant operator-valued weight T , and a right-invariant valued weight T' ; let ν be a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7. Let $x \in M \cap \alpha(N)'$ and $y \in M \cap \beta(N)'$. Then :*

(i) x belongs to $\beta(N)$ if and only if we have :

$$\Gamma(x) = 1_{\beta \otimes_{\alpha} x}$$

(ii) y belongs to $\alpha(N)$ if and only if we have :

$$\Gamma(y) = y_{\beta \otimes_{\alpha} 1}$$

More generally, if x_1, x_2 are in $M \cap \alpha(N)'$ and such that $\Gamma(x_1) = 1_{\beta \otimes_{\alpha} x_2}$, then $x_1 = x_2 \in \beta(N)$.

Proof. The proof is given in [L2], 4.4. □

B.2. Proposition. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7. Then, there exists a unique one-parameter group of automorphisms γ_t^L of N such that, for all $t \in \mathbb{R}$ and $n \in N$, we have :*

$$\sigma_t^T(\beta(n)) = \beta(\gamma_t^L(n))$$

$$\sigma_t^{RTR}(\alpha(n)) = \alpha(\gamma_{-t}^L(n))$$

Moreover, the automorphism groups γ_t^L and σ_t^ν commute, and there exists a positive self-adjoint non-singular operator $h_L \in Z(N) \cap N^{\gamma^L}$ such that, for any $x \in N^+$ and $t \in \mathbb{R}$, we have :

$$\nu \circ \gamma_t^L(x) = \nu(h_L^t x)$$

Using the operator-valued weights $RT'R$ and RTR , we obtain another one-parameter group of automorphisms γ_t^R of N , such that we have :

$$\sigma_t^{RT'R}(\beta(n)) = \beta(\gamma_t^R(n))$$

$$\sigma_t^{T'}(\alpha(n)) = \alpha(\gamma_{-t}^R(n))$$

and a positive self-adjoint non-singular operator $h_R \in Z(N) \cap N^{\gamma^R}$ such that we have :

$$\nu \circ \gamma_t^R(x) = \nu(h_R^t x)$$

Proof. The existence of γ_t^L is given by [L2], 4.5; moreover, from the formula $\sigma_t^\Phi \circ \sigma_s^\Psi(\beta(n)) = \sigma_s^\Psi \circ \sigma_t^\Phi(\beta(n))$, we obtain :

$$\beta(\gamma_t^L \circ \sigma_{-s}^\nu(n)) = \beta(\sigma_{-s}^\nu \circ \gamma_t^L(n))$$

which gives the commutation of γ_t^L and σ_{-s}^ν . The existence of h_L is then straightforward. The construction of γ^R and h_R is just the application of the preceeding results to $RT'R$, RTR and ν . \square

B.3. Proposition. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7. Let T_L (resp. T_R) be another left (resp. right)-invariant operator-valued weight; we shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$, $\Phi' = \nu \circ \alpha^{-1} \circ T_L$, $\Psi = \nu \circ \beta^{-1} \circ T'$ and $\Psi' = \nu \circ \beta^{-1} \circ T_R$ the lifted normal semi-finite weights on M ; then, we have :*

$$\begin{aligned}\beta(h_L^{ist}) &= (D\Psi' \circ \sigma_t^\Phi : D\Psi' \circ \tau_t)_s \\ \alpha(h_R^{ist}) &= (D\Phi' \circ \sigma_{-t}^\Psi : D\Phi' \circ \tau_t)_s\end{aligned}$$

where τ_s is the scaling group constructed from T , T' and ν as well from $RT'R$, RTR and ν (A.5 and A.10).

Proof. From A.5, we get, for all $t \in \mathbb{R}$, $\Gamma \circ \sigma_t^\Phi \tau_{-t} = (id_{\beta * \alpha} \sigma_t^\Phi \tau_{-t})\Gamma$, and, therefore, by the right-invariance of T_R , we get, for all $x \in \mathfrak{M}_{T_R}^+$, that $\tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) = T_R(x)$; let now $x \in \mathfrak{M}_{\Psi'}^+$; $T_R(x)$ is an element of the positive extended part of $\beta(N)$ which can be written :

$$\int_0^\infty \lambda de_\lambda + (1-p)\infty$$

where p is a projection in $\beta(N)$, and e_λ is a resolution of p . As x belongs to $\mathfrak{M}_{\Psi'}^+$, it is well known that $p = 1$, and $T_R(x) = \int_0^\infty \lambda de_\lambda$. There exists also a projection q and a resolution of q such that :

$$\tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) = \int_0^\infty \lambda df_\lambda + (1-q)\infty$$

and, for all $\mu \in \mathbb{R}^+$, we have, because $e_\mu x e_\mu$ belongs to $\mathfrak{M}_{T_R}^+$:

$$\begin{aligned}e_\mu \left(\int_0^\infty \lambda df_\lambda \right) e_\mu + e_\mu (1-q) e_\mu \infty &= e_\mu \tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) e_\mu \\ &= \tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(e_\mu x e_\mu) \\ &= T_R(e_\mu x e_\mu) \\ &= \int_0^\mu \lambda de_\lambda\end{aligned}$$

from which we infer that $(1-q)e_\mu = 0$, and, therefore, that $q = 1$; then, we get that $e_\mu \tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) e_\mu$ is increasing with μ towards $T_R(x)$. Therefore, we get that :

$$\tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) \subset T_R(x)$$

and, finally, the equality, for all $x \in \mathfrak{M}_{\Psi'}^+$:

$$\tau_t \sigma_{-t}^\Phi T_R \sigma_t^\Phi \tau_{-t}(x) = T_R(x)$$

Moreover, as we have, for all $n \in N$:

$$\tau_t \sigma_{-t}^\Phi(\beta(n)) = \beta(\sigma_t^\nu \gamma_{-t}^L(n))$$

we get, using B.2, that, for all $x \in \mathfrak{M}_{\Psi'}^+$:

$$\Psi'(\beta(h_L^{-t/2}) \sigma_t^\Phi \tau_{-t}(x) \beta(h_L^{-t/2})) = \Psi'(x)$$

and, therefore, that, for all $x \in M^+$:

$$\Psi'(\beta(h_L^{-t/2})\sigma_t^\Phi\tau_{-t}(x)\beta(h_L^{-t/2})) \leq \Psi'(x)$$

A similar calculation (with $\tau_t\sigma_{-t}^\Phi$ instead of $\sigma_t^\Phi\tau_{-t}$) leads to :

$$\Psi'(\beta(h_L^{t/2})\tau_t\sigma_{-t}^\Phi(x)\beta(h_L^{t/2})) \leq \Psi'(x)$$

which leads to the equality, from which we get the first result.

Applying this result to $RT_R R$, $RT_L R$ and ν , we get, using again [A.10](#) :

$$\begin{aligned} \beta(h_R^{ist}) &= (D\Phi' \circ R \circ \sigma_t^{\Psi \circ R} : D\Phi' \circ R \circ \tau_t)_s \\ &= (D\Phi' \circ \sigma_{-t}^\Psi \circ R : D\Phi' \circ \tau_t \circ R)_s \\ &= R[((D\Phi' \circ \sigma_{-t}^\Psi : D\Phi' \circ \tau_t)_{-s})^*] \end{aligned}$$

which leads to the result. \square

B.4. Corollary. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule; let us suppose that there exists a left-invariant operator-valued weight T , a right-invariant valued weight T' and ν a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of [3.7](#). We shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M , R the coinverse and τ_t the scaling group constructed in [A.6](#) and [A.5](#); we shall denote λ the scaling operator of Ψ with respect to Φ ([2.6](#)), h_L and h_R the operators constructed in [B.2](#). Then, for all s, t in \mathbb{R} :*

- (i) $(D\Psi : D\Psi \circ \tau_t)_s = \lambda^{ist}\beta(h_L^{ist})$
- (ii) $(D\Phi : D\Phi \circ \tau_t)_s = \lambda^{ist}\alpha(h_R^{ist})$
- (iii) $(D\Phi : D\Phi \circ \sigma_{-t}^{\Phi \circ R})_s = \lambda^{ist}\alpha(h_R^{ist})\alpha(h_L^{-ist})$
- (iv) $(D\Psi : D\Psi \circ \sigma_t^{\Psi \circ R})_s = \lambda^{ist}\beta(h_L^{ist})\beta(h_R^{-ist})$.

Proof. Applying [B.3](#) with $T_R = T'$, as $(D\Psi \circ \sigma_t^\Phi : D\Psi)_s = \lambda^{-ist}$ ([2.6](#)), we obtain (i). Applying [B.3](#) with $T_L = T$, as $(D\Phi : D\Phi \circ \sigma_{-t}^\Psi)_s = \lambda^{ist}$, we obtain (ii). Applying [B.3](#) with $T_R = RTR$, we obtain :

$$\begin{aligned} \beta(h_L^{ist}) &= (D\Phi \circ R \circ \sigma_t^\Phi : D\Phi \circ R \circ \tau_t)_s \\ &= (D\Phi \circ \sigma_{-t}^{\Phi \circ R} \circ R : D\Phi \circ \tau_t \circ R)_s \\ &= R((D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi \circ \tau_t)_{-s}^*) \end{aligned}$$

and, therefore $\alpha(h_L^{ist}) = (D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi \circ \tau_t)_{-s}^*$ from which one gets :

$$\alpha(h_L^{ist}) = (D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi \circ \tau_t)_s$$

Using (ii), we get :

$$(D\Phi : D\Phi \circ \sigma_{-t}^{\Phi \circ R})_s = \lambda^{ist}\alpha(h_R^{ist})\alpha(h_L^{-ist})$$

which is (iii). And applying [B.3](#) with $T_L = RT'R$, we obtain (iv). \square

B.5. Lemma. *Let M be a von Neumann algebra, Φ a normal semi-finite faithful weight on M , θ_t a one parameter group of automorphisms of M . Let us suppose that there exists a positive non singular operator μ affiliated to M^Φ such that, for all s, t in \mathbb{R} , we have*

$$(D\Phi \circ \theta_t : D\Phi)_s = \mu^{ist}$$

We have then, for all $t \in \mathbb{R}$, $\theta_t(\mu) = \mu$. Let us write $\mu = \int_0^\infty \lambda d\epsilon_\lambda$ the spectral decomposition of μ , and let us define $f_n = \int_{1/n}^n d\epsilon_\lambda$. We have then, for all a in \mathfrak{N}_Φ , t in \mathbb{R} , n in \mathbb{N} :

$$\omega_{J_\Phi \Lambda_\Phi(a f_n)} \circ \theta_t = \omega_{J_\Phi \Lambda_\Phi(\theta_{-t}(a) f_n \mu^{t/2})}$$

Proof. Let us remark first that $\theta_t(\mu) = \mu$, and, therefore, $\theta_t(f_n) = f_n$. On the other hand, for any a in M , we have :

$$\theta_{-t}\sigma_s^\Phi\theta_t(x) = \sigma_s^{\Phi\circ\theta_t}(x) = \mu^{ist}\sigma_s^\Phi(x)\mu^{-ist}$$

and then :

$$\theta_{-t}\sigma_s^\Phi(x) = \mu^{ist}\sigma_s^\Phi\theta_{-t}(x)\mu^{-ist}$$

If now x is analytic with respect to Φ , we get that $\theta_{-t}(f_n x f_m)$ is analytic with respect to Φ and that :

$$f_n\theta_{-t}\sigma_{i/2}^\Phi(x)f_m = \mu^{-t/2}f_n\sigma_{i/2}^\Phi(\theta_{-t}(x))f_m\mu^{t/2}$$

Let us take now a in \mathfrak{N}_Φ , analytic with respect to Φ ; we have, for any y in M :

$$\begin{aligned}\omega_{J_\Phi\Lambda_\Phi(f_n a f_m)} \circ \theta_t(y) &= (\theta_t(y)J_\Phi\Lambda_\Phi(f_n a f_m)|J_\Phi\Lambda_\Phi(f_n a f_m)) \\ &= (\theta_t(y)\Lambda_\Phi(f_m\sigma_{-i/2}^\Phi(a^*)f_n)|\Lambda_\Phi(f_m\sigma_{-i/2}^\Phi(a^*)f_n)) \\ &= \Phi(f_n\sigma_{i/2}^\Phi(a)f_m\theta_t(y)f_m\sigma_{-i/2}^\Phi(a^*)f_n)\end{aligned}$$

which, using the preceeding remarks, is equal to :

$$\Phi \circ \theta_t(\mu^{-t/2}f_n\sigma_{i/2}^\Phi(\theta_{-t}(a))f_m\mu^{t/2}y\mu^{t/2}f_m\sigma_{-i/2}^\Phi(\theta_{-t}(a^*))f_n\mu^{-t/2})$$

and, making now f_n increasing to 1, we get that $\omega_{J_\Phi\Lambda_\Phi(a f_m)} \circ \theta_t(y)$ is equal to :

$$\begin{aligned}\Phi(\sigma_{i/2}^\Phi(\theta_{-t}(a))f_m\mu^{t/2}y\mu^{t/2}f_m\sigma_{-i/2}^\Phi(\theta_{-t}(a^*))) \\ = (y\Lambda_\Phi(f_m\mu^{t/2}\sigma_{-i/2}^\Phi(\theta_{-t}(a^*)))|\Lambda_\Phi(f_m\mu^{t/2}\sigma_{-i/2}^\Phi(\theta_{-t}(a^*))) \\ = (yJ_\Phi\Lambda_\Phi(\theta_{-t}(a)f_m\mu^{t/2})|J_\Phi\Lambda_\Phi(\theta_{-t}(a)f_m\mu^{t/2}))\end{aligned}$$

from which we get the result. \square

B.6. Lemma. Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule, equipped with a left-invariant operator-valued weight T , and a right-invariant valued weight T' ; let ν be a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7. We shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M , R the coinverse and τ_t the scaling group constructed in A.6 and A.5. Then, we have :

(i) there exists a positive non singular operator μ_1 affiliated to M^Φ and invariant under τ_t , such that $(D\Phi \circ \tau_t : D\Phi)_s = \mu_1^{ist}$; let us write $\mu_1 = \int_0^\infty \lambda d\epsilon_\lambda$ and $f_n = \int_{1/n}^n d\epsilon_\lambda$; we have then, for all a in \mathfrak{N}_Φ , t in \mathbb{R} , n in \mathbb{N} and x in M^+ :

$$\omega_{J_\Phi\Lambda_\Phi(\tau_t(a)f_n)} = \omega_{J_\Phi\Lambda_\Phi(a f_n \mu_1^{t/2})} \circ \tau_{-t}$$

$$T \circ \tau_t(x) = \alpha \circ \sigma_t^\nu \circ \alpha^{-1}(T(\mu_1^{t/2} x \mu_1^{-t/2}))$$

(ii) there exists a positive non singular operator μ_2 affiliated to M^Φ and invariant under $\sigma_t^{\Phi \circ R}$, such that $(D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi)_s = \mu_2^{ist}$; let us write $\mu_2 = \int_0^\infty \lambda d\epsilon'_\lambda$ and $f'_n = \int_{1/n}^n d\epsilon'_\lambda$; we have then, for all b in \mathfrak{N}_Φ , t in \mathbb{R} and n in \mathbb{N} :

$$\omega_{J_\Phi\Lambda_\Phi(b f'_n)} \circ \sigma_t^{\Phi \circ R} = \omega_{J_\Phi\Lambda_\Phi(\sigma_{-t}^{\Phi \circ R}(b) f'_n \mu_2^{-t/2})}$$

$$T(\sigma_{-t}^{\Phi \circ R}(\mu_1^{-t/2} x \mu_1^{t/2})) = \alpha \circ \gamma_t^L \circ \alpha^{-1}(T(x))$$

Moreover, we have $\mu_1^{is} = \lambda^{-is} \alpha(h_R^{-is})$, $\mu_2^{is} = \mu_1^{is} \alpha(h_L^{is})$, and μ_1^{is} , μ_2^{is} , $\alpha(h_L^{is})$ belong to $\alpha(N)' \cap M^\Phi$. The non-singular operators μ_1 , μ_2 and $\alpha(h_L)$ commute two by two.

Proof. By B.4(ii), we get that $(D\Phi \circ \tau_t : D\Phi)_s = \lambda^{-ist} \alpha(h_R^{-ist})$, as λ is positive non singular, affiliated to the center $Z(M)$, and h_R is positive non singular affiliated to the center of N , we get there exists μ_1 positive non singular, affiliated to M^Φ such that :

$$\mu_1^{ist} = \lambda^{-ist} \alpha(h_R^{-ist}) = (D\Phi \circ \tau_t : D\Phi)_s$$

We can then apply B.5 to τ_t and $\tau_t(a)f_n$ (which belongs to \mathfrak{N}_Φ) to get the first formula of (i). On the other hand, we get that $\alpha \circ \sigma_{-t}^\nu \circ \alpha^{-1} \circ T \circ \tau_t$ is a normal semi-finite operator-valued weight which verify, for all $x \in M^+$

$$\alpha \circ \sigma_{-t}^\nu \circ \alpha^{-1} \circ T \circ \tau_t(x) = T(\mu_1^{t/2} x \mu_1^{t/2})$$

from which we get the second formula of (i).

By B.4(iii), we get that $(D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi)_s = \lambda^{-ist} \alpha(h_R^{-ist}) \alpha(h_L^{ist})$; with the same arguments, we get that there exists μ_2 positive non singular, affiliated to M^Φ such that:

$$\mu_2^{ist} = \lambda^{-ist} \alpha(h_R^{-ist}) \alpha(h_L^{ist}) = (D\Phi \circ \sigma_{-t}^{\Phi \circ R} : D\Phi)_s$$

and we get the first formula of (ii) by applying again B.5 with $\sigma_{-t}^{\Phi \circ R}$.

On the other hand, using B.2, we get that $\alpha \circ \gamma_{-t}^L \circ \alpha^{-1} \circ T \circ \sigma_{-t}^{\Phi \circ R}$ is an operator-valued weight which verify, for all $x \in M^+$:

$$\begin{aligned} \nu \circ \gamma_{-t}^L \circ \alpha^{-1} \circ T \circ \sigma_{-t}^{\Phi \circ R}(x) &= \nu(h_L^{-t/2} \alpha^{-1}(T \sigma_{-t}^{\Phi \circ R}(x)) h_L^{-t/2}) \\ &= \Phi(\alpha(h_L^{-t/2} \sigma_{-t}^{\Phi \circ R}(x) \alpha(h_L^{-t/2})) \\ &= \Phi \circ \sigma_{-t}^{\Phi \circ R}[\alpha(h_L^{-t/2}) x \alpha(h_L^{-t/2})] \\ &= \Phi(\mu_2^{t/2} \alpha(h_L^{-t/2}) x \alpha(h_L^{-t/2}) \mu_2^{t/2}) \end{aligned}$$

from which we get, because $\mu_2^{t/2} \alpha(h_L^{-t/2})$ commutes with $\alpha(N)$:

$$\alpha \circ \gamma_{-t}^L \circ \alpha^{-1} \circ T \circ \sigma_{-t}^{\Phi \circ R}(x) = T(\mu_2^{t/2} \alpha(h_L^{-t/2}) x \alpha(h_L^{-t/2}) \mu_2^{t/2})$$

or :

$$T(\sigma_{-t}^{\Phi \circ R}(x)) = \alpha \circ \gamma_t^L \circ \alpha^{-1}(T(\mu_1^{t/2} x \mu_1^{t/2}))$$

from which we finish the proof. \square

B.7. Proposition. *Let $(N, \alpha, \beta, M, \Gamma)$ be a Hopf-bimodule, equipped with a left-invariant operator-valued weight T , and a right-invariant valued weight T' ; let ν be a normal semi-finite faithful weight on N , relatively invariant with respect to T and T' in the sense of 3.7. We shall denote $\Phi = \nu \circ \alpha^{-1} \circ T$ and $\Psi = \nu \circ \beta^{-1} \circ T'$ the two lifted normal semi-finite weights on M , R the coinverse and τ_t the scaling group constructed in A.6 and A.5; let λ be the scaling operator of Ψ with respect to Φ (2.6), γ^L and γ^R the two one-parameter automorphism groups of N introduced in B.2; then :*

(i) *for all $t \in \mathbb{R}$, we have $\Gamma \circ \tau_t = (\sigma_t^\Phi \underset{N}{\beta^* \alpha} \sigma_{-t}^{\Phi \circ R}) \Gamma = (\sigma_t^{\Psi \circ R} \underset{N}{\beta^* \alpha} \sigma_{-t}^\Psi) \Gamma$.*

(ii) *we have $h_L = h_R = 1$, and $\nu \circ \gamma^L = \nu \circ \gamma^R = \nu$.*

(iii) *for all s, t in \mathbb{R} , we have $(D\Phi : D\Phi \circ \tau_t)_s = (D\Psi : D\Psi \circ \tau_t)_s = \lambda^{ist}$.*

(iv) *for all s, t in \mathbb{R} , we have $(D\Phi \circ \sigma_t^{\Phi \circ R} : D\Phi)_s = \lambda^{ist}$.*

Therefore, the modular automorphism groups σ^Φ and $\sigma^{\Phi \circ R}$ commute, the weight ν is relatively invariant with respect to Φ and $\Phi \circ R$ and λ is the scaling operator of $\Phi \circ R$ with respect to Φ ; and we have $\tau_t(\lambda) = \lambda$, $R(\lambda) = \lambda$;

(v) *there exists a non singular positive operator q affiliated to $Z(N)$ such that $\lambda = \alpha(q) = \beta(q)$.*

Proof. As, for all $n \in N$, we have :

$$\sigma_{-t}^{\Phi \circ R}(\alpha(n)) = R\sigma_t^\Phi R(\alpha(n)) = \alpha(\gamma_t^L(n))$$

and, by definition, $\sigma_t^\Phi(\beta(n)) = \beta(\gamma_t^L(n))$, using a remark made in 2.4, we may consider the automorphism $\sigma_{-t}^{\Phi \circ R} \beta^*_{\alpha} \sigma_t^{\Phi \circ R}$ on $M \beta^*_{\alpha} M$; let's take a and b in $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$; let's write $h_L = \int_O^\infty \lambda de_\lambda^L$ and let us write $h_p = \int_{1/p}^p de_\lambda^L$; moreover, let's use the notations of B.6; we get that :

$$(id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m)}) (\sigma_{-t}^{\Phi \circ R} \beta^*_{\alpha} \sigma_t^{\Phi \circ R}) \Gamma \circ \tau_t(f_n a^* a f_n)$$

is equal to :

$$\sigma_{-t}^\Phi (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m)} \circ \sigma_t^{\Phi \circ R}) \Gamma \circ \tau_t(f_n a^* a f_n)$$

which, thanks to B.6(ii), can be written, because $\alpha(h_p)$ belongs to $\alpha(N)' \cap M^\Phi$, and therefore $b\alpha(h_p)$ belongs to \mathfrak{N}_Φ :

$$\sigma_{-t}^\Phi (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(\sigma_{-t}^{\Phi \circ R}(b\alpha(h_p))f'_m \mu_2^{-t/2})}) \Gamma \circ \tau_t(f_n a^* a f_n)$$

or :

$$R\sigma_t^{\Phi \circ R} R(id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(\sigma_{-t}^{\Phi \circ R}(b\alpha(h_p))f'_m \mu_2^{-t/2})}) \Gamma \circ \tau_t(f_n a^* a f_n)$$

By B.6 and 2.2.2, we know that $a f_n \mu_1^{t/2}$ belongs to $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$; using now B.6(i), we get that $\tau_t(a f_n) = \tau_t(a) f_n$ belongs to $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$.

On the other hand, by B.6 and 2.2.2, we know that $b\alpha(h_p)f'_m$ belongs to $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$; using now B.6(ii), we get that :

$$\sigma_{-t}^{\Phi \circ R}(b\alpha(h_p)f'_m \mu_1^{-t/2}) = \sigma_{-t}^{\Phi \circ R}(b)f'_m \mu_2^{-t/2} \alpha(h_p) \alpha(h_L^{t/2})$$

belongs to $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$, and so, using again 2.2.2,

$$\sigma_{-t}^{\Phi \circ R}(b)f'_m \mu_2^{-t/2} \alpha(h_p) = \sigma_{-t}^{\Phi \circ R}(b)f'_m \mu_2^{-t/2} \alpha(h_p) \alpha(h_L^{t/2}) \alpha(h_p) \alpha(h_L^{-t/2})$$

belongs also to $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$; therefore, we can use A.6, and we get it is equal to :

$$R\sigma_t^{\Phi \circ R} (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(\tau_t(a)f_n)}) \Gamma(\mu_2^{-t/2} f'_m \alpha(h_p) \sigma_{-t}^{\Phi \circ R}(b^* b) \alpha(h_p) f'_m \mu_2^{-t/2})$$

which can be written, thanks to B.6(i) :

$$R\sigma_t^{\Phi \circ R} (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(a f_n \mu_1^{t/2})} \circ \tau_{-t}) \Gamma(\mu_2^{-t/2} f'_m \alpha(h_p) \sigma_{-t}^{\Phi \circ R}(b^* b) \alpha(h_p) f'_m \mu_2^{-t/2})$$

or, $\alpha(h_p)$, as well as $\mu_2^{-t/2} f'_m$, being invariant under $\sigma_t^{\Phi \circ R}$:

$$R(id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(a f_n \mu_1^{t/2})}) (\sigma_t^{\Phi \circ R} \beta^*_{\alpha} \tau_{-t}) \Gamma \circ \sigma_{-t}^{\Phi \circ R} (\mu_2^{-t/2} f'_m \alpha(h_p) b^* b \alpha(h_p) f'_m \mu_2^{-t/2})$$

and using A.5, and again A.6, we get it is equal to :

$$\begin{aligned} R[(id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(a f_n \mu_1^{t/2})}) \Gamma(\mu_2^{-t/2} f'_m \alpha(h_p) b^* b \alpha(h_p) f'_m \mu_2^{-t/2})] \\ = (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m \mu_2^{-t/2})}) \Gamma(\mu_1^{t/2} f_n a^* a f_n \mu_1^{t/2}) \end{aligned}$$

Finally, we have proved that, for all a, b in $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$, m, n, p in \mathbb{N} , we have :

$$\begin{aligned} (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m)}) (\sigma_{-t}^{\Phi \circ R} \beta^*_{\alpha} \sigma_t^{\Phi \circ R}) \Gamma \circ \tau_t(f_n a^* a f_n) = \\ (id \beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m \mu_2^{-t/2})}) \Gamma(\mu_1^{t/2} f_n a^* a f_n \mu_1^{t/2}) \end{aligned}$$

But, for all $x, y \in M$, we have :

$$\omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m)}(x) = \omega_{J_\Phi \Lambda_\Phi(b)}(\alpha(h_p)f'_m x f'_m \alpha(h_p))$$

$$\omega_{J_\Phi \Lambda_\Phi(b\alpha(h_p)f'_m \mu_2^{-t/2})}(y) = \omega_{J_\Phi \Lambda_\Phi(b)}(\alpha(h_p)f'_m \mu_2^{-t/2} x \mu_2^{-t/2} f'_m \alpha(h_p))$$

and, therefore, we get that :

$$(id_{\beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b)}})[(1_{\beta \otimes_{\alpha} \alpha(h_p)f'_m})(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(f_n a^* a f_n)(1_{\beta \otimes_{\alpha} f'_m \alpha(h_p)})]$$

is equal to :

$$(id_{\beta^*_{\alpha} \omega_{J_\Phi \Lambda_\Phi(b)}})[(1_{\beta \otimes_{\alpha} \alpha(h_p)f'_m \mu_2^{-t/2}})\Gamma(\mu_1^{t/2} f_n a^* a f_n \mu_1^{t/2})(1_{\beta \otimes_{\alpha} \mu_2^{-t/2} f'_m \alpha(h_p)})]$$

and, by density, we get that :

$$(1_{\beta \otimes_{\alpha} \alpha(h_p)f'_m})(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(f_n a^* a f_n)(1_{\beta \otimes_{\alpha} f'_m \alpha(h_p)})$$

is equal to :

$$(1_{\beta \otimes_{\alpha} \alpha(h_p)f'_m \mu_2^{-t/2}})\Gamma(\mu_1^{t/2} f_n a^* a f_n \mu_1^{t/2})(1_{\beta \otimes_{\alpha} \mu_2^{-t/2} f'_m \alpha(h_p)})$$

and, after making p going to ∞ , we obtain that :

$$(1_{\beta \otimes_{\alpha} f'_m})(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(f_n a^* a f_n)(1_{\beta \otimes_{\alpha} f'_m})$$

is equal to (*):

$$(1_{\beta \otimes_{\alpha} f'_m \mu_2^{-t/2}})\Gamma(\mu_1^{t/2} f_n a^* a f_n \mu_1^{t/2})(1_{\beta \otimes_{\alpha} \mu_2^{-t/2} f'_m})$$

Let's now take a file a_i in $\mathfrak{N}_\Phi \cap \mathfrak{N}_T$ weakly converging to 1; going to the limit, we get that :

$$(1_{\beta \otimes_{\alpha} f'_m})(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(f_n)(1_{\beta \otimes_{\alpha} f'_m}) = (1_{\beta \otimes_{\alpha} f'_m \mu_2^{-t/2}})\Gamma(\mu_1^{t/2} f_n \mu_1^{t/2})(1_{\beta \otimes_{\alpha} \mu_2^{-t/2} f'_m})$$

When n goes to ∞ , then f_n is increasing to 1, the first is increasing to $1_{\beta \otimes_{\alpha} f'_m}$, and the

second is increasing to $(1_{\beta \otimes_{\alpha} f'_m \mu_2^{-t/2}})\Gamma(\mu_1^t)(1_{\beta \otimes_{\alpha} \mu_2^{-t/2} f'_m})$ which is therefore bounded.

Taking now m going to ∞ , we get that the two non-singular operators $\Gamma(\mu_1^t)$ and $1_{\beta \otimes_{\alpha} \mu_2^t}$ are equal. Using [B.1](#), we get then that μ_1 is equal to μ_2 (and is affiliated to $\beta(N)$), from which we get, using [B.6](#), that $h_L = 1$. Applying all these calculations to $(N, \alpha, \beta, M, \Gamma, RT'R, T', \nu)$, we get that $h_R = 1$, which is (ii).

Let's come back to the equality (*) above; we obtain that :

$$(1_{\beta \otimes_{\alpha} f'_m})(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(f_n a^* a f_n)(1_{\beta \otimes_{\alpha} f'_m})$$

is equal to :

$$(1_{\beta \otimes_{\alpha} f'_m})\Gamma(f_n a^* a f_n)(1_{\beta \otimes_{\alpha} f'_m})$$

So, when n and m go to ∞ , we obtain :

$$(\sigma_{-t}^{\Phi} \beta^*_{\alpha} \sigma_t^{\Phi \circ R})\Gamma \circ \tau_t(a^* a) = \Gamma(a^* a)$$

which, by density, gives the first formula of (i), the second being given then by [A.10](#).

From (ii) and [B.4](#) (i) and (ii), we get (iii).

From (ii) and [B.4](#)(iii), we get that $(D\Phi \circ \sigma_t^{\Phi \circ R} : D\Phi)_s = \lambda^{ist}$; therefore, as λ is affiliated

to $Z(M)$, we get the commutation of the modular groups σ^Φ and $\sigma^{\Phi \circ R}$. Using 2.6, we get that there exists λ_R positive non singular affiliated to $Z(M)$ and δ_R positive non singular affiliated to M such that $(D\Phi \circ R : D\Phi)_t = \lambda_R^{it^2/2} \delta_R^{it}$, and the properties of R allows us to write that $R(\lambda_R) = \lambda_R$. But, on the other hand, the formula $(D\Phi \circ \sigma_t^{\Phi \circ R} : D\Phi)_s = \lambda_R^{ist}$ (2.6), gives that $\lambda_R = \lambda$ and, therefore, we get that $R(\lambda) = \lambda$. The formula $\tau_t(\lambda) = \lambda$ comes from (iii), which finishes the proof of (iv).

By (i), we have $\lambda = \mu_1 = \mu_2$, and, as we had proved that μ_1 is affiliated to $\beta(N)$, we get that λ is affiliated to $\beta(N)$; as $R(\lambda) = \lambda$ by (iv), we get (v). \square

B.8. Theorem. *Let $(N, M, \alpha, \beta, \Gamma, T, T', \nu)$ be a measured quantum groupoid in the sense of 3.7, and let us denote R (resp. τ_t) the co-inverse (resp. the scaling group) constructed in A.6 (resp. A.5). Then :*

(i) $(N, M, \alpha, \beta, \Gamma, T, RTR, \nu)$ (resp. $(N, M, \alpha, \beta, \Gamma, RT'R, T', \nu)$) is a measured quantum groupoid . Moreover, R (resp. τ_t) remains the co-inverse (resp. the scaling group) of this measured quantum groupoid.

(ii) $(N, M, \alpha, \beta, \Gamma, T, R, \tau, \nu)$ is a measured quantum groupoid in the sense of [L2], 4.1.

(iii) conversely if $(N, M, \alpha, \beta, \Gamma, T, R, \tau, \nu)$ is a measured quantum groupoid in the sense of [L2], 4.1, then $(N, M, \alpha, \beta, \Gamma, T, RTR, \nu)$ is a measured quantum groupoid in the sense of 3.7.

Proof. Result (i) is given by B.7(iv); (ii) is given by A.6, A.8 and B.7(ii); and (iii) is given by [L2], 5.3. \square

REFERENCES

- [AR] C. Anantharaman-Delaroche and J. Renault : Amenable groupoids, Monographies de l'Enseignement Mathématique, **36** (L'enseignement mathématique, Genève, 2000) [19](#)
- [BS] S. Baaĵ and G. Skandalis : Unitaires multiplicatifs et dualité pour les produits croisés de \mathbb{C}^* -algèbres, *Ann. Sci. ENS*, **26** (1993), 425-488. [6](#)
- [BSz1] G. Böhm and K. Szlachányi : A Coassociative \mathbb{C}^* -Quantum group with Non Integral Dimensions, *Lett. Math. Phys.*, **38** (1996), 437-456. [6](#), [19](#)
- [BSz2] G. Böhm and K. Szlachányi : Weak \mathbb{C}^* -Hopf Algebras : the coassociative symmetry of non-integral dimensions, in Quantum Groups and Quantum spaces *Banach Center Publications*, **40** (1997), 9-19. [6](#), [19](#)
- [C1] A. Connes : On the spatial theory of von Neumann algebras, *J. Funct. Analysis*, **35** (1980), 153-164. [2](#), [8](#), [9](#), [12](#)
- [C2] A. Connes : Non commutative Geometry, Academic Press, 1994 [19](#)
- [E1] M. Enock : Produit croisé d'une algèbre de von Neumann par une algèbre de Kac, *J. Funct. Analysis*, **26** (1977), 16-46 [6](#), [49](#)
- [E2] M. Enock : Inclusions irréductibles de facteurs et unitaires multiplicatifs II, *J. Funct. Analysis*, **137** (1996), 466-543. [6](#), [95](#)
- [E3] M. Enock : Quantum groupoids of compact type, *J. Inst. Math. Jussieu*, **4** (2005), 29-133. [6](#), [13](#), [21](#), [22](#), [25](#), [37](#)
- [E4] M. Enock : Inclusions of von Neumann algebras and quantum groupoids III, *J. Funct. Analysis*, **223** (2005), 311-364. [2](#), [7](#), [11](#), [16](#), [17](#), [29](#), [30](#), [91](#)
- [E5] M. Enock : Continuous fields of \mathbb{C}^* -algebras and Lesieur's quantum groupoids, math.OA/0605512 [28](#), [44](#)
- [E6] M. Enock : On Lesieur's measured quantum groupoids, math.OA/07061472 [2](#), [6](#), [7](#), [23](#)
- [EN] M. Enock, R. Nest : Inclusions of factors, multiplicative unitaries and Kac algebras, *J. Funct. Analysis*, **137** (1996), 466-543. [6](#), [8](#), [10](#), [11](#), [14](#), [29](#), [95](#)
- [ES1] M. Enock, J.-M. Schwartz : Produit croisé d'une algèbre de von Neumann par une algèbre de Kac II, *Publ. RIMS Kyoto*, **16** (1980), 189-232. [6](#), [50](#), [73](#), [80](#)
- [ES2] M. Enock, J.-M. Schwartz : Kac Algebras and Duality of locally compact Groups, Springer-Verlag, Berlin, 1989. [6](#), [30](#)
- [EV] M. Enock, J.-M. Vallin : Inclusions of von Neumann algebras and quantum groupoids, *J. Funct. Analysis*, **172** (2000), 249-300. [2](#), [6](#), [7](#), [15](#), [16](#), [18](#), [19](#), [20](#), [21](#), [29](#), [30](#), [46](#), [48](#), [63](#), [91](#)
- [H] U. Haagerup : Operator valued weights in von Neumann algebras I, *J. Funct. Analysis*, **32** (1979), 175-206. [60](#), [69](#)
- [J] V. Jones : Index for subfactors, *Invent. Math.*, **72** (1983), 1-25. [10](#)
- [KV1] J. Kustermans and S. Vaes : Locally compact quantum groups, *Ann. Sci. ENS*, **33** (2000), 837-934. [6](#), [29](#)
- [KV2] J. Kustermans and S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting, *Math. Scand.*, **92** (2003), 68-92. [6](#), [7](#), [29](#)
- [L1] F. Lesieur : thesis, University of Orleans, available at : <http://tel.ccsd.cnrs.fr/documents/archives0/00/00/55/05> [2](#), [6](#), [22](#), [23](#)
- [L2] F. Lesieur : Measured Quantum Groupoids, math.OA/0504104, to be published in *Mémoires de la SMF* [2](#), [6](#), [7](#), [22](#), [23](#), [25](#), [27](#), [28](#), [96](#), [97](#), [98](#), [99](#), [100](#), [107](#)
- [MN] T. Masuda and Y. Nakagami : A von Neumann Algebra framework for the duality of the quantum groups, *Publ. RIMS Kyoto*, **30** (1994), 799-850. [6](#)
- [MNW] T. Masuda, Y. Nakagami and S.L. Woronowicz : A \mathbb{C}^* -algebraic framework for quantum groups, *Internat. J. math.*, **14** (2003), 903-1001. [6](#)
- [NV1] D. Nikshych, L. Vainerman : Algebraic versions of a finite dimensional quantum groupoid, in Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, 2000. [6](#)
- [NV2] D. Nikshych, L. Vainerman : A characterization of depth 2 subfactors of II_1 factors, *J. Funct. Analysis*, **171** (2000), 278-307. [6](#)
- [OR] K. Oty and A. Ramsay : Actions and coactions of measured groupoids on W^* -algebras, *J. Operator Theory*, **56** (2006), 199-217. [46](#)

- [P] A. Paterson : Groupoids, Inverse semi-groups, and their Operator algebras, Birkhauser, Boston, 1999. [19](#)
- [Ra] A. Ramsay : Virtual groups and group actions, *Adv. Math.*, **6** (1971), 253-232. [19](#)
- [R1] J. Renault : A Groupoid Approach to \mathbb{C}^* -Algebras, *Lecture Notes in Math.* **793**, Springer-Verlag [19](#), [22](#)
- [R2] J. Renault : The Fourier algebra of a measured groupoid and its multipliers, *J. Funct. Analysis*, **145** (1997), 455-490. [19](#)
- [S1] J.-L. Sauvageot : Produit tensoriel de Z -modules et applications, in Operator Algebras and their Connections with Topology and Ergodic Theory, Proceedings Buşteni, Romania, 1983, *Lecture Notes in Math.* **1132**, Springer-Verlag, 468-485. [8](#), [9](#), [12](#), [16](#)
- [S2] J.-L. Sauvageot : Sur le produit tensoriel relatif d'espaces de Hilbert, *J. Operator Theory*, **9** (1983), 237-352. [2](#), [8](#), [12](#), [13](#), [14](#), [15](#)
- [St] Ş. Strătilă : Modular theory in Operator Algebras, Abacus Press, Turnbridge Wells, England, 1981
- [Sz] K. Szlachányi : Weak Hopf algebras, in Operators Algebras and Quantum Field Theory, S. Doplicher, R. Longo, J.E. Roberts, L. Zsidó editors, International Press, 1996. [6](#), [19](#)
- [T] M. Takesaki : Theory of Operator Algebras II, Springer, Berlin, 2003. [2](#), [8](#), [10](#), [12](#)
- [V1] S. Vaes : A Radon-Nikodym theorem for von Neumann algebras, *J. Operator Theory*, **46** (2001), 477-489. [2](#), [15](#), [18](#), [96](#)
- [V2] S. Vaes : The unitary implementation of a locally compact Quantum Group action; *J. Funct. Analysis*, **180** (2001), 426-480. [6](#), [30](#), [46](#), [68](#), [95](#)
- [V3] S. Vaes : Strictly outer actions of groups and quantum groups, *J. reine angew. Math.*, **578** (2005), 147-184. [95](#)
- [Va] Leonid Vainerman : A note on quantum groupoids, *C.R. Acad. Sci. Paris*, **315** (1992), 1125-1132. [95](#)
- [Val1] J.-M. Vallin : \mathbb{C}^* -algèbres de Hopf et \mathbb{C}^* -algèbres de Kac, *Proc. London Math. Soc.*, **50** (1985), 131-174. [13](#)
- [Val2] J.-M. Vallin : Bimodules de Hopf et Poids opératoriels de Haar, *J. Operator theory*, **35** (1996), 39-65 [6](#), [18](#), [19](#), [22](#)
- [Val3] J.-M. Vallin : Unitaire pseudo-multiplicatif associé à un group-oïde; applications à la moyennabilité, *J. Operator theory*, **44** (2000), 347-368. [6](#), [22](#)
- [Val4] J.-M. Vallin : Groupoïdes quantiques finis, *J. Algebra*, **239** (2001), 215-261. [6](#)
- [Val5] J.-M. Vallin : Multiplicative partial isometries and finite quantum groupoids, in Locally Compact Quantum Groups and Groupoids, IRMA Lectures in Mathematics and Theoretical Physics 2, V. Turaev, L. Vainerman editors, de Gruyter, 2002. [6](#)
- [W1] S.L. Woronowicz : Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ group. *Invent. Math.*, **93** (1988), 35-76. [6](#)
- [W2] S.L. Woronowicz : Compact quantum group, in "Symétries quantiques" (Les Houches, 1995), North-Holland, Amsterdam (1998), 845-884. [6](#)
- [W3] S.L. Woronowicz : From multiplicative unitaries to quantum groups, *Int. J. Math.*, **7** (1996), 127-149. [6](#)
- [Y1] T. Yamanouchi : Crossed product by groupoid actions and their smooth flows of weights, *Publ. RIMS*, **28** (1992), 535-578. [6](#), [19](#), [63](#), [64](#), [85](#)
- [Y2] T. Yamanouchi : Dual weights on crossed products by groupoid actions, *Publ. RIMS*, bf 28 (1992), 653-678. [6](#), [19](#), [63](#), [64](#), [85](#), [86](#)
- [Y3] T. Yamanouchi : Duality for actions and coactions of measured Groupoids on von Neumann Algebras, *Memoirs of the A.M.S.*, **101** (1993), 1-109. [6](#), [19](#), [22](#), [28](#), [46](#), [63](#), [64](#), [85](#)

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNITÉ MIXTE PARIS 6 / PARIS 7 / CNRS DE RECHERCHE 7586, 175, RUE DU CHEVALERET, PLATEAU 7E, F-75013 PARIS
E-mail address: enock@math.jussieu.fr